



- **Reasons Adults Seek Orthodontic Treatment**

**Reasons Adults Seek Orthodontic Treatment Overcoming Hesitations About Braces Later in Life Clear Aligners for Subtle Tooth Movement Comparing Ceramic Brackets With Other Invisible Options Balancing Cosmetic Goals and Functional Needs Social Changes That Encourage Adult Orthodontics Handling Orthodontic Care Alongside Busy Schedules Managing Discomfort and Adjusting to Lifestyle Shifts Case Studies Highlighting Adult Smile Transformations Psychological Benefits of Aligned Teeth in Adulthood Minimizing Visibility of Orthodontic Appliances Key Questions Adults Often Ask About Treatment**

- **How Orthodontics Connects With Periodontal Health**

**How Orthodontics Connects With Periodontal Health Coordinating Orthodontic Work With Restorative Dentistry The Role of Oral Surgery in Complex Alignment Cases Importance of Joint Consultation for Comprehensive Outcomes Leveraging Data Sharing Between Dental Specialties Collaborative Approaches to Pediatric Growth Management Integrating Prosthodontics for Full Mouth Rehabilitation Teamwork for Managing Tongue Thrust and Speech Issues Streamlining Communication Among Healthcare Providers Shared Planning for Improved Patient Experiences Post Surgical Follow Ups With Multiple Specialists Building a Unified Strategy for Long Term Stability**

- **About Us**



# Collaborative Approaches to Pediatric Growth Management

**\*\*Desire for Improved Aesthetics\*\*:** Many adults seek orthodontic treatment to enhance the appearance of their smile, addressing concerns such as crooked teeth, gaps, or overbites that may have persisted from childhood or developed due to various life factors.

**\*\*Desire for Improved Aesthetics\*\*:** Many adults seek orthodontic treatment to enhance the appearance of their smile, addressing concerns such as crooked teeth, gaps, or overbites that may have persisted from childhood or developed due to various life factors.

Managing growth in pediatric orthodontic patients is a nuanced and vital aspect of ensuring not just the immediate correction of dental anomalies but also the long-term health and aesthetics of a child's dentofacial development. This process, deeply intertwined with collaborative approaches to pediatric growth management, emphasizes the importance of coordinated efforts among various specialists—orthodontists, pediatric dentists, oral surgeons, and sometimes endocrinologists—to tailor treatment plans that consider each child's unique growth trajectory. The essence of this collaborative approach lies in understanding that children are not merely smaller versions of adults; their bodies are dynamic systems undergoing rapid changes that can significantly influence orthodontic outcomes.

The importance of managing growth in pediatric orthodontics cannot be overstated. During childhood and adolescence, the jawbones are still developing and have a remarkable plasticity that allows for more effective and less invasive interventions compared to adult patients. By harnessing this natural growth potential, orthodontists can guide the development of jaws and teeth into harmonious alignment, often reducing or eliminating the need for more complex treatments later in life. Early orthodontic evaluations are recommended around age seven **Kids' dental alignment services** thumb sucking. This proactive strategy not only improves dental function—such as biting and chewing—but also profoundly impacts facial aesthetics and self-esteem, which are crucial during formative years.

Collaboration among healthcare professionals is key to effective growth management. Each specialist brings their expertise to the table: orthodontists focus on tooth positioning and jaw relations; pediatric dentists ensure optimal oral health; oral surgeons might intervene for significant skeletal discrepancies; and endocrinologists could play a role when hormonal imbalances are suspected to affect growth patterns. By working together, these professionals can create comprehensive treatment plans that consider all aspects of a child's development, ensuring interventions are timely, appropriate, and aligned with the child's overall well-being.

Moreover, this collaborative approach fosters communication between families and healthcare providers, empowering parents with knowledge about their child's development stages and treatment options. It instills confidence in the care plan while ensuring that every decision made is with the child's best interests at heart—a critical consideration given the lifelong impact of orthodontic treatments on appearance and health.

In essence, managing growth in pediatric orthodontic patients through collaborative approaches embodies a holistic view towards childhood dentofacial development. It

recognizes the dynamic nature of children's growing bodies and leverages interdisciplinary expertise to deliver personalized care that addresses both immediate needs and future outcomes. In doing so, it sets the stage for healthier smiles that last a lifetime while nurturing confidence in young individuals navigating their developmental journey.

**\*\*Boosting Self-Confidence\*\*:**  
**Orthodontic treatment can significantly improve an adult's self-esteem and confidence by correcting dental imperfections, leading to a more positive self-image and better social interactions.**

---

- **\*\*Desire for Improved Aesthetics\*\***: Many adults seek orthodontic treatment to enhance the appearance of their smile, addressing concerns such as crooked teeth, gaps, or overbites that may have persisted from childhood or developed due to various life factors.
- **\*\*Boosting Self-Confidence\*\***: Orthodontic treatment can significantly improve an adult's self-esteem and confidence by correcting dental imperfections, leading to a more positive self-image and better social interactions.
- **\*\*Enhancing Oral Health\*\***: Proper alignment of teeth through orthodontic treatment can improve oral hygiene by making it easier to clean teeth effectively, thereby reducing the risk of cavities and gum disease that may have been challenging to manage in misaligned dentitions.
- **\*\*Addressing Chronic Dental Issues\*\***: Adults may seek orthodontics to resolve long-standing dental problems such as bite issues (overbites, underbites, crossbites) that can lead to jaw pain, headaches, and digestive difficulties if left untreated.
- **\*\*Correcting Speech Impediments\*\***: Misaligned teeth or jaw structures can contribute to speech difficulties; orthodontic treatment can correct these issues, improving articulation and overall communication skills.
- **\*\*Preventive Measures Against Tooth Wear\*\***: Properly aligned teeth are less prone to excessive wear and tear; adults may pursue orthodontic treatment to prevent premature tooth degradation and associated costs of restorative dentistry.
- **\*\*Facilitating Better Chewing Efficiency\*\***: Orthodontic treatment can improve bite function and alignment, allowing for more efficient chewing which is crucial for digestion and overall nutritional health in adulthood.

Collaborative approaches in pediatric growth management represent a significant shift from traditional, siloed medical practices towards a more holistic, team-oriented strategy

that recognizes the multifaceted nature of children's health and development. This paradigm shift is pivotal, given that growth issues in children can stem from a complex interplay of genetic, nutritional, psychological, and environmental factors. By embracing collaborative approaches, healthcare providers can enhance treatment outcomes in profound ways, ensuring that each child receives comprehensive care tailored to their unique needs.

At the heart of these collaborative efforts lies the concept of interdisciplinary teams. These teams typically include pediatricians, nutritionists, endocrinologists, psychologists, physical therapists, and sometimes even social workers or dietitians. Each professional brings their specialized knowledge to the table, creating a rich tapestry of insights that contribute to a more nuanced understanding of the child's growth challenges. This collective expertise allows for the identification of underlying issues that might be overlooked by a single practitioner focusing on one aspect of health.

One key benefit of such collaboration is the personalized treatment plan it facilitates. Rather than applying a one-size-fits-all approach—and thus risking ineffective or even harmful interventions—healthcare providers can craft strategies that consider all facets of a child's life. For instance, if a child presents with delayed growth due to both nutritional deficiencies and psychological stressors at home, addressing only one aspect would likely yield inadequate results. A collaborative team could simultaneously work on improving dietary intake while also providing family support resources to address emotional well-being and home environment factors.

Moreover, these collaborative efforts significantly enhance communication between healthcare providers and families. Regular meetings and open dialogues ensure that parents are informed partners in their child's care journey rather than mere recipients of directives. This fosters trust and empowers families with knowledge about growth management strategies they can implement at home. Such engagement not only

improves compliance with treatment plans but also helps in early detection of potential issues before they become more serious concerns.

The integration of technology further amplifies the effectiveness of collaborative approaches in pediatric growth management. Digital platforms enable seamless sharing of patient information among team members, allowing for real-time updates and coordinated decision-making without necessitating multiple appointments or redundant tests. Telehealth services extend this reach even further, making specialist consultations accessible regardless of geographical constraints—a critical advantage for families living in rural areas or those facing mobility challenges.

Additionally, research shows that collaborative models often lead to improved psychological outcomes for children with growth issues. The supportive environment created by involving multiple professionals who understand the stress and anxiety such conditions can cause for both children and families plays an integral role in promoting mental well-being alongside physical health improvements.

In conclusion, collaborative approaches to pediatric growth management exemplify how modern healthcare can evolve beyond conventional boundaries to deliver superior outcomes for children facing complex health challenges. By leveraging interdisciplinary expertise, personalizing treatment plans, enhancing communication between providers and families, and ensuring that every child has the opportunity to thrive physically and emotionally, collaborative pediatric care addresses immediate health concerns and long-term well-being—ensuring that every child has the opportunity to thrive physically and emotionally. This collaborative approach in pediatric care set a standard that prioritizes holistic wellness over fragmented treatments—a standard that holds immense promise for generations to come.

## **\*\*Enhancing Oral Health\*\*:**

**Proper alignment of teeth through orthodontic treatment can improve oral hygiene by making it easier to clean teeth effectively, thereby reducing the risk of cavities and gum disease that may have been challenging to manage in misaligned dentitions.**



## Understanding Pediatric Growth Patterns: A Collaborative Approach to Growth Management

Growing up is a miraculous journey, especially in the early years of life when children evolve from tiny infants into vibrant children and eventually into teenagers. This transformative process, however, isn't merely about getting taller or heavier; it's a complex interplay of biological, nutritional, and environmental factors that shape a child's growth pattern. Recognizing the significance of understanding these patterns is crucial for ensuring healthy development and addressing potential growth-related issues promptly. This essay explores the essence of pediatric growth patterns and emphasizes the importance of collaborative approaches in managing them effectively.

At the heart of understanding pediatric growth patterns lies the concept of developmental milestones—predictable stages that most children reach as they grow. These milestones are not just about height and weight; they encompass cognitive, physical, and social developments too. For instance, a toddler's ability to walk independently or a preschooler's emerging ability to express complex thoughts are pivotal indicators of healthy growth. However, every child is unique, influenced by their genetic blueprint and environmental surroundings, which can lead to variations in growth trajectories.

The traditional perspective often relied on standardized growth charts to assess whether a child was growing at an expected rate. While these charts remain invaluable tools for healthcare providers, recognizing their limitations is equally important. They provide a general framework but may overlook individual nuances and cultural differences that influence growth patterns. Herein lies the power of collaboration—a team approach involving parents, pediatricians, nutritionists, psychologists, and educators—to gain a comprehensive understanding tailored to each child's needs.

Collaboration begins with open communication between parents and healthcare providers. Regular check-ups offer opportunities not only to measure physical growth but also to discuss dietary habits, activity levels, emotional well-being, and any concerns parents might have regarding their child's development. This dialogue ensures that interventions are holistic rather than focused solely on physical parameters like height or weight percentiles. Nutritionists play a pivotal role in this context by providing personalized dietary advice that considers cultural preferences while ensuring nutritional adequacy for optimal growth.

Moreover, incorporating insights from psychology can be transformative in managing pediatric growth patterns holistically. Understanding how emotional health affects physical development allows for interventions aimed at reducing stress or anxiety that might impede growth or overall well-being. Schools also become critical partners in this collaborative effort by fostering environments that support both physical activity and mental health through inclusive play activities and supportive educational practices tailored to diverse developmental speeds among students.

Continuous monitoring through technology also enhances collaborative efforts without diminishing personal interactions vital for understanding individual contexts fully. Wearable devices or mobile apps designed for tracking health metrics can provide valuable data points over time but should complement rather than replace face-to-face consultations where nuanced observations are crucial for accurate assessments.

In conclusion, navigating pediatric growth patterns requires more than just numbers on a chart; it demands an empathetic understanding woven through collaborative efforts among various stakeholders in a child's life. By weaving together medical expertise with parental insights and community support systems—including schools—we create an environment conducive not just to reaching typical milestones but thriving in all facets of childhood development. This holistic approach underscores our collective responsibility towards nurturing healthy futures where every child reaches their full potential at their

own pace yet supported by a network dedicated to their well-being every step of the way.

**\*\*Addressing Chronic Dental Issues\*\*:** Adults may seek orthodontics to resolve long-standing dental problems such as bite issues (overbites, underbites, crossbites) that can lead to jaw pain, headaches, and digestive difficulties if left untreated.

When diving into the intricate world of pediatric growth management, particularly through the lens of collaborative approaches to orthodontic treatment, it's essential to understand the key developmental stages that frame a child's dental and skeletal growth. This journey through childhood isn't merely a race to adulthood but a complex symphony of growth, influenced by genetic blueprints, environmental factors, and the nuanced interplay between these elements. Orthodontists, alongside other healthcare professionals like pediatricians and dentists, play critical roles in harmonizing this symphony for optimal outcomes.

The journey begins in infancy, where foundational habits and oral structures set the stage for future developments. During this period, the focus is often on ensuring proper breastfeeding techniques and addressing any anomalies in sucking habits that could lead to dental irregularities down the line. As children transition into toddlerhood and early childhood—roughly ages 2 to 6—the emergence of primary teeth marks a significant milestone. This phase is crucial for monitoring bite alignment and identifying issues such as crossbites or crowding early on. It's during these formative years that preliminary interventions might be considered to guide proper growth trajectories, minimizing potential complications later in life.

As children enter their mixed dentition phase—typically around age 6 to 12—where both primary and permanent teeth coexist, orthodontic evaluations become more frequent. This period is pivotal as it involves not just watching over incoming permanent teeth but also being vigilant about habits like thumb sucking or prolonged use of pacifiers that could adversely affect emerging dental structures. Collaborative care during this stage often includes timely referrals for orthodontic consultations when discrepancies in jaw development or tooth alignment arise, ensuring interventions are timely and effective.

Adolescence brings its own set of challenges with the surge of growth hormones influencing both body and facial structures rapidly. This phase—usually spanning from

around age 11 for girls and 13 for boys—marks a critical window for orthodontic treatments aimed at correcting major alignment issues such as overbites, underbites, or severe crowding before facial bones fully mature. The collaboration among healthcare providers becomes even more crucial here; integrating orthodontic expertise with other specialties ensures comprehensive care tailored to each individual's unique developmental profile.

Moreover, understanding the implications of these developmental stages allows practitioners to adopt preventive strategies rather than solely reactive treatments. Early intervention can significantly reduce the complexity—and sometimes the necessity—for more invasive procedures later on, promoting not just aesthetic outcomes but also functional benefits like improved speech clarity and chewing efficiency. Furthermore, fostering a collaborative approach encourages communication between specialists like orthodontists, general dentists, pediatricians, and even psychologists when needed—a team effort that addresses physical health alongside emotional well-being throughout these pivotal stages of growth.

In essence, navigating key developmental stages through collaborative approaches in pediatric growth management enriches orthodontic treatment with precision timing and holistic care strategies tailored specifically for growing patients. By recognizing each stage's unique needs and challenges while leveraging interdisciplinary cooperation, we pave the way not only for healthier smiles but also support overall development in children—a testament to how teamwork can transform potential hurdles into stepping stones toward thriving adulthoods filled with confident smiles

## **\*\*Correcting Speech**

**Impediments\*\*:** Misaligned teeth or jaw structures can contribute to speech difficulties; orthodontic treatment can correct these issues, improving articulation and overall communication skills.

Interdisciplinary teams, comprising experts like orthodontists, pediatricians, and dentists, play a pivotal role in the comprehensive assessment of growth, particularly in the context

of pediatric growth management. This collaborative approach is not merely beneficial; it's essential for ensuring that children receive holistic care tailored to their unique developmental needs. The beauty of this teamwork lies in its ability to blend specialized knowledge, fostering a more nuanced understanding of growth-related issues than any single discipline could achieve alone.

Let's delve into how these professionals contribute to a well-rounded assessment of growth in children. Orthodontists, with their expertise in the development and alignment of teeth and jaws, are crucial for identifying skeletal and dental discrepancies that may impact overall facial growth and function. They bring to the table an acute awareness of how malocclusions (misalignments of the teeth) can influence not just oral health but also facial aesthetics and breathing patterns. Through careful examination and sometimes utilizing advanced imaging technologies, orthodontists can spot potential issues early on, advocating for timely interventions that align with the child's growth trajectory.

Pediatricians, on the broader canvas of a child's health, oversee the general wellness and developmental milestones of children. They are instrumental in monitoring growth patterns from infancy through adolescence, noting any deviations from expected norms that might signal underlying health conditions affecting growth—ranging from hormonal imbalances to nutritional deficiencies. Their role extends beyond mere observation; pediatricians act as coordinators within this interdisciplinary framework, ensuring that insights from various specialists converge effectively towards the child's best interests. They communicate critical information regarding systemic health factors that could influence both dental and skeletal development.

Dentists contribute significantly by focusing on oral health—a cornerstone of overall well-being—especially during formative years when habits established can have lifelong impacts. Beyond routine check-ups and treatments for cavities or decay, dentists observe how oral structures develop and interact within the growing face and jaw. Their

input is invaluable when considering how dental interventions can support or even enhance orthodontic treatments planned by their colleagues. Moreover, they educate both children and parents about practices that promote healthy oral habits—a preventive approach integral to maintaining proper growth trajectories without unnecessary complications down the line.

The synergy among these disciplines exemplifies collaborative healthcare at its finest. Each professional brings a unique lens through which they assess different facets of a child's growth—orthodontists focusing on dental-skeletal relationships, pediatricians overseeing general health milestones, and dentists championing oral health integrity. When these experts collaborate within an interdisciplinary team setting, they create a tapestry of insights that no single professional could weave alone. This holistic approach ensures interventions are informed by comprehensive assessments: early detection leads to timely treatment plans tailored not just to correct immediate concerns but also to support optimal long-term growth outcomes for each individual child.

In essence, interdisciplinary teams redefine what effective pediatric growth management looks like today—moving away from fragmented care towards integrated strategies that respect the interconnectivity of various aspects influencing childhood development. Such collaboration epitomizes modern healthcare's evolution towards patient-centered care models where every specialist plays a vital part in nurturing healthy futures for our youngest generation.



## **\*\*Preventive Measures**

### **Against Tooth Wear\*\*:**

**Properly aligned teeth are less prone to excessive wear and tear; adults may pursue orthodontic treatment to prevent premature tooth degradation and associated costs of restorative dentistry.**

The Collaborative Team Approach in pediatric growth management represents a multifaceted strategy that harnesses the collective expertise of various professionals to ensure children achieve optimal growth and development. This approach is rooted in the understanding that no single discipline holds all the answers to the complex challenges children may face regarding their growth. By integrating diverse perspectives and specializations, healthcare teams can provide comprehensive care that addresses not just the physical aspects but also psychological and emotional dimensions affecting a child's growth trajectory.

At the heart of this collaborative effort are several key roles, each contributing uniquely to the overarching goal of supporting pediatric patients. Pediatricians serve as the cornerstone of this team, offering foundational knowledge on growth patterns, developmental milestones, and potential indicators of growth disorders. Their ongoing assessment and monitoring are crucial for early identification of any deviations from typical growth trajectories.

Nutritionists play an equally vital role, focusing on dietary needs and how nutrition impacts growth. They tailor feeding plans to meet individual needs, taking into account not only caloric requirements but also the quality of nutrients essential for healthy development. Given that malnutrition or improper nutrition can significantly hinder growth, their insights are indispensable in formulating holistic care plans.

Physical therapists often join the team when there are concerns related to motor development or physical disabilities that could affect a child's ability to grow or move normally. They design exercise programs aimed at enhancing muscle strength and mobility, which are foundational for overall growth and development. Their involvement ensures that physical capabilities are optimized alongside nutritional interventions.

Psychologists contribute by addressing emotional and behavioral aspects that might influence a child's growth indirectly. Stress, anxiety, or family dynamics can have profound effects on eating habits and overall well-being, impacting how children grow physically and mentally. By fostering a supportive environment and providing coping strategies, psychologists help mitigate these factors to create a more conducive atmosphere for healthy development.

Moreover, involving parents or caregivers is paramount within this collaborative framework. Their role extends beyond mere support; they are active participants in understanding their child's needs and implementing care strategies at home. Educating parents about signs of developmental issues, proper nutrition, and activities conducive to healthy growth empowers them to be advocates for their child's well-being outside clinical settings.

This collaborative team approach underscores communication and coordination as pivotal elements for success. Regular multidisciplinary meetings facilitate knowledge sharing, ensuring all team members are aligned with the child's evolving needs and progress. Such synergy allows for timely adjustments in treatment plans based on comprehensive assessments from all angles—medical, nutritional, physical, psychological—and parental insights.

In essence, adopting a collaborative team approach to pediatric growth management exemplifies modern healthcare's shift towards patient-centered care models that recognize individual uniqueness while leveraging collective expertise for improved outcomes. This model not only enhances the effectiveness of interventions but also fosters an environment where every stakeholder feels valued and integral to achieving optimal health outcomes for children navigating the journey of growth and development amidst an array of potential challenges.

**\*\*Facilitating Better Chewing Efficiency\*\*:** Orthodontic treatment can improve bite function and alignment, allowing for more efficient chewing which is crucial for digestion and overall nutritional health in adulthood.

Orthodontists, as specialists in the field of dentistry, play a pivotal role in the realm of pediatric growth management, particularly through the planning and execution of growth-modifying treatments. This specialized area intertwines intricately with the broader concept of collaborative approaches, highlighting the significance of teamwork among healthcare professionals to ensure optimal outcomes for young patients. Let's delve into how orthodontists navigate this complex landscape, emphasizing the collaborative spirit that underpins successful pediatric growth management.

To begin with, understanding the developmental stages of children is fundamental for orthodontists. The jaw and facial structures are not static; they evolve significantly from childhood into adolescence. Orthodontists leverage this knowledge to identify the ideal moments for intervention—times when growth can be harnessed to correct malocclusions, improve facial aesthetics, and ensure proper function of the jaws. This timing is crucial; early intervention can prevent more complex issues from arising later on, thereby reducing the need for more invasive treatments in adulthood.

The planning phase is where collaboration begins to take center stage. Orthodontists often work alongside general dentists, paediatricians, and sometimes even maxillofacial surgeons to form a comprehensive treatment plan tailored to each child's unique needs. These discussions are not merely about aligning teeth but also involve considering how dental health impacts overall growth and development. For instance, addressing a crossbite or overjet might require insights from a paediatrician regarding underlying medical conditions that could influence growth patterns.

In executing these growth-modifying treatments, orthodontists employ various tools and techniques—from traditional braces to cutting-edge appliances like palatal expanders and functional appliances. Each device serves a specific purpose: some guide jaw growth, others correct tooth positioning, and a few combine these functions to achieve holistic

improvements in oral health and facial aesthetics. Here again, collaboration is key; regular consultations with other specialists ensure that each aspect of treatment complements the others effectively.

Moreover, involving parents and caregivers in the treatment process fosters an environment of shared responsibility and understanding. Orthodontists educate these primary support figures about what to expect during different phases of treatment, how to manage homecare routines effectively, and why certain decisions are made regarding their child's care plan. This educational component strengthens trust between families and healthcare providers while empowering parents to actively participate in their child's journey toward better oral health.

The beauty of collaborative approaches in pediatric growth management lies not just in improved clinical outcomes but also in enhancing patient experience and satisfaction. Children who feel supported by a cohesive team of professionals are more likely to adhere to treatment plans with enthusiasm rather than reluctance—a critical factor considering that compliance directly influences treatment efficacy. Furthermore, such collaborative efforts often lead to interdisciplinary learning opportunities for all involved professionals, continually refining practices based on collective experiences and latest research findings.

In conclusion, orthodontists stand at the forefront of collaborative approaches to pediatric growth management by integrating their specialized knowledge with insights from various healthcare disciplines. Through meticulous planning and execution of growth-modifying treatments—enriched by ongoing collaboration—they not only address immediate dental concerns but also contribute significantly to shaping healthier futures for children navigating their formative years. The synergy among professionals ensures that every child receives personalized care that respects their unique developmental trajectory while nurturing healthy habits that last a lifetime.

Pediatricians play a pivotal role in the intricate tapestry of children's health, serving not just as healthcare providers but as partners in the journey of growth and development. The topic of "Collaborative Approaches to Pediatric Growth Management" encapsulates a multifaceted strategy aimed at ensuring children reach their potential, both physically and developmentally. This approach is underpinned by the understanding that a child's growth is influenced by a myriad of factors—genetic, environmental, nutritional, and emotional—each requiring careful consideration and coordinated action.

The essence of collaborative approaches lies in recognizing that no single professional holds all the answers or resources to manage pediatric growth effectively. It calls for a synergy among various healthcare specialists, including pediatricians, dietitians, endocrinologists, psychologists, and physical therapists, along with the indispensable involvement of families. This teamwork ensures that each child benefits from a comprehensive perspective tailored specifically to their needs.

At the heart of this collaboration is regular monitoring and assessment. Pediatricians are often the primary coordinators who track growth through standardized measures like height-for-age percentiles and weight-for-length percentiles, but they do not work in isolation. They actively engage with dietitians to evaluate nutritional intake and make dietary recommendations that support healthy growth patterns. When concerns about growth retardation or obesity arise—a common challenge in today's fast-paced world—pediatricians collaborate with endocrinologists to explore hormonal imbalances or other medical conditions impacting growth.

Moreover, recognizing the profound impact emotions and psychological factors have on developmental milestones, pediatricians frequently partner with psychologists or child behavior specialists. These collaborations are crucial for addressing issues like anxiety or behavioral challenges that can indirectly affect a child's eating habits or physical activity levels. By weaving together these diverse expertise threads, healthcare teams create a

supportive network around each child, fostering an environment where growth isn't just measured in inches and pounds but also in confidence and well-being.

Parental involvement is another cornerstone of effective collaboration in pediatric growth management. Parents are vital partners who provide insights into their child's daily routines, eating habits, and emotional states—information critical for crafting personalized care plans. Pediatricians strive to educate parents about healthy lifestyle choices while also listening to their observations and concerns. This two-way communication builds trust and empowers families to play an active role in managing their child's growth journey.

In essence, collaborative approaches to pediatric growth management embody a holistic philosophy that transcends traditional boundaries of care delivery. It acknowledges that growing up is not merely a biological process but an experience deeply intertwined with emotional health, nutrition, physical activity, and familial support. As pediatricians navigate this complex landscape alongside fellow professionals and families, they illuminate pathways toward healthier futures for children—a future where every milestone is celebrated not just as a measure of physical development but as an achievement woven into the rich fabric of life itself. Through such unified efforts, we honor the beautiful complexity of childhood development and reaffirm our commitment to nurturing tomorrow's generation with compassionate intelligence and collective wisdom.

The intersection of dental care and pediatric growth management presents a fascinating landscape where the health of a child's teeth and gums is intricately linked to their overall growth and development. Dentists, traditionally viewed as specialists in oral health, are increasingly recognizing their role as integral members of a broader team focused on collaborative approaches to managing pediatric growth. This shift reflects an evolving understanding that oral health is not isolated but rather a cornerstone supporting the well-being of children as they develop.



At the heart of this collaboration lies the recognition that issues in the mouth can significantly impact a child's ability to eat, speak, and even breathe comfortably, all of which are fundamental to their growth and development. For instance, untreated dental cavities or misalignments can lead to pain and discomfort, affecting a child's nutritional intake and potentially leading to malnutrition—an issue that directly influences growth patterns. Moreover, conditions like obstructive sleep apnea, sometimes rooted in anatomical issues within the oral cavity or airway, can disrupt sleep quality, impacting hormonal balances crucial for proper growth.

Dentists contribute uniquely to this collaborative approach by providing specialized insights into the oral manifestations of broader health issues. Their expertise allows them to identify signs of developmental disorders early on—such as certain patterns in tooth eruption or jaw alignment—that might otherwise be overlooked until more severe symptoms arise. By working hand-in-hand with pediatricians, orthodontists, nutritionists, and other healthcare professionals, dentists ensure that oral health considerations are integrated into comprehensive growth management plans for children.

Furthermore, preventive care remains a powerful tool in this collaborative framework. Regular dental check-ups not only help maintain healthy teeth and gums but also serve as opportunities for early intervention. Dentists educate parents and caregivers about the importance of good oral hygiene practices suited for different developmental stages, emphasizing how these habits contribute to overall health. Fluoride treatments and sealants can be pivotal in preventing decay before it starts, safeguarding children's teeth while supporting their general well-being without invasive procedures often associated with corrective treatments later on.

The role dentists play extends beyond just treating conditions; they actively participate in monitoring growth milestones related to oral development. By tracking how a child's jaw grows and teeth erupt over time, dentists provide valuable data points for assessing

overall developmental progress. This ongoing dialogue between dental professionals and other members of the healthcare team ensures that interventions are timely and tailored to each child's unique needs—a critical aspect of nurturing healthy growth trajectories.

In essence, dentists embody a vital piece in the puzzle of pediatric growth management through their specialized focus on oral health—a facet closely intertwined with overall development outcomes. Their collaborative approach underscores the importance of viewing healthcare holistically; understanding that what happens in the mouth can have far-reaching effects on a child's journey toward becoming a healthy adult. As we continue to appreciate this interconnectedness, we foster environments where every aspect of a child's development is considered—ensuring they grow not just taller but healthier too.

### **Early Intervention Strategies in Collaborative Approaches to Pediatric Growth Management**

The journey of childhood growth is a delicate and intricate process, often requiring a multifaceted approach to ensure that children reach their full potential. Early intervention strategies stand as pivotal measures in fostering healthy growth patterns, particularly for children who may face challenges due to various medical, nutritional, or developmental issues. By adopting collaborative approaches involving healthcare professionals, families, and sometimes educators and community resources, we can weave a supportive net around these young individuals, encouraging optimal growth and development from the very start.

At the heart of early intervention is the principle of **timeliness**. The earlier we identify potential growth issues—be they related to nutrition, hormonal imbalances, or genetic conditions—the better equipped we are to address them. Regular monitoring by pediatricians and specialized health teams allows for timely detection of deviations from typical growth trajectories. This proactive stance is crucial; it ensures that interventions

can be tailored and implemented before minor issues escalate into more significant complications.

Collaboration among various professionals forms the backbone of effective early intervention strategies. Pediatricians often lead the charge but work hand-in-hand with dietitians who assess nutritional needs and craft personalized meal plans that resonate with a child's cultural background and preferences, ensuring adherence isn't just a goal but an achievable reality. Physical therapists might join the team to address any motor skill delays that could indirectly impact growth through reduced activity levels. Psychologists play a vital role too, supporting both children and parents through emotional challenges that can accompany growth disorders, fostering a positive mindset essential for successful outcomes.

Family involvement is another cornerstone of these collaborative approaches. Parents and caregivers are not merely recipients of advice; they are integral partners in the care process. Educating families about growth milestones, nutritional choices, and activity levels empowers them to become active contributors to their child's development at home. This partnership builds trust and ensures consistency between clinical recommendations and day-to-day practices—a critical factor in achieving sustainable progress.

Moreover, recognizing the significance of community resources expands our intervention network beyond clinical settings. Schools can become allies in monitoring growth trends through regular check-ups integrated into their health services while also promoting physical activities suitable for every child's ability level. Community programs focusing on nutrition education can offer accessible resources for families seeking guidance on balanced diets within their economic means. These broader networks reinforce the messages received from healthcare providers, creating a seamless web of support around growing children.

In conclusion, early intervention strategies within collaborative approaches to pediatric growth management represent a holistic embrace—one that acknowledges the complexities involved in nurturing healthy development during childhood's formative years. By fostering an environment rich with timely expertise, family engagement, educational support from schools, and leverage on community resources, we not only address immediate health concerns but lay down foundational pathways toward lifelong wellness for our children. It's about recognizing that every child deserves not just survival but thriving—and together, through collaboration and early action, we can make this vision a reality.

The journey of a child's growth is a remarkable and delicate process, fraught with moments of rapid change and development that shape not only their physical stature but also their overall health and well-being. In the intricate landscape of pediatric growth management, the importance of timely assessments and interventions during critical growth periods cannot be overstated. This perspective underscores a collaborative approach, one that weaves together the expertise of healthcare professionals, the insights of parents, and the unique needs of each child to foster optimal development.

During infancy and early childhood, children experience explosive growth spurts that lay the foundation for their future health. These critical periods are windows of opportunity where interventions can have profound effects on long-term outcomes. Timely assessments serve as essential checkpoints to monitor growth patterns, nutritional status, and developmental milestones. They allow healthcare providers to identify deviations from typical growth trajectories early on—whether due to genetic predispositions, environmental factors, or underlying medical conditions—that could otherwise go unnoticed until more severe issues arise.

The collaborative approach here is vital because it pools knowledge and perspectives from various specialists—pediatricians, nutritionists, endocrinologists, psychologists—and families themselves. Each brings a unique lens through which to view a child's growth

journey. Parents often notice subtle changes in behavior or appetite before they are clinically apparent; their insights are invaluable in crafting personalized intervention strategies. Moreover, collaboration ensures that interventions are holistic and address not just physical growth but also emotional and cognitive development—a crucial aspect often intertwined with nutritional and health statuses.

Consideration of cultural practices and family dynamics within this collaborative framework fosters an environment where interventions resonate with the child's lived experiences, enhancing adherence and effectiveness. For instance, dietary recommendations must align with cultural eating habits while still promoting healthy growth; similarly, behavioral strategies should respect familial values while encouraging positive developmental outcomes.

Timely interventions during these critical windows can mitigate risks associated with conditions such as malnutrition, obesity, or hormonal imbalances—conditions that could impact a child's quality of life far into adulthood if not addressed promptly. By intervening early through tailored nutritional plans, physical activity programs designed by pediatric experts in collaboration with families, or even psychological support when necessary, we pave the way for healthier futures for these children.

In essence, nurturing children through their pivotal growth stages requires vigilance and partnership among all stakeholders involved in their care—a collective effort aimed at recognizing potential challenges early on and addressing them proactively through informed assessments and targeted interventions. Such an approach not only supports immediate health needs but equips children with the best possible foundations for lifelong well-being—a testament to what collaborative approaches to pediatric growth management can achieve when timed perfectly with nature's own rhythm of development.

When delving into the intricate world of pediatric growth management, particularly concerning the development of the jaw, it's essential to explore the multifaceted role that functional appliances play. These devices are not merely tools; they represent a cornerstone in collaborative approaches aimed at guiding and optimizing jaw development in growing children. The essence of their application lies in harnessing the malleable nature of a child's developing skeletal structure to foster healthier growth patterns, thereby mitigating potential issues that could arise later in life.

Functional appliances, broadly speaking, are orthodontic devices designed to influence the growth of the jaws and teeth alignment by utilizing the natural forces generated during functions like swallowing, speaking, and chewing. Unlike traditional orthodontic treatments that might focus solely on moving teeth into desired positions post-growth phases, functional appliances work proactively during crucial developmental stages. They encourage optimal positioning and growth of both the upper and lower jaws by guiding them into more harmonious relationships with each other and with the rest of the facial structure.

The beauty—and challenge—of using functional appliances lies in their reliance on patient compliance and timing. Since these appliances capitalize on the body's natural growth processes, their effectiveness is intricately tied to when they are introduced and how consistently they are used. This necessitates a collaborative approach involving parents, orthodontists, pediatricians, and sometimes other specialists like oral surgeons or ENT specialists. Each professional brings unique insights to ensure that treatment is not only effective but also integrated seamlessly into the child's overall health plan.

One fascinating aspect of employing functional appliances is their adaptability to address a variety of conditions, from Class II malocclusions—where the upper jaw protrudes ahead of the lower jaw—to cases requiring expansion of a narrow palate or correction of abnormal swallowing patterns. By actively engaging with these issues early on, practitioners can

potentially reduce or eliminate the need for more invasive procedures later in life, offering children not just immediate benefits but also long-term advantages for their oral health and aesthetics.

Moreover, this collaborative approach emphasizes education—not just for parents but for children themselves—as an integral part of successful treatment outcomes. Understanding why these appliances are important and how they work can motivate younger patients to adhere to wearing schedules and care instructions meticulously. It cultivates a sense of involvement in their own health journey from an early age—a lesson beneficial beyond dental care alone.

In conclusion, functional appliances serve as a testament to how thoughtful collaboration across disciplines can lead to innovative solutions in pediatric growth management. Their role in guiding jaw development isn't just about correcting misalignments; it's about nurturing balanced facial growth that supports overall health and confidence throughout life's stages. As we continue refining these techniques and expanding interdisciplinary cooperation, we pave the way for even more personalized and effective care for our youngest patients navigating these critical developmental years.

The integration of orthodontic treatments with other medical specialties represents a pivotal shift in how we approach pediatric growth management, particularly when considering the intricate interplay between dental development and overall health. This collaborative approach is not merely a trend; it's a recognition that the health and well-being of children are multifaceted, requiring a team effort that transcends traditional disciplinary boundaries.

At the heart of this integrative strategy lies the understanding that orthodontic issues can have far-reaching effects beyond mere aesthetics or functional concerns related to teeth alignment. Conditions like malocclusions, jaw discrepancies, and developmental anomalies

can impact a child's breathing, speech, chewing efficiency, and even self-esteem. By integrating orthodontics with other specialties such as pediatrics, otolaryngology (ear, nose, and throat), pulmonology (breathing disorders), and sometimes even psychology, healthcare providers can offer comprehensive care that addresses the whole child rather than isolated symptoms.

One of the most significant benefits of this collaborative approach is in managing growth-related issues early on. Pediatricians, being the primary caregivers during early childhood, play a crucial role in identifying potential orthodontic concerns alongside monitoring general health milestones. Early detection allows for timely referrals to orthodontists who can then work in tandem with pediatricians to create individualized treatment plans that consider both immediate dental needs and long-term growth trajectories.

Moreover, incorporating insights from specialists like otolaryngologists becomes essential when dealing with conditions such as sleep apnea or airway obstruction issues which are often linked to craniofacial anomalies. Here, an integrated approach ensures that while correcting dental misalignments through braces or expanders might be part of the solution, addressing underlying airway problems could involve surgical interventions or specific therapeutic strategies recommended by experts in respiratory conditions. This holistic view ensures that treatments are not only effective but also considerate of their broader impact on a child's health and development.

Additionally, psychological aspects cannot be overlooked when discussing pediatric growth management. Children facing orthodontic challenges may experience social stigma or emotional distress related to their appearance or difficulties in daily activities like eating or speaking clearly. Collaborating with psychologists or child development experts ensures that these emotional needs are met alongside physical treatments. This support system fosters resilience and confidence in children undergoing treatment, making the process less daunting for them and their families.



In essence, the integration of orthodontic treatments within a broader framework of pediatric growth management exemplifies modern medicine's move towards personalized and holistic care. It underscores the importance of interdisciplinary collaboration where professionals from diverse backgrounds pool their expertise to craft solutions tailored to each child's unique needs. As we continue to refine this collaborative approach, we not only enhance treatment outcomes but also contribute significantly to improving the quality of life for children navigating their formative years—a testament to how interconnected our approaches to health truly are.

Collaboration with endocrinologists in managing conditions affecting growth, such as dwarfism and gigantism, is a cornerstone of pediatric growth management. This collaborative approach represents a multidisciplinary effort that brings together the expertise of various specialists to ensure children receive comprehensive care tailored to their unique needs. Let's delve into why this partnership is not just beneficial but essential in navigating the complexities of pediatric growth disorders.

At the heart of these conditions lies an intricate balance of hormones that regulate growth, primarily influenced by the endocrine system. Endocrinologists, specialists in hormonal disorders, play a pivotal role in diagnosing and treating conditions like dwarfism—often due to genetic mutations affecting growth hormone production or responsiveness—and gigantism, which is typically characterized by excessive production of growth hormone, usually due to a tumor on the pituitary gland. The nuances of these hormonal imbalances require a deep understanding that general pediatricians might not possess in depth, underscoring the necessity for endocrinological expertise.

The collaborative approach begins with accurate diagnosis. Endocrinologists utilize specialized tests including hormone level assessments and imaging studies to pinpoint the exact cause and nature of growth anomalies. This precise diagnostic step is crucial; it sets the stage for developing an effective treatment plan tailored specifically to each child's condition and potential complications.

Treatment itself is multifaceted, often involving medication adjustments to regulate hormone levels, surgical interventions when tumors are involved, and psychological support to address the social and emotional challenges these conditions can pose. Here's where collaboration truly shines: endocrinologists work hand-in-hand with pediatric endocrinology nurses, nutritionists, physical therapists, psychologists, and sometimes surgeons or orthopedic specialists. Each professional contributes their expertise to create a holistic care plan that addresses not just the physiological aspects but also the psychosocial challenges faced by children with growth disorders and their families.

Moreover, this collaborative framework fosters continuous education and innovation. Regular meetings among team members facilitate discussions on emerging research findings, treatment advancements, and shared case studies, ensuring that every child benefits from cutting-edge knowledge and practices. For parents navigating these complex medical landscapes alongside their children, having a cohesive team communicates a united front dedicated to their child's well-being—a powerful reassurance during what can be an uncertain journey.

In conclusion, collaboration between endocrinologists and other healthcare professionals in managing conditions affecting growth such as dwarfism and gigantism exemplifies modern pediatric care at its best. It embodies an integrative approach that recognizes no single specialty holds all the answers for such complex medical challenges. Through this synergy of expertise—hormonal mastery meeting surgical precision alongside psychological support—these multidisciplinary teams empower children with growth disorders to lead healthier, more fulfilling lives while supporting their families every step of the way.

Coordinating efforts with geneticists in the context of congenital anomalies is a cornerstone of effective pediatric growth management, particularly when we delve into the intricate world of collaborative approaches. This intersection of expertise is vital in ensuring that children with congenital anomalies receive comprehensive care that

addresses not just their immediate health needs but also their long-term growth and development.

To understand the significance of this collaboration, one must first grasp the complexity of congenital anomalies—birth defects that originate during fetal development and can affect virtually any part of the body. These anomalies range from relatively minor to life-threatening conditions, impacting physical, cognitive, and sometimes emotional growth. As such, they necessitate a multidisciplinary approach where geneticists play a pivotal role alongside pediatricians, surgeons, therapists, and other specialists.

Geneticists bring an invaluable perspective to the table due to their deep understanding of how genetic variations contribute to congenital anomalies. By identifying specific genetic factors at play, they can provide insights into the potential course of a child's condition, guiding treatment strategies and offering families crucial information about risks for future offspring. This knowledge is foundational for personalized medicine approaches that tailor interventions to the unique genetic makeup of each child.

In practice, collaboration between geneticists and other healthcare providers involved in pediatric growth management might manifest in several ways. Firstly, early intervention is key; upon diagnosing a congenital anomaly through prenatal screening or post-birth examination, prompt consultation with a geneticist can clarify the nature of the anomaly and its implications for growth patterns. This allows for timely dietary adjustments, hormonal therapies, or surgical interventions tailored to correct or mitigate developmental delays associated with the condition.

Moreover, ongoing communication between geneticists and other team members ensures that as new research emerges—be it about gene therapy advancements or better understanding hereditary patterns—this information seamlessly integrates into the child's

ongoing care plan. Such coordinated efforts underscore a holistic approach where every aspect of a child's health is considered within a broader familial and genetic context.

Beyond medical interventions, psychological support becomes equally critical given the emotional toll that congenital anomalies can take on families. Genetic counselors often work hand-in-hand with these teams to provide guidance not only on medical prognoses but also on navigating complex decisions regarding reproductive options for future pregnancies—a testament to how deeply interwoven these fields become in supporting affected families.

In essence, collaborative approaches involving coordination with geneticists are not merely beneficial but essential in managing pediatric growth amidst congenital anomalies. This synergy amplifies our ability to address these complex conditions comprehensively—embracing medical science's cutting edge while grounding care in empathy and understanding tailored to each unique family journey. It's this comprehensive teamwork that promises brighter futures for children facing such challenges—a future where growth management isn't just about overcoming obstacles but thriving despite them.

In the ever-evolving landscape of pediatric growth management, the integration of technology has emerged not merely as a trend but as an essential pillar in fostering enhanced collaboration among healthcare professionals, families, and the young patients themselves. This synergy between human expertise and technological innovation marks a significant shift in how we approach the complex journey of monitoring and supporting children's growth, ensuring that every step is informed, connected, and tailored to individual needs.

At the heart of this transformation lies the ability of technology to bridge distances and break down barriers that traditionally hindered collaborative efforts. Telemedicine platforms, for instance, have revolutionized access to specialized care. Imagine a scenario

where a rural family can consult with leading pediatric endocrinologists without leaving their hometown. Through high-definition video conferencing and secure sharing of medical records, these families receive expert advice tailored to their unique circumstances, making collaborative care more inclusive and accessible than ever before.

Moreover, technological advancements have equipped healthcare teams with sophisticated tools for data collection and analysis. Wearable devices and mobile health applications allow continuous monitoring of vital growth indicators such as height, weight, and developmental milestones. This real-time data not only empowers parents to actively participate in their child's growth management but also enables healthcare professionals to observe trends and intervene promptly when necessary. The immediacy of this feedback loop fosters a more dynamic and responsive collaborative environment where decisions are made based on up-to-date information rather than historical data alone.

Furthermore, digital platforms designed specifically for pediatric growth management facilitate seamless communication among all stakeholders—doctors, nurses, dietitians, physical therapists, and parents. These platforms often feature shared electronic health records (EHRs), allowing each team member to contribute insights and updates in one cohesive space. This collective view ensures everyone is on the same page regarding treatment plans, progress reports, and adjustments needed along the way. Such collaboration not only enhances the quality of care but also strengthens trust among families who feel supported by a cohesive team dedicated to their child's well-being.

Educational resources available online further enrich this collaborative ecosystem. Parents can access a wealth of information about pediatric growth patterns, nutritional guidelines tailored to different ages and activity levels, as well as emotional support forums where they connect with other families navigating similar journeys. Armed with knowledge facilitated by technology, parents become active partners in decision-making processes rather than passive recipients of advice—a crucial shift towards empowering family-

centered care.

However, it's vital to acknowledge that while technology amplifies our collaborative capabilities significantly, it must be implemented thoughtfully to ensure it complements rather than replaces the invaluable human touch in pediatric care. The warmth of a nurse's reassuring words or the nuanced understanding conveyed through eye contact remain irreplaceable aspects of caregiving that technology cannot fully replicate. Thus, embracing technology for enhanced collaboration should be viewed as augmenting traditional methods rather than replacing them entirely.

In conclusion, "Utilizing Technology for Enhanced Collaboration" within the realm of Pediatric Growth Management embodies a promise—a future where cutting-edge tools work hand-in-hand with compassionate human interaction to navigate the intricate pathways of childhood development effectively. As we continue to innovate within this domain, let us remember that our ultimate goal is not just improved metrics or streamlined processes but healthier outcomes rooted in comprehensive collaboration among all those who play a role in nurturing our youngest generation's potential into reality.

The integration of digital imaging and 3D modeling into the realm of pediatric growth management represents a transformative leap in how healthcare professionals approach the complex task of monitoring and supporting children's development. This innovative application marries technology with the nuanced understanding required in pediatric care, facilitating precise planning and enhancing communication among multidisciplinary team members—a cornerstone for effective collaborative approaches.

To begin with, digital imaging, including advanced techniques like MRI, CT scans, and ultrasound, provides an unparalleled window into a child's internal anatomy and growth patterns. These tools allow for detailed visualizations that were once impossible to

achieve with traditional methods. For instance, clinicians can assess bone density, organ development, and even subtle changes in craniofacial structures with precision that directly translates to more personalized care plans. The ability to observe growth over time through serial imaging offers invaluable insights into the effectiveness of interventions and adjustments needed along the way.

Parallel to this, 3D modeling takes the data gleaned from these imaging technologies and transforms it into tangible representations. Imagine being able to hold a physical model of a child's skeletal system or a detailed anatomical structure right on your desk—that's what 3D modeling offers. These models not only aid in clearer visualization but also serve as excellent educational tools for both medical teams and patients (and their families). They simplify complex anatomical concepts, making discussions about growth abnormalities or necessary surgical interventions more accessible and understandable. This clarity fosters better-informed consent processes and empowers families by visualizing potential outcomes and treatment paths.

The synergy between digital imaging and 3D modeling significantly enhances collaborative efforts among team members—pediatricians, surgeons, radiologists, physical therapists, nutritionists, psychologists, and more. With shared digital platforms showcasing up-to-date models and imaging results, teams can convene virtually or in person around a common reference point. This unified perspective ensures that everyone is on the same page regarding patient progress and upcoming strategies. It streamlines decision-making processes by providing objective data that can be discussed openly within the team framework, reducing miscommunications often associated with relying solely on textual reports or memory recall of previous images.

Moreover, this technological marriage doesn't just stop at improving intra-team collaboration; it extends its benefits to remote consultations. With high-resolution digital files easily shareable across vast distances via secure networks, opinions from around the

globe can be sought swiftly. This capability is particularly crucial for rare conditions requiring specialized knowledge outside local expertise, ensuring children receive comprehensive care tailored to their unique needs.

In conclusion, the incorporation of digital imaging and 3D modeling into pediatric growth management epitomizes a forward-thinking approach towards collaborative patient care. By furnishing precise diagnostic capabilities, these technologies enhance clinical outcomes and enrich the overall experience for children.

Teleorthodontics, a burgeoning field at the intersection of telemedicine and orthodontics, has revolutionized the way professionals approach patient care, especially in complex areas like pediatric growth management. This innovative approach leverages technology to facilitate remote consultations and foster collaboration among specialists, marking a significant shift from traditional, geographically bound practices. The essence of teleorthodontics lies in its ability to bridge distances, making expert advice and collaborative efforts more accessible than ever before, particularly for addressing the nuanced needs of growing children.

The concept of collaborative approaches to pediatric growth management is rooted in recognizing that a child's dental and skeletal development is not isolated but intricately linked with their overall growth and health. Orthodontists, pediatric dentists, maxillofacial surgeons, endocrinologists, and nutritionists are just a few of the specialties that might converge in managing a child's growth trajectory. Traditionally, coordinating between these diverse professionals could be cumbersome, requiring physical presence in the same location—a logistical nightmare that could potentially delay critical interventions.

Enter teleorthodontics: this technology enables real-time video consultations where specialists can discuss cases collaboratively without the constraints of physical proximity. Imagine an orthodontist in New York reviewing X-rays and 3D models alongside a pediatric



endocrinologist based in California while discussing a patient's growth patterns influenced by hormonal factors. Such seamless interaction not only expedites decision-making but enriches it with multidisciplinary insights crucial for comprehensive care plans tailored specifically for pediatric patients.

Moreover, teleorthodontics enhances accessibility for families who might otherwise face significant challenges traveling to specialized centers regularly. Children with unique growth management needs often require frequent monitoring; teleorthodontics allows for periodic check-ins that are both convenient and less disruptive to their daily lives. This continuity of care is pivotal as it ensures that any adjustments in treatment plans can be discussed promptly among the involved specialists, fostering an adaptable approach suited to each child's evolving needs.

Furthermore, this collaborative platform facilitates knowledge sharing on cutting-edge techniques and research findings pertinent to pediatric growth management. Continuing education becomes less about individual pursuit and more about collective advancement—a scenario where insights gleaned from various disciplines can immediately benefit patient outcomes. The synergy created through such interactions catalyzes innovation within orthodontic practices aimed at optimizing growth trajectories for children under their care.

However, as we embrace the benefits of teleorthodontics in collaborative approaches to pediatric growth management, it's essential to acknowledge potential challenges—issues ranging from technological disparities across regions to ensuring robust data security measures are upheld during virtual exchanges of sensitive health information. Overcoming these hurdles requires concerted efforts from technology providers, healthcare institutions, and regulatory bodies to create an environment conducive to safe and effective remote collaboration.

In conclusion, teleorthodontics stands as a testament to how modern technology can transform specialized fields like orthodontics into dynamic platforms for interdisciplinary collaboration—especially beneficial for intricate areas such as pediatric growth management. By breaking down geographical barriers and fostering rich dialogues among diverse specialists, we pave the way for nurturing healthier futures for our youngest patients through coordinated care that's both informed and responsive to individual needs. As we move forward into this digitally interconnected era of healthcare delivery, embracing such innovations promises not just advancements in treatment methodologies but also profound enhancements in patient experiences and outcomes alike.

When we delve into the realm of pediatric growth management, it becomes evident that a one-size-fits-all approach is not just less effective but often counterproductive. The essence of fostering healthy growth in children lies in recognizing and embracing their unique journeys—each child a distinct tapestry woven from genetics, environment, lifestyle, and personal health history. This realization underscores the critical importance of customized treatment plans, meticulously tailored to meet the individual needs of each young patient.

In collaborative approaches to pediatric growth management, this personalization isn't merely an idealistic goal; it's a pivotal strategy for achieving optimal outcomes. At the heart of these customized treatment plans is a thorough assessment that goes beyond mere measurements of height and weight. It involves understanding the child's medical history, dietary habits, physical activity levels, psychological well-being, and even family dynamics. By weaving together insights from these various threads, healthcare providers can craft a comprehensive picture of the child's growth trajectory and potential hurdles they might face.

Collaboration here extends beyond just doctors and parents; it includes engaging with nutritionists, physical therapists, psychologists, and educators to create a support network that addresses all facets of the child's development. Each professional brings

their expertise to the table, contributing pieces to a holistic puzzle aimed at enhancing growth while ensuring overall well-being. For instance, a nutritionist might work closely with the family to develop meal plans that not only promote growth but also cater to any food preferences or allergies the child may have. Meanwhile, a physical therapist could design age-appropriate exercises that encourage active growth without overwhelming or injuring developing bodies.

Moreover, personalized treatment plans emphasize regular monitoring and adaptability—they are living documents that evolve as the child grows and circumstances change. Regular check-ins allow for timely adjustments in response to progress or new challenges that may arise. This dynamic approach ensures that interventions remain relevant and effective throughout different stages of childhood and adolescence when growth patterns can shift dramatically.

The benefits of such tailored strategies are manifold. Children who receive customized care often experience improved adherence to treatment plans since they are more likely to engage with approaches that feel understood and relevant to their lives. There's also evidence suggesting better health outcomes—whether in terms of reaching healthy growth milestones or managing conditions like obesity or hormonal imbalances more effectively. Above all, this individualized focus fosters an environment where children feel supported rather than simply 'treated,' nurturing their confidence and self-esteem as they navigate their developmental journey.

In conclusion, customized treatment plans represent a paradigm shift towards more humane and effective pediatric growth management—a shift grounded in collaboration and personalization. By honoring each child's unique path with bespoke strategies crafted through teamwork among various specialists, we not only promote healthier physical development but also contribute significantly to their emotional well-being and future quality of life. It's about recognizing that every step forward in pediatric care should be

guided by compassionate understanding tailored precisely to who each growing child is at their core.

When we delve into the realm of pediatric growth management, it becomes increasingly clear that a one-size-fits-all approach is not only impractical but often ineffective. The importance of personalized strategies in this context cannot be overstated, especially when considering each child's unique growth trajectory. This individualized approach acknowledges the intricate tapestry of factors—genetic, environmental, nutritional, and even psychological—that influence a child's development. By embracing collaborative approaches that tailor interventions to the specific needs of each child, healthcare providers, educators, and families can weave together a supportive network that fosters optimal growth and development.

Every child is born with an inherent potential shaped by their genetic blueprint, yet this potential unfolds differently for each individual. Some children may experience rapid growth in the early years, while others might have a more gradual pattern. Environmental factors such as socioeconomic status, access to nutritious food, and exposure to toxins can markedly alter this natural trajectory. Additionally, lifestyle choices and family dynamics play pivotal roles in shaping a child's health outcomes. Acknowledging these variables compels us to move beyond generic guidelines towards customized care plans that respect and respond to each child's unique circumstances.

Collaborative approaches to pediatric growth management exemplify the power of teamwork among healthcare professionals, parents, and sometimes even the children themselves. By bringing together pediatricians, nutritionists, psychologists, physical therapists, and educators under one umbrella of shared goals—ensuring the child thrives physically, mentally, and emotionally—we create a comprehensive support system tailored precisely to the child's evolving needs. Such collaboration allows for continuous assessment and adjustment of strategies as the child grows and faces new developmental milestones or challenges.

Moreover, involving children in their own growth management journey cultivates self-awareness and empowerment from an early age. As they understand their unique needs and participate actively in setting goals with their care team, they become advocates for their health—a skill beneficial throughout life. This collaborative spirit encourages open communication between all parties involved; parents share insights about their child's behavior at home while professionals offer expert guidance based on observed patterns and data-driven recommendations.

In essence, recognizing the significance of personalized strategies within collaborative frameworks for pediatric growth management reflects a profound respect for human complexity and diversity. It underscores our commitment to nurturing not just physical health but holistic well-being through understanding and responding to each child's singular journey toward maturity. By prioritizing individuality in growth management strategies, we pave the way for healthier futures where every child reaches their full potential in their own time and way.

Certainly, exploring the nuances of collaborative management in pediatric growth management through case studies offers a compelling narrative of how teamwork and shared strategies can lead to remarkable outcomes in children's health. The essence of collaborative approaches lies in merging the expertise of various professionals—pediatricians, dietitians, psychologists, physical therapists, and sometimes even educators—to address the multifaceted needs of growing children. Let's delve into a couple of illustrative case studies that highlight the success of such integrated efforts.

## **Case Study 1: The Growing Together Program**

In a bustling metropolitan city, the "Growing Together" program was initiated to tackle the rising concerns of childhood obesity and its associated growth abnormalities. Recognizing that addressing this issue required more than just nutritional advice, the program brought

together a diverse team including pediatric endocrinologists, nutritionists, school cafeteria managers, and community fitness instructors.

The core strategy involved comprehensive assessments of each child's growth trajectory alongside dietary habits and physical activity levels. What made this initiative stand out was its community-centric approach; parents were actively involved in workshops designed to educate them on balanced diets and engaging activities for their kids. Schools played a pivotal role by integrating healthier meal options and encouraging daily physical exercises. Over two years, participants not only showed significant improvements in BMI percentiles but also reported enhanced family cohesion around health goals—a testament to how collaborative efforts can weave together personal health with community well-being.

## **Case Study 2: The Pediatric Growth Clinic Model**

Another inspiring example emerges from a rural clinic that transformed its approach to pediatric growth management through a collaborative model dubbed the "Pediatric Growth Clinic." Here, doctors partnered with local agricultural experts to address food security issues directly impacting children's nutrition. The clinic didn't just treat symptoms; it sought to understand and mitigate the root causes affecting growth patterns among local children.

By working closely with farmers to promote locally grown nutritious foods and educating families on simple yet effective home gardening techniques, the clinic fostered an environment where healthy eating became accessible and culturally relevant. This collaboration extended beyond healthcare providers; it included local government agencies that supported policies favoring sustainable farming practices beneficial for child nutrition. Within three years, not only did childhood obesity rates decline notably

within the clinic's catchment area, but there was also an increase in overall vegetable consumption among children—a clear indicator of successful collaborative outcomes transcending traditional healthcare boundaries.

## Reflections on Collaborative Success

These case studies underscore several key elements crucial for successful collaborative management in pediatric growth: **inclusivity**—bringing together diverse expertise; **community engagement**—ensuring solutions are rooted in cultural relevance and practical applicability; and **systemic thinking**—addressing broader socio-economic factors influencing childhood nutrition and growth. Such collaborations not only improve individual health outcomes but also cultivate community resilience against future challenges related to pediatric health management. As we look towards nurturing healthier generations, embracing these collaborative frameworks becomes increasingly vital—a reminder that together, we can accomplish much more than alone.

### Monitoring Progress and Adjusting Treatments Collaboratively: A Cornerstone of Pediatric Growth Management

The journey of pediatric growth management is a tapestry woven with threads of medical expertise, familial involvement, and the unique growth narrative of each child. At its heart lies a fundamental principle: monitoring progress and adjusting treatments collaboratively. This approach is pivotal not just for effectiveness but for ensuring that every child receives care that is as personalized as their fingerprints.

In the realm of pediatric growth management, which encompasses a spectrum of conditions from growth hormone deficiencies to genetic disorders affecting development,

the initial steps often involve a comprehensive evaluation by a multidisciplinary team. This team typically includes endocrinologists, pediatricians, nutritionists, psychologists, and sometimes even surgeons, depending on the child's specific needs. The first meeting sets the stage; it's where parents and caregivers share their observations and concerns, laying the groundwork for a partnership rooted in mutual trust and understanding. Here, the significance of collaboration becomes evident: it's not just about diagnosing and treating; it's about understanding the whole child within their family context.

As treatment plans are formulated—often involving medications, dietary adjustments, or physical therapies—the importance of continuous monitoring cannot be overstated. Regular follow-ups are scheduled not merely as checkpoints but as opportunities for open dialogue. These visits allow healthcare providers to assess physical growth alongside emotional and social development, ensuring that interventions remain holistically beneficial. For instance, while growth hormone therapy can significantly impact height, its effects on bone density or metabolic health require vigilant monitoring. Equally crucial is how these treatments affect a child's self-esteem and interactions with peers—a consideration often brought to light through collaborative discussions with families.

Adjustments to treatment plans are an inevitable part of this journey. Children grow not only physically but also in their understanding of their condition and treatment needs. What works at one stage might need recalibration as they enter puberty or face new challenges. This adaptability hinges on ongoing communication between healthcare providers and families. Parents become acute observers of their child's responses to treatment—both positive and negative—and their insights are invaluable in fine-tuning approaches for optimal outcomes. When children feel heard and involved in discussions about their care—age-appropriately—they develop a sense of agency that can empower them throughout their lives.



Moreover, this collaborative model fosters resilience among families navigating what can be an intricate landscape of medical appointments and lifestyle adjustments. It builds networks of support—peer connections through support groups or online communities—that share experiences and strategies for managing day-to-day challenges effectively. The collective wisdom gathered through these interactions enriches individual care plans, introducing practical solutions that might not emerge within traditional clinical settings alone.

In essence, monitoring progress and adjusting treatments collaboratively is more than a methodical process; it's an embrace of partnership in which every stakeholder contributes to the richness of the child's developmental story. It reaffirms that growing up shouldn't be done in isolation but rather within a supportive ecosystem where every voice counts—and where each step forward is celebrated collectively, reflecting the multifaceted journey toward achieving optimal pediatric growth management.

When considering the topic of "Collaborative Approaches to Pediatric Growth Management," it's essential to highlight the significance of implementing regular multidisciplinary reviews as a cornerstone strategy. This approach is not merely a checkbox on a checklist but rather a vital, dynamic process that ensures children receive the most appropriate and effective care as they grow and develop.

The essence of pediatric growth management lies in its inherently complex nature, intertwining physical health, nutrition, psychology, and developmental milestones. Given this complexity, relying on a single perspective or discipline can lead to gaps in understanding and intervention. Herein lies the brilliance of adopting collaborative approaches—bringing together a diverse team of specialists under a unified goal: the optimal growth and well-being of the child.

Regular multidisciplinary reviews serve as pivotal moments in this journey. These gatherings are more than mere meetings; they are opportunities for deep dialogue, rich with insights from various experts including pediatricians, dietitians, psychologists, endocrinologists, and physical therapists, among others. Each professional contributes their unique lens through which to view the child's progress, ensuring that no stone is left unturned in assessing growth patterns, addressing challenges, and celebrating achievements.

The frequency of these reviews is crucial; they should be scheduled not just periodically but as needed, responding to the evolving needs of each child. For instance, during critical growth phases or when significant changes in health status occur—such as after introducing new treatments or dietary adjustments—these reviews become even more indispensable. They allow for timely adaptations in management strategies to align with the child's current trajectory, thereby enhancing outcomes significantly.

Moreover, these collaborative sessions foster an environment where team members not only share their observations but also learn from one another. This cross-disciplinary exchange enriches everyone's understanding of pediatric growth management and encourages innovative problem-solving approaches tailored specifically to each child's needs. It cultivates a culture of continuous learning and improvement within the team itself—a powerful driver for excellence in patient care.

Parents and caregivers play an integral role in these discussions too. Their insights into the child's behavior at home complement clinical observations perfectly. By involving them actively in these reviews, healthcare providers ensure that strategies developed are not only clinically sound but also practically feasible within the family's daily life—crucial for long-term success.

In conclusion, regular multidisciplinary reviews are far more than procedural formalities; they are essential conduits for delivering comprehensive and adaptive care in pediatric growth management. By weaving together diverse expertise through collaborative approaches, we create a robust support system that addresses every facet of a child's development—physical, emotional, nutritional—and positions them for thriving throughout their formative years and beyond. This human-centric model underscores our commitment to not just managing growth but nurturing it holistically for each unique individual entrusted to our care.

Engaging parents and patients in the decision-making process is a cornerstone of effective pediatric growth management, particularly when it comes to ensuring adherence to treatment plans. This collaborative approach not only empowers families but also significantly enhances the outcomes of treatment by fostering a sense of ownership and understanding over the child's health journey. The essence of this strategy lies in recognizing that children, especially as they grow older, along with their parents, are integral parts of the healthcare team.

The foundation of engaging parents and patients begins with open communication. Healthcare providers must create an environment where discussions are welcoming, informative, and respectful of the family's perspectives and concerns. This means taking the time to explain medical terms in layman's terms, detailing how various treatments work, their potential benefits, risks, and what adherence entails. When parents understand these facets deeply, they are better equipped to support their child's treatment plan actively.

Moreover, involving children in their care from an early age instills a sense of responsibility and curiosity about their health. As children mature, they can be gradually introduced to more complex aspects of their treatment plans—understanding why certain medications or lifestyle changes are necessary can motivate them to stick to these plans voluntarily rather than out of obligation. This autonomy-supportive approach is crucial; it encourages

young patients to view themselves as active participants in managing their growth and health rather than passive recipients of care.

Furthermore, establishing clear goals together with the family sets a roadmap for success. These goals should be realistic, tailored to the individual needs and circumstances of each patient, and regularly revisited to reflect progress or adjust strategies as necessary. By setting achievable milestones—be it related to medication adherence, dietary modifications, or physical activity—families feel a sense of accomplishment which reinforces positive behaviors and adherence patterns.

Support systems also play a pivotal role in this collaborative approach. Parents often benefit from resources like support groups or educational materials that provide insights into other families' experiences with similar challenges. These networks can offer practical tips on managing daily routines around treatment plans while providing emotional support during tough times. Healthcare providers can facilitate access to such resources or even organize community workshops where experts share knowledge directly with families.

In addition to these strategies, leveraging technology can greatly enhance engagement efforts. Mobile apps designed for pediatric health management allow for easy tracking of medication schedules, growth milestones, and communication between healthcare providers and families. Such tools not only remind but also educate users about best practices in growth management in an interactive manner that appeals particularly well to tech-savvy youth and their parents alike.


In conclusion, engaging parents and patients in decision-making regarding pediatric growth management is not just beneficial—it's essential for successful treatment adherence and long-term health outcomes. By fostering an environment rooted in collaboration, open communication, education, goal-setting, support networks—and now aided by technology—healthcare providers empower families to take charge of their child's

health journey collectively. This approach ensures that every step taken towards managing pediatric growth is informed by those who know the child best—their parents—and the child themselves—a synergy that lays a solid foundation for healthy development throughout life's stages.

### About pediatrics

This article is about the branch of medicine. For the journal, see Pediatrics (journal). For the branch of dentistry, see Pedodontics.

### Pediatrics



A pediatrician examines a neonate.


<b>Focus</b>	<b>Infants, Children, Adolescents, and Young Adults</b>
<b>Subdivisions</b>	<i>Paediatric cardiology, neonatology, critical care, pediatric oncology, hospital medicine, primary care, others (see below)</i>
<b>Significant diseases</b>	Congenital diseases, Infectious diseases, Childhood cancer, Mental disorders
<b>Significant tests</b>	World Health Organization Child Growth Standards
<b>Specialist</b>	<b>Pediatrician</b>
<b>Glossary</b>	Glossary of medicine

**Pediatrics** (American English) also spelled **paediatrics** (British English), is the branch of medicine that involves the medical care of infants, children, adolescents, and

young adults. In the United Kingdom, pediatrics covers many of their youth until the age of 18.<sup>[1]</sup> The American Academy of Pediatrics recommends people seek pediatric care through the age of 21, but some pediatric subspecialists continue to care for adults up to 25.<sup>[2]</sup><sup>[3]</sup> Worldwide age limits of pediatrics have been trending upward year after year.<sup>[4]</sup> A medical doctor who specializes in this area is known as a **pediatrician**, or **paediatrician**. The word *pediatrics* and its cognates mean "healer of children", derived from the two Greek words: *παιδίον* (*paîdion* "child") and *ἰατρός* (*iátrōs* "doctor, healer"). Pediatricians work in clinics, research centers, universities, general hospitals and children's hospitals, including those who practice pediatric subspecialties (e.g. neonatology requires resources available in a NICU).

## History

[edit]



Part of Great Ormond Street Hospital in London, United Kingdom, which was the first pediatric hospital in the English-speaking world.

The earliest mentions of child-specific medical problems appear in the *Hippocratic Corpus*, published in the fifth century B.C., and the famous *Sacred Disease*. These publications discussed topics such as childhood epilepsy and premature births. From the first to fourth centuries A.D., Greek philosophers and physicians Celsus, Soranus of Ephesus, Aretaeus, Galen, and Oribasius, also discussed specific illnesses affecting children in their works, such as rashes, epilepsy, and meningitis.<sup>[5]</sup> Already Hippocrates, Aristotle, Celsus, Soranus, and Galen<sup>[6]</sup> understood the differences in

growing and maturing organisms that necessitated different treatment: *Ex toto non sic pueri ut viri curari debent* ("In general, boys should not be treated in the same way as men").<sup>[7]</sup> Some of the oldest traces of pediatrics can be discovered in Ancient India where children's doctors were called *kumara bhrtya*.<sup>[6]</sup>

Even though some pediatric works existed during this time, they were scarce and rarely published due to a lack of knowledge in pediatric medicine. *Sushruta Samhita*, an ayurvedic text composed during the sixth century BCE, contains the text about pediatrics.<sup>[8]</sup> Another ayurvedic text from this period is *Kashyapa Samhita*.<sup>[9][10]</sup> A second century AD manuscript by the Greek physician and gynecologist Soranus of Ephesus dealt with neonatal pediatrics.<sup>[11]</sup> Byzantine physicians Oribasius, Aëtius of Amida, Alexander Trallianus, and Paulus Aegineta contributed to the field.<sup>[6]</sup> The Byzantines also built *brephotrophia* (crèches).<sup>[6]</sup> Islamic Golden Age writers served as a bridge for Greco-Roman and Byzantine medicine and added ideas of their own, especially Haly Abbas, Yahya Serapion, Abulcasis, Avicenna, and Averroes. The Persian philosopher and physician al-Razi (865–925), sometimes called the father of pediatrics, published a monograph on pediatrics titled *Diseases in Children*.<sup>[12][13]</sup> Also among the first books about pediatrics was *Libellus [Opusculum] de aegritudinibus et remediis infantium* 1472 ("Little Book on Children Diseases and Treatment"), by the Italian pediatrician Paolo Bagellardo.<sup>[14][5]</sup> In sequence came Bartholomäus Metlinger's *Ein Regiment der Jungerkinder* 1473, Cornelius Roelans (1450–1525) no title Buchlein, or Latin compendium, 1483, and Heinrich von Louffenburg (1391–1460) *Versehung des Leibs* written in 1429 (published 1491), together form the *Pediatric Incunabula*, four great medical treatises on children's physiology and pathology.<sup>[6]</sup>

While more information about childhood diseases became available, there was little evidence that children received the same kind of medical care that adults did.<sup>[15]</sup> It was during the seventeenth and eighteenth centuries that medical experts started offering specialized care for children.<sup>[5]</sup> The Swedish physician Nils Rosén von Rosenstein (1706–1773) is considered to be the founder of modern pediatrics as a

medical specialty,<sup>[16][17]</sup> while his work *The diseases of children, and their remedies* (1764) is considered to be "the first modern textbook on the subject"<sup>[18]</sup> However, it was not until the nineteenth century that medical professionals acknowledged pediatrics as a separate field of medicine. The first pediatric-specific publications appeared between the 1790s and the 1920s.<sup>[19]</sup>

## Etymology

[edit]

The term pediatrics was first introduced in English in 1859 by Abraham Jacobi. In 1860, he became "the first dedicated professor of pediatrics in the world."<sup>[20]</sup> Jacobi is known as the *father of American pediatrics* because of his many contributions to the field.<sup>[21][22]</sup> He received his medical training in Germany and later practiced in New York City.<sup>[23]</sup>

The first generally accepted pediatric hospital is the *Hôpital des Enfants Malades* (French: *Hospital for Sick Children*), which opened in Paris in June 1802 on the site of a previous orphanage.<sup>[24]</sup> From its beginning, this famous hospital accepted patients up to the age of fifteen years,<sup>[25]</sup> and it continues to this day as the pediatric division of the Necker-Enfants Malades Hospital, created in 1920 by merging with the nearby *Necker Hospital*, founded in 1778.<sup>[26]</sup>

In other European countries, the Charité (a hospital founded in 1710) in Berlin established a separate Pediatric Pavilion in 1830, followed by similar institutions at Saint Petersburg in 1834, and at Vienna and Breslau (now Wrocław, Poland), both in 1837. In 1852 Britain's first pediatric hospital, the Hospital for Sick Children, Great Ormond Street was founded by Charles West.<sup>[24]</sup> The first Children's hospital in Scotland opened in 1860 in Edinburgh.<sup>[27]</sup> In the US, the first similar institutions were the Children's Hospital of Philadelphia, which opened in 1855, and then Boston Children's Hospital (1869).<sup>[28]</sup> Subspecialties in pediatrics were created at the Harriet Lane Home at Johns Hopkins by Edwards A. Park.<sup>[29]</sup>



## Differences between adult and pediatric medicine

[edit]

The body size differences are paralleled by maturation changes. The smaller body of an infant or neonate is substantially different physiologically from that of an adult. Congenital defects, genetic variance, and developmental issues are of greater concern to pediatricians than they often are to adult physicians. A common adage is that children are not simply "little adults". The clinician must take into account the immature physiology of the infant or child when considering symptoms, prescribing medications, and diagnosing illnesses.<sup>[30]</sup>

Pediatric physiology directly impacts the pharmacokinetic properties of drugs that enter the body. The absorption, distribution, metabolism, and elimination of medications differ between developing children and grown adults.<sup>[30][31][32]</sup> Despite completed studies and reviews, continual research is needed to better understand how these factors should affect the decisions of healthcare providers when prescribing and administering medications to the pediatric population.<sup>[30]</sup>

### Absorption

[edit]

Many drug absorption differences between pediatric and adult populations revolve around the stomach. Neonates and young infants have increased stomach pH due to decreased acid secretion, thereby creating a more basic environment for drugs that are taken by mouth.<sup>[31][30][32]</sup> Acid is essential to degrading certain oral drugs before systemic absorption. Therefore, the absorption of these drugs in children is greater than in adults due to decreased breakdown and increased preservation in a less acidic gastric space.<sup>[31]</sup>

Children also have an extended rate of gastric emptying, which slows the rate of drug absorption.<sup>[31][32]</sup>

Drug absorption also depends on specific enzymes that come in contact with the oral drug as it travels through the body. Supply of these enzymes increase as children continue to develop their gastrointestinal tract.<sup>[31][32]</sup> Pediatric patients have underdeveloped proteins, which leads to decreased metabolism and increased serum concentrations of specific drugs. However, prodrugs experience the opposite effect because enzymes are necessary for allowing their active form to enter systemic circulation.<sup>[31]</sup>

## **Distribution**

[edit]

Percentage of total body water and extracellular fluid volume both decrease as children grow and develop with time. Pediatric patients thus have a larger volume of distribution than adults, which directly affects the dosing of hydrophilic drugs such as beta-lactam antibiotics like ampicillin.<sup>[31]</sup> Thus, these drugs are administered at greater weight-based doses or with adjusted dosing intervals in children to account for this key difference in body composition.<sup>[31][30]</sup>

Infants and neonates also have fewer plasma proteins. Thus, highly protein-bound drugs have fewer opportunities for protein binding, leading to increased distribution.<sup>[30]</sup>

## **Metabolism**

[edit]

Drug metabolism primarily occurs via enzymes in the liver and can vary according to which specific enzymes are affected in a specific stage of development.<sup>[31]</sup> Phase I and Phase II enzymes have different rates of maturation and development, depending on their specific mechanism of action (i.e. oxidation, hydrolysis, acetylation, methylation, etc.). Enzyme capacity, clearance, and half-life are all

factors that contribute to metabolism differences between children and adults<sup>[31]</sup><sup>[32]</sup> Drug metabolism can even differ within the pediatric population, separating neonates and infants from young children.<sup>[30]</sup>

## **Elimination**

[edit]

Drug elimination is primarily facilitated via the liver and kidneys<sup>[31]</sup> In infants and young children, the larger relative size of their kidneys leads to increased renal clearance of medications that are eliminated through urine<sup>[32]</sup> In preterm neonates and infants, their kidneys are slower to mature and thus are unable to clear as much drug as fully developed kidneys. This can cause unwanted drug build-up, which is why it is important to consider lower doses and greater dosing intervals for this population.<sup>[30]</sup><sup>[31]</sup> Diseases that negatively affect kidney function can also have the same effect and thus warrant similar considerations.<sup>[31]</sup>

## **Pediatric autonomy in healthcare**

[edit]

A major difference between the practice of pediatric and adult medicine is that children, in most jurisdictions and with certain exceptions, cannot make decisions for themselves. The issues of guardianship, privacy, legal responsibility, and informed consent must always be considered in every pediatric procedure. Pediatricians often have to treat the parents and sometimes, the family, rather than just the child. Adolescents are in their own legal class, having rights to their own health care decisions in certain circumstances. The concept of legal consent combined with the non-legal consent (assent) of the child when considering treatment options, especially in the face of conditions with poor prognosis or complicated and painful procedures/surgeries, means the pediatrician must take into account the desires of many people, in addition to those of the patient.<sup>[citation needed]</sup>

## History of pediatric autonomy

[edit]

The term autonomy is traceable to ethical theory and law, where it states that autonomous individuals can make decisions based on their own logic.<sup>[33]</sup>

Hippocrates was the first to use the term in a medical setting. He created a code of ethics for doctors called the *Hippocratic Oath* that highlighted the importance of putting patients' interests first, making autonomy for patients a top priority in health care.<sup>[34]</sup>

In ancient times, society did not view pediatric medicine as essential or scientific.<sup>[35]</sup> Experts considered professional medicine unsuitable for treating children. Children also had no rights. Fathers regarded their children as property, so their children's health decisions were entrusted to them.<sup>[5]</sup> As a result, mothers, midwives, "wise women", and general practitioners treated the children instead of doctors.<sup>[35]</sup> Since mothers could not rely on professional medicine to take care of their children, they developed their own methods, such as using alkaline soda ash to remove the vernix at birth and treating teething pain with opium or wine. The absence of proper pediatric care, rights, and laws in health care to prioritize children's health led to many of their deaths. Ancient Greeks and Romans sometimes even killed healthy female babies and infants with deformities since they had no adequate medical treatment and no laws prohibiting infanticide.<sup>[5]</sup>

In the twentieth century, medical experts began to put more emphasis on children's rights. In 1989, in the United Nations Rights of the Child Convention, medical experts developed the Best Interest Standard of Child to prioritize children's rights and best interests. This event marked the onset of pediatric autonomy. In 1995, the American Academy of Pediatrics (AAP) finally acknowledged the Best Interest Standard of a Child as an ethical principle for pediatric decision-making, and it is still being used today.<sup>[34]</sup>

## Parental authority and current medical issues

[edit]

The majority of the time, parents have the authority to decide what happens to their child. Philosopher John Locke argued that it is the responsibility of parents to raise their children and that God gave them this authority. In modern society, Jeffrey Blustein, modern philosopher and author of the book *Parents and Children: The Ethics of Family*, argues that parental authority is granted because the child requires parents to satisfy their needs. He believes that parental autonomy is more about parents providing good care for their children and treating them with respect than parents having rights.<sup>[36]</sup> The researcher Kyriakos Martakis, MD, MSc, explains that research shows parental influence negatively affects children's ability to form autonomy. However, involving children in the decision-making process allows children to develop their cognitive skills and create their own opinions and, thus, decisions about their health. Parental authority affects the degree of autonomy the child patient has. As a result, in Argentina, the new National Civil and Commercial Code has enacted various changes to the healthcare system to encourage children and adolescents to develop autonomy. It has become more crucial to let children take accountability for their own health decisions.<sup>[37]</sup>

In most cases, the pediatrician, parent, and child work as a team to make the best possible medical decision. The pediatrician has the right to intervene for the child's welfare and seek advice from an ethics committee. However, in recent studies, authors have denied that complete autonomy is present in pediatric healthcare. The same moral standards should apply to children as they do to adults. In support of this idea is the concept of paternalism, which negates autonomy when it is in the patient's interests. This concept aims to keep the child's best interests in mind regarding autonomy. Pediatricians can interact with patients and help them make decisions that will benefit them, thus enhancing their autonomy. However, radical theories that question a child's moral worth continue to be debated today<sup>[37]</sup>

Authors often question whether the treatment and equality of a child and an adult should be the same. Author Tamar Schapiro notes that children need nurturing and cannot exercise the same level of authority as adults.<sup>[38]</sup> Hence, continuing the discussion on whether children are capable of making important health decisions until this day.

## **Modern advancements**


[edit]

According to the Subcommittee of Clinical Ethics of the Argentinean Pediatric Society (SAP), children can understand moral feelings at all ages and can make reasonable decisions based on those feelings. Therefore, children and teens are deemed capable of making their own health decisions when they reach the age of 13. Recently, studies made on the decision-making of children have challenged that age to be 12.<sup>[37]</sup>

Technology has made several modern advancements that contribute to the future development of child autonomy, for example, unsolicited findings (U.F.s) of pediatric exome sequencing. They are findings based on pediatric exome sequencing that explain in greater detail the intellectual disability of a child and predict to what extent it will affect the child in the future. Genetic and intellectual disorders in children make them incapable of making moral decisions, so people look down upon this kind of testing because the child's future autonomy is at risk. It is still in question whether parents should request these types of testing for their children. Medical experts argue that it could endanger the autonomous rights the child will possess in the future. However, the parents contend that genetic testing would benefit the welfare of their children since it would allow them to make better health care decisions.<sup>[39]</sup> Exome sequencing for children and the decision to grant parents the right to request them is a medically ethical issue that many still debate today.

## **Education requirements**

[edit]

The examples and perspective in this section **deal primarily with United States and do not represent a worldwide view of the subject**. You may  improve this section, discuss the issue on the talk page, or create a new section, as appropriate. *(September 2019) (Learn how and when to remove this message)*

Aspiring medical students will need 4 years of undergraduate courses at a college or university, which will get them a BS, BA or other bachelor's degree. After completing college, future pediatricians will need to attend 4 years of medical school (MD/DO/MBBS) and later do 3 more years of residency training, the first year of which is called "internship." After completing the 3 years of residency, physicians are eligible to become certified in pediatrics by passing a rigorous test that deals with medical conditions related to young children.<sup>[*citation needed*]</sup>

In high school, future pediatricians are required to take basic science classes such as biology, chemistry, physics, algebra, geometry, and calculus. It is also advisable to learn a foreign language (preferably Spanish in the United States) and be involved in high school organizations and extracurricular activities. After high school, college students simply need to fulfill the basic science course requirements that most medical schools recommend and will need to prepare to take the MCAT (Medical College Admission Test) in their junior or early senior year in college. Once attending medical school, student courses will focus on basic medical sciences like human anatomy, physiology, chemistry, etc., for the first three years, the second year of which is when medical students start to get hands-on experience with actual patients.<sup>[40]</sup>

## Training of pediatricians

[edit]

Pediatrics

image not found or type unknown



## Occupation

### Names

- Pediatrician
- Paediatrician

**Occupation type** Specialty

**Activity sectors** Medicine

## Description

### Education required

- Doctor of Medicine
- Doctor of Osteopathic Medicine
- Bachelor of Medicine, Bachelor of Surgery (MBBS/MBChB)

### Fields of employment

Hospitals, Clinics

The training of pediatricians varies considerably across the world. Depending on jurisdiction and university, a medical degree course may be either undergraduate-entry or graduate-entry. The former commonly takes five or six years and has been usual in the Commonwealth. Entrants to graduate-entry courses (as in the US), usually lasting four or five years, have previously completed a three- or four-year university degree, commonly but by no means always in sciences. Medical graduates hold a degree specific to the country and university in and from which they graduated. This degree qualifies that medical practitioner to become licensed or registered under the laws of that particular country, and sometimes of several countries, subject to requirements for "internship" or "conditional registration".



Pediatricians must undertake further training in their chosen field. This may take from four to eleven or more years depending on jurisdiction and the degree of specialization.

In the United States, a medical school graduate wishing to specialize in pediatrics must undergo a three-year residency composed of outpatient, inpatient, and critical care rotations. Subspecialties within pediatrics require further training in the form of 3-year fellowships. Subspecialties include critical care, gastroenterology, neurology, infectious disease, hematology/oncology, rheumatology, pulmonology, child abuse, emergency medicine, endocrinology, neonatology, and others.<sup>[41]</sup>

In most jurisdictions, entry-level degrees are common to all branches of the medical profession, but in some jurisdictions, specialization in pediatrics may begin before completion of this degree. In some jurisdictions, pediatric training is begun immediately following the completion of entry-level training. In other jurisdictions, junior medical doctors must undertake generalist (unstreamed) training for a number of years before commencing pediatric (or any other) specialization. Specialist training is often largely under the control of *pediatric organizations* (see below) rather than universities and depends on the jurisdiction.

## **Subspecialties**

[edit]

Subspecialties of pediatrics include:

*(not an exhaustive list)*

- Addiction medicine (multidisciplinary)
- Adolescent medicine
- Child abuse pediatrics
- Clinical genetics
- Clinical informatics
- Developmental-behavioral pediatrics

- Headache medicine
- Hospital medicine
- Medical toxicology
- Metabolic medicine
- Neonatology/Perinatology
- Pain medicine (multidisciplinary)
- Palliative care (multidisciplinary)
- Pediatric allergy and immunology
- Pediatric cardiology
  - Pediatric cardiac critical care
- Pediatric critical care
  - Neurocritical care
  - Pediatric cardiac critical care
- Pediatric emergency medicine
- Pediatric endocrinology
- Pediatric gastroenterology
  - Transplant hepatology
- Pediatric hematology
- Pediatric infectious disease
- Pediatric nephrology
- Pediatric oncology
  - Pediatric neuro-oncology
- Pediatric pulmonology
- Primary care
- Pediatric rheumatology
- Sleep medicine (multidisciplinary)
- Social pediatrics
- Sports medicine

### **Other specialties that care for children**

[edit]

*(not an exhaustive list)*

- Child neurology
  - Addiction medicine (multidisciplinary)
  - Brain injury medicine
  - Clinical neurophysiology
  - Epilepsy
  - Headache medicine
  - Neurocritical care
  - Neuroimmunology
  - Neuromuscular medicine
  - Pain medicine (multidisciplinary)
  - Palliative care (multidisciplinary)
  - Pediatric neuro-oncology
  - Sleep medicine (multidisciplinary)
- Child and adolescent psychiatry, subspecialty of psychiatry
- Neurodevelopmental disabilities
- Pediatric anesthesiology, subspecialty of anesthesiology
- Pediatric dentistry, subspecialty of dentistry
- Pediatric dermatology, subspecialty of dermatology
- Pediatric gynecology
- Pediatric neurosurgery, subspecialty of neurosurgery
- Pediatric ophthalmology, subspecialty of ophthalmology
- Pediatric orthopedic surgery, subspecialty of orthopedic surgery
- Pediatric otolaryngology, subspecialty of otolaryngology
- Pediatric plastic surgery, subspecialty of plastic surgery
- Pediatric radiology, subspecialty of radiology
- Pediatric rehabilitation medicine, subspecialty of physical medicine and rehabilitation
- Pediatric surgery, subspecialty of general surgery
- Pediatric urology, subspecialty of urology

## See also

[edit]

- American Academy of Pediatrics
- American Osteopathic Board of Pediatrics
- Center on Media and Child Health (CMCH)
- Children's hospital
- List of pediatric organizations
- List of pediatrics journals
- Medical specialty
- Pediatric Oncall
- Pain in babies
- Royal College of Paediatrics and Child Health
- Pediatric environmental health

## References

[edit]

1. <sup>^</sup> "Paediatrics" (PDF). *nhs.uk*. Archived (PDF) from the original on 13 July 2020. Retrieved 2 July 2020.
2. <sup>^</sup> "Choosing a Pediatrician for Your New Baby (for Parents) - Nemours KidsHealth". *kidshealth.org*. Archived from the original on 14 July 2020. Retrieved 13 July 2020.
3. <sup>^</sup> "Age limits of pediatrics". *Pediatrics*. **81** (5): 736. May 1988. doi:10.1542/peds.81.5.736. PMID 3357740. S2CID 245164191. Archived from the original on 19 April 2017. Retrieved 18 April 2017.
4. <sup>^</sup> Sawyer, Susan M.; McNeil, Robyn; Francis, Kate L.; Matskarofski, Juliet Z.; Patton, George C.; Bhutta, Zulfiqar A.; Esangbedo, Dorothy O.; Klein, Jonathan D. (1 November 2019). "The age of paediatrics". *The Lancet Child & Adolescent Health*. **3** (11): 822–830. doi:10.1016/S2352-4642(19)30266-4. ISSN 2352-4642. PMID 31542355. S2CID 202732818.
5. <sup>^</sup> **a b c d e** Duffin, Jacalyn (2010). *History of Medicine, Second Edition: A Scandalously Short Introduction*. University of Toronto Press.
6. <sup>^</sup> **a b c d e** Colón, A. R.; Colón, P. A. (January 1999). *Nurturing children: a history of pediatrics*. Greenwood Press. ISBN 978-0-313-31080-5. Retrieved 20 October 2012.

7. ^ Celsus, *De Medicina*, Book 3, Chapter 7, § 1.
8. ^ John G. Raffensperger. *Children's Surgery: A Worldwide History*. McFarland. p. 21.
9. ^ David Levinson; Karen Christensen. *Encyclopedia of modern Asia*. Vol. 4. Charles Scribner's Sons. p. 116.
10. ^ Desai, A.B. *Textbook Of Paediatrics*. Orient blackswan. p. 1.
11. ^ Dunn, P. M. (1995). "Soranus of Ephesus (Circa AD 98-138) and perinatal care in Roman times". *Archives of Disease in Childhood. Fetal and Neonatal Edition*. **73** (1): F51 – F52. doi:10.1136/fn.73.1.f51. PMC 2528358. PMID 7552600.
12. ^ Elgood, Cyril (2010). *A Medical History of Persia and The Eastern Caliphate (1st ed.)*. London: Cambridge. pp. 202–203. ISBN 978-1-108-01588-2. "By writing a monograph on 'Diseases in Children' he may also be looked upon as the father of paediatrics."
13. ^ U.S. National Library of Medicine, "Islamic Culture and the Medical Arts, Al-Razi, the Clinician" [1] Archived 5 January 2018 at the Wayback Machine
14. ^ "Achar S *Textbook Of Pediatrics (Third Edition)*". A. B. Desai (ed.) (1989). p.1. ISBN 81-250-0440-8
15. ^ Stern, Alexandra Minna; Markel, Howard (2002). *Formative Years: Children's Health in the United States, 1880-2000*. University of Michigan Press. pp. 23–24. doi:10.3998/mpub.17065. ISBN 978-0-472-02503-9. Archived from the original on 30 November 2021. Retrieved 30 November 2021.
16. ^ Lock, Stephen; John M. Last; George Dunea (2001). *The Oxford illustrated companion to medicine*. Oxford University Press US. p. 173. ISBN 978-0-19-262950-0. Retrieved 9 July 2010. "Rosen von Rosenstein."
17. ^ Roberts, Michael (2003). *The Age of Liberty: Sweden 1719–1772*. Cambridge University Press. p. 216. ISBN 978-0-521-52707-1. Retrieved 9 July 2010.
18. ^ Dallas, John. "Classics of Child Care". Royal College of Physicians of Edinburgh. Archived from the original on 27 July 2011. Retrieved 9 July 2010.
19. ^ Duffin, Jacalyn (29 May 2010). *History of Medicine, Second Edition: A Scandalously Short Introduction*. University of Toronto Press.
20. ^ Stern, Alexandra Minna; Markel, Howard (2002). *Formative Years: Children's Health in the United States, 1880-2000*. University of Michigan Press. pp. 23–24. doi:10.3998/mpub.17065. ISBN 978-0-472-02503-9. Archived from the original on 30 November 2021. Retrieved 30 November 2021.
21. ^ "Broadribb's *Introductory Pediatric Nursing*". Nancy T. Hatfield (2007). p.4. ISBN 0-7817-7706-2

22. ^ "Jacobi Medical Center - General Information". Archived from the original on 18 April 2006. Retrieved 6 April 2006.
23. ^ Kutzsche, Stefan (8 April 2021). "Abraham Jacobi (1830–1919) and his transition from political to medical activist". *Acta Paediatrica*. **110** (8): 2303–2305. doi:10.1111/apa.15887. ISSN 0803-5253. PMID 33963612. S2CID 233998658. Archived from the original on 7 May 2023. Retrieved 7 May 2023.
24. ^ **a b** Ballbriga, Angel (1991). "One century of pediatrics in Europe (section: development of pediatric hospitals in Europe)". In Nichols, Burford L.; et al. (eds.). *History of Paediatrics 1850–1950*. Nestlé Nutrition Workshop Series. Vol. 22. New York: Raven Press. pp. 6–8. ISBN 0-88167-695-0.
25. ^ official history site (in French) of nineteenth century paediatric hospitals in Paris
26. ^ "Introducing the Necker-Enfants Malades Hospital". *Hôpital des Necker-Enfants Malades*.
27. ^ Young, D.G. (August 1999). "The Mason Brown Lecture: Scots and paediatric surgery". *Journal of the Royal College of Surgeons Edinburgh*. **44** (4): 211–5. PMID 10453141. Archived from the original on 14 July 2014.
28. ^ Pearson, Howard A. (1991). "Pediatrics in the United States". In Nichols, Burford L.; et al. (eds.). *History of Paediatrics 1850–1950*. Nestlé Nutrition Workshop Series. Vol. 22. New York: Raven Press. pp. 55–63. ISBN 0-88167-695-0.
29. ^ "Commentaries: Edwards A Park". *Pediatrics*. **44** (6). American Academy of Pediatrics: 897–901. 1969. doi:10.1542/peds.44.6.897. PMID 4903838. S2CID 43298798.
30. ^ **a b c d e f g h** O'Hara, Kate (2016). "Paediatric pharmacokinetics and drug doses". *Australian Prescriber*. **39** (6): 208–210. doi:10.18773/austprescr.2016.071. ISSN 0312-8008. PMC 5155058. PMID 27990048.
31. ^ **a b c d e f g h i j k l m** Wagner, Jonathan; Abdel-Rahman, Susan M. (2013). "Pediatric pharmacokinetics". *Pediatrics in Review*. **34** (6): 258–269. doi:10.1542/pir.34-6-258. ISSN 1526-3347. PMID 23729775.
32. ^ **a b c d e f** Batchelor, Hannah Katharine; Marriott, John Francis (2015). "Paediatric pharmacokinetics: key considerations". *British Journal of Clinical Pharmacology*. **79** (3): 395–404. doi:10.1111/bcp.12267. ISSN 1365-2125. PMC 4345950. PMID 25855821.
33. ^ Katz, Aviva L.; Webb, Sally A.; COMMITTEE ON BIOETHICS; Macauley, Robert C.; Mercurio, Mark R.; Moon, Margaret R.; Okun, Alexander L.; Opel, Douglas J.; Statter, Mindy B. (1 August 2016). "Informed Consent in Decision-Making in Pediatric Practice". *Pediatrics*. **138** (2): e20161485. doi:10.1542/peds.2016-1485. ISSN 0031-4005. PMID 27456510. S2CID 7951515.

34. ^ **a b** Mazur, Kate A.; Berg, Stacey L., eds. (2020). *Ethical Issues in Pediatric Hematology/Oncology*. pp. 13–21. doi:10.1007/978-3-030-22684-8. ISBN 978-3-030-22683-1. S2CID 208302429.
35. ^ **a b** Stern, Alexandra Minna; Markel, Howard (2002). *Formative Years: Children's Health in the United States, 1880-2000*. University of Michigan Press. pp. 23–24. doi:10.3998/mpub.17065. ISBN 978-0-472-02503-9. Archived from the original on 30 November 2021. Retrieved 30 November 2021.
36. ^ Friedman, Lainie Ross (2004). *Children, families, and health care decision making*. Clarendon Press. ISBN 0-19-925154-1. OCLC 756393117.
37. ^ **a b c** Martakis, K.; Schröder-Bäck, P.; Brand, H. (1 June 2018). "Developing child autonomy in pediatric healthcare: towards an ethical model". *Archivos Argentinos de Pediatría*. **116** (3): e401 – e408. doi:10.5546/aap.2018.eng.e401. ISSN 0325-0075. PMID 29756714. S2CID 46889502.
38. ^ Schapiro, Tamar (1 July 1999). "What Is a Child?". *Ethics*. **109** (4): 715–738. doi:10.1086/233943. ISSN 0014-1704. S2CID 170129444. Archived from the original on 30 November 2021. Retrieved 30 November 2021.
39. ^ Dondorp, W.; Bolt, I.; Tibben, A.; De Wert, G.; Van Summeren, M. (1 September 2021). "'We Should View Him as an Individual': The Role of the Child's Future Autonomy in Shared Decision-Making About Unsolicited Findings in Pediatric Exome Sequencing". *Health Care Analysis*. **29** (3): 249–261. doi:10.1007/s10728-020-00425-7. ISSN 1573-3394. PMID 33389383. S2CID 230112761.
40. ^ "What Education Is Required to Be a Pediatrician?". Archived from the original on 7 June 2017. Retrieved 14 June 2017.
41. ^ "CoPS". *www.pedsubs.org*. Archived from the original on 18 September 2013. Retrieved 14 August 2015.


## Further reading

[edit]


- *BMC Pediatrics* – open access
- *Clinical Pediatrics*
- *Developmental Review* – partial open access
- *JAMA Pediatrics*
- *The Journal of Pediatrics* – partial open access

## External links


[edit]



Wikimedia Commons has media related to ***Pediatrics***.



Wikibooks has a book on the topic of: ***Pediatrics***



Look up ***paediatrics*** or ***pediatrics*** in Wiktionary, the free dictionary.

- [Pediatrics Directory at Curlie](#)
- [Pediatric Health Directory at OpenMD](#)
- [v](#)
- [t](#)
- [e](#)

Medicine



## **Surgery**

- Cardiac surgery
- Cardiothoracic surgery
- Endocrine surgery
- Eye surgery
- General surgery
  - Colorectal surgery
  - Digestive system surgery
- Neurosurgery
- Oral and maxillofacial surgery
- Orthopedic surgery
- Hand surgery
- Otolaryngology
  - ENT
- Pediatric surgery
- Plastic surgery
- Reproductive surgery
- Surgical oncology
- Transplant surgery
- Trauma surgery
- Urology
  - Andrology
- Vascular surgery
- Allergy / Immunology
- Angiology
- Cardiology
- Endocrinology
- Gastroenterology
  - Hepatology





## **Internal medicine**

- Geriatrics
- Hematology
- Hospital medicine
- Infectious diseases
- Nephrology
- Oncology

## Medical education

- Medical school
- Bachelor of Medicine, Bachelor of Surgery
- Bachelor of Medical Sciences
- Master of Medicine
- Master of Surgery
- Doctor of Medicine
- Doctor of Osteopathic Medicine
- MD–PhD
  - Medical Scientist Training Program
- Alternative medicine
- Allied health
- Molecular oncology
- Nanomedicine
- Personalized medicine
- Public health
- Rural health
- Therapy
- Traditional medicine
- Veterinary medicine
- Physician
  - Chief physician
- History of medicine

## Related topics

-  **Category**
-  **Commons**
-  **WikiProject**
-  **Portal**
-  **Outline**

- **v**
- **t**
- **e**

### Health (Pediatrics)

- Baby food
- Birth weight
- Breast pump
- Breastfeeding
- Breastfeeding and medications
- Breastfeeding and mental health
- Bottle feeding
- Colic
- Cradle cap
- Esotropia
- Failure to thrive
- Immunization
- Infant and toddler safety
- Infant bathing
- Infant feeding
- Infant food safety
- Infant formula
- Infant massage
- Infant respiratory distress syndrome
- Infant sleep training
- Neonatal intensive care unit
- Newborn care and safety
- Oral rehydration therapy
  - Pedialyte
- Preterm birth
- Shaken baby syndrome
- Soy formula
- SIDS

## **Development**

- Attachment parenting
- Baby-led weaning
- Baby talk
- Babbling
- Birth defect
- Childbirth
- Crawling
- Gestational age
- Infant visual development
- Irritant diaper dermatitis
- Infant cognitive development
- Infant crying
- Kangaroo care
- Mother
- Nursery rhyme
- Object permanence
- Parent
- Parenting
- Peekaboo
- Play
- Prenatal development
- Prenatal development table
- Teething
- Walking
- Weaning

## **Socialization and Culture**

- Attachment
- Babysitting
- Child abuse
- Child care
- Child custody
- Children's rights
  - UN Child rights
- Circumcision
- Foster care
- Grandparent visitation
- Infant swimming
- Milk bank
- Nanny
- Wet nurse

## **Infant care and equipment**

- Baby bouncer
- Baby gate
- Baby monitor/Hidden camera
- Baby powder
- Baby shampoo
- Baby toy
- Baby walker
- Bib
- Baby swing
- Baby transport
- Bassinet
- Car seat safety
- Cloth diaper
- Cradle board
- Diaper
- Diaper bag
- Baby wipes
- Haberman Feeder
- High chair
- Infant bed (*American 'crib' and 'cradle', British 'cot'*)
- Infant carrier
- Infant clothing
- Pacifier
- Playpen
- Stroller
- Supplemental nursing system
- Swaddling
- Swim diaper
- Teether
- Travel cot

## Other topics

- Baby shower
- Babywearing
- Child neglect
- Closed adoption
- Cry room
- Infant ear piercing
- Open adoption
- Prenatal cocaine exposure
- Neonatal withdrawal syndrome
- Parental child abduction
- Parental responsibility
- Parenting plan
- Paternity
  - Paternity fraud

## Authority control databases Image not found or type unknown Edit this at Wikidata

- Germany
- Japan
- National**
  - Czech Republic
  - Latvia
  - Israel
- Other**
  - NARA


## About health professional

A **health professional**, **healthcare professional**, or **healthcare worker** (sometimes abbreviated **HCW**)<sup>[1]</sup> is a provider of health care treatment and advice based on formal training and experience. The field includes those who work as a nurse,


physician (such as family physician, internist, obstetrician, psychiatrist, radiologist, surgeon etc.), physician assistant, registered dietitian, veterinarian, veterinary technician, optometrist, pharmacist, pharmacy technician, medical assistant, physical therapist, occupational therapist, dentist, midwife, psychologist, audiologist, or healthcare scientist, or who perform services in allied health professions. Experts in public health and community health are also health professionals.

## Fields

[edit]



NY College of Health Professions massage therapy class




US Navy doctors deliver a healthy baby

- v
- t
- e

**Health practitioners  
and professionals**



- Athletic trainer
- Audiologist
- Chiropractor
- Clinical coder
- Clinical nurse specialist
- Clinical officer
- Community health worker
- Dentist
- Dietitian and nutritionist
- Emergency medical technician
- Feldsher
- Health administrator
- Medical assistant
- Medical laboratory scientist
- Medical transcriptionist
- Nurse anesthetist
- Nurse practitioner
- Nurse midwife
- Nurse
- Occupational Therapist
- Optometrist
- Paramedic
- Pharmacist
- Pharmaconomist
- Pharmacy technician
- Phlebotomist
- Physician
- Physician assistant
- Podiatrist
- Psychologist
- Psychotherapist
- Physical therapist
- Radiographer
- Radiology assistant



70% of global health and social care workers are women, 30% of leaders in the global health sector are women

The healthcare workforce comprises a wide variety of professions and occupations who provide some type of healthcare service, including such direct care practitioners as physicians, nurse practitioners, physician assistants, nurses, respiratory therapists, dentists, pharmacists, speech-language pathologist, physical therapists, occupational therapists, physical and behavior therapists, as well as allied health professionals such as phlebotomists, medical laboratory scientists, dieticians, and social workers. They often work in hospitals, healthcare centers and other service delivery points, but also in academic training, research, and administration. Some provide care and treatment services for patients in private homes. Many countries have a large number of community health workers who work outside formal healthcare institutions. Managers of healthcare services, health information technicians, and other assistive personnel and support workers are also considered a vital part of health care teams.<sup>[2]</sup>

Healthcare practitioners are commonly grouped into health professions. Within each field of expertise, practitioners are often classified according to skill level and skill specialization. "Health professionals" are highly skilled workers, in professions that usually require extensive knowledge including university-level study leading to the award of a first degree or higher qualification.<sup>[3]</sup> This category includes physicians,

physician assistants, registered nurses, veterinarians, veterinary technicians, veterinary assistants, dentists, midwives, radiographers, pharmacists, physiotherapists, optometrists, operating department practitioners and others. Allied health professionals, also referred to as "health associate professionals" in the International Standard Classification of Occupations, support implementation of health care, treatment and referral plans usually established by medical, nursing, respiratory care, and other health professionals, and usually require formal qualifications to practice their profession. In addition, unlicensed assistive personnel assist with providing health care services as permitted.<sup>[citation needed]</sup>

Another way to categorize healthcare practitioners is according to the sub-field in which they practice, such as mental health care, pregnancy and childbirth care, surgical care, rehabilitation care, or public health.<sup>[citation needed]</sup>

## **Mental health**

[edit]

Main article: Mental health professional

A mental health professional is a health worker who offers services to improve the mental health of individuals or treat mental illness. These include psychiatrists, psychiatry physician assistants, clinical, counseling, and school psychologists, occupational therapists, clinical social workers, psychiatric-mental health nurse practitioners, marriage and family therapists, mental health counselors, as well as other health professionals and allied health professions. These health care providers often deal with the same illnesses, disorders, conditions, and issues; however, their scope of practice often differs. The most significant difference across categories of mental health practitioners is education and training.<sup>[4]</sup> There are many damaging effects to the health care workers. Many have had diverse negative psychological symptoms ranging from emotional trauma to very severe anxiety. Health care workers have not been treated right and because of that their mental, physical, and emotional health has been affected by it. The SAGE author's said that there were

94% of nurses that had experienced at least one PTSD after the traumatic experience. Others have experienced nightmares, flashbacks, and short and long term emotional reactions.<sup>[5]</sup> The abuse is causing detrimental effects on these health care workers. Violence is causing health care workers to have a negative attitude toward work tasks and patients, and because of that they are "feeling pressured to accept the order, dispense a product, or administer a medication"<sup>[6]</sup> Sometimes it can range from verbal to sexual to physical harassment, whether the abuser is a patient, patient's families, physician, supervisors, or nurses.<sup>[citation needed]</sup>

## **Obstetrics**

[edit]

Main articles: Obstetrics, Midwifery, and Birth attendant

A maternal and newborn health practitioner is a health care expert who deals with the care of women and their children before, during and after pregnancy and childbirth. Such health practitioners include obstetricians, physician assistants, midwives, obstetrical nurses and many others. One of the main differences between these professions is in the training and authority to provide surgical services and other life-saving interventions.<sup>[7]</sup> In some developing countries, traditional birth attendants, or traditional midwives, are the primary source of pregnancy and childbirth care for many women and families, although they are not certified or licensed. According to research, rates for unhappiness among obstetrician-gynecologists (Ob-Gyns) range somewhere between 40 and 75 percent<sup>[8]</sup>

## **Geriatrics**

[edit]

Main articles: Geriatrics and Geriatric care management

A geriatric care practitioner plans and coordinates the care of the elderly and/or disabled to promote their health, improve their quality of life, and maintain their

independence for as long as possible.<sup>[9]</sup> They include geriatricians, occupational therapists, physician assistants, adult-gerontology nurse practitioners, clinical nurse specialists, geriatric clinical pharmacists, geriatric nurses, geriatric care managers, geriatric aides, nursing aides, caregivers and others who focus on the health and psychological care needs of older adults.<sup>[citation needed]</sup>

## **Surgery**

[edit]

A surgical practitioner is a healthcare professional and expert who specializes in the planning and delivery of a patient's perioperative care, including during the anaesthetic, surgical and recovery stages. They may include general and specialist surgeons, physician assistants, assistant surgeons, surgical assistants, veterinary surgeons, veterinary technicians, anesthesiologists, anesthesiologist assistants, nurse anesthetists, surgical nurses, clinical officers, operating department practitioners, anaesthetic technicians, perioperative nurses, surgical technologists, and others.<sup>[citation needed]</sup>

## **Rehabilitation**

[edit]

A rehabilitation care practitioner is a health worker who provides care and treatment which aims to enhance and restore functional ability and quality of life to those with physical impairments or disabilities. These include physiatrists, physician assistants, rehabilitation nurses, clinical nurse specialists, nurse practitioners, physiotherapists, chiropractors, orthotists, prosthetists, occupational therapists, recreational therapists, audiologists, speech and language pathologists, respiratory therapists, rehabilitation counsellors, physical rehabilitation therapists, athletic trainers, physiotherapy technicians, orthotic technicians, prosthetic technicians, personal care assistants, and others.<sup>[10]</sup>

## Optometry

[edit]

Main article: Optometry

Optometry is a field traditionally associated with the correction of refractive errors using glasses or contact lenses, and treating eye diseases. Optometrists also provide general eye care, including screening exams for glaucoma and diabetic retinopathy and management of routine or eye conditions. Optometrists may also undergo further training in order to specialize in various fields, including glaucoma, medical retina, low vision, or paediatrics. In some countries, such as the United Kingdom, United States, and Canada, Optometrists may also undergo further training in order to be able to perform some surgical procedures.

## Diagnostics


[edit]

Main article: Medical diagnosis

Medical diagnosis providers are health workers responsible for the process of determining which disease or condition explains a person's symptoms and signs. It is most often referred to as diagnosis with the medical context being implicit. This usually involves a team of healthcare providers in various diagnostic units. These include radiographers, radiologists, Sonographers, medical laboratory scientists, pathologists, and related professionals.<sup>*[citation needed]*</sup>

## Dentistry

[edit]



Dental assistant on the right supporting a dental operator on the left, during a procedure.

Main article: Dentistry

A dental care practitioner is a health worker and expert who provides care and treatment to promote and restore oral health. These include dentists and dental surgeons, dental assistants, dental auxiliaries, dental hygienists, dental nurses, dental technicians, dental therapists or oral health therapists, and related professionals.

## **Podiatry**

[edit]

Care and treatment for the foot, ankle, and lower leg may be delivered by podiatrists, chiropodists, pedorthists, foot health practitioners, podiatric medical assistants, podiatric nurse and others.

## **Public health**

[edit]

A public health practitioner focuses on improving health among individuals, families and communities through the prevention and treatment of diseases and injuries, surveillance of cases, and promotion of healthy behaviors. This category includes community and preventive medicine specialists, physician assistants, public health nurses, pharmacist, clinical nurse specialists, dietitians, environmental health officers (public health inspectors), paramedics, epidemiologists, public health dentists, and

others.<sup>[*citation needed*]</sup>

## Alternative medicine


[edit]

In many societies, practitioners of alternative medicine have contact with a significant number of people, either as integrated within or remaining outside the formal health care system. These include practitioners in acupuncture, Ayurveda, herbalism, homeopathy, naturopathy, Reiki, Shamballa Reiki energy healing Archived 2021-01-25 at the Wayback Machine, Siddha medicine, traditional Chinese medicine, traditional Korean medicine, Unani, and Yoga. In some countries such as Canada, chiropractors and osteopaths (not to be confused with doctors of osteopathic medicine in the United States) are considered alternative medicine practitioners.

## Occupational hazards

[edit]

See also: Occupational hazards in dentistry and Nursing § Occupational hazards



A healthcare professional wears an air sampling device to investigate exposure to airborne influenza

A video describing the Occupational Health and Safety Network, a tool for monitoring occupational hazards to health care workers

The healthcare workforce faces unique health and safety challenges and is recognized by the National Institute for Occupational Safety and Health (NIOSH) as a



priority industry sector in the National Occupational Research Agenda (NORA) to identify and provide intervention strategies regarding occupational health and safety issues.<sup>[11]</sup>

## Biological hazards

[edit]

Exposure to respiratory infectious diseases like tuberculosis (caused by *Mycobacterium tuberculosis*) and influenza can be reduced with the use of respirators; this exposure is a significant occupational hazard for health care professionals<sup>[12]</sup> Healthcare workers are also at risk for diseases that are contracted through extended contact with a patient, including scabies.<sup>[13]</sup> Health professionals are also at risk for contracting blood-borne diseases like hepatitis B, hepatitis C, and HIV/AIDS through needlestick injuries or contact with bodily fluids.<sup>[14]</sup><sup>[15]</sup> This risk can be mitigated with vaccination when there is a vaccine available, like with hepatitis B.<sup>[15]</sup> In epidemic situations, such as the 2014–2016 West African Ebola virus epidemic or the 2003 SARS outbreak, healthcare workers are at even greater risk, and were disproportionately affected in both the Ebola and SARS outbreaks.<sup>[16]</sup>

In general, appropriate personal protective equipment (PPE) is the first-line mode of protection for healthcare workers from infectious diseases. For it to be effective against highly contagious diseases, personal protective equipment must be watertight and prevent the skin and mucous membranes from contacting infectious material. Different levels of personal protective equipment created to unique standards are used in situations where the risk of infection is different. Practices such as triple gloving and multiple respirators do not provide a higher level of protection and present a burden to the worker, who is additionally at increased risk of exposure when removing the PPE. Compliance with appropriate personal protective equipment rules may be difficult in certain situations, such as tropical environments or low-resource settings. A 2020 Cochrane systematic review found low-quality evidence that using more breathable fabric in PPE, double gloving, and

active training reduce the risk of contamination but that more randomized controlled trials are needed for how best to train healthcare workers in proper PPE use<sup>[16]</sup>

## **Tuberculosis screening, testing, and education**

[edit]

Based on recommendations from The United States Center for Disease Control and Prevention (CDC) for TB screening and testing the following best practices should be followed when hiring and employing Health Care Personnel<sup>[17]</sup>

When hiring Health Care Personnel, the applicant should complete the following<sup>[18]</sup> a TB risk assessment,<sup>[19]</sup> a TB symptom evaluation for at least those listed on the Signs & Symptoms page,<sup>[20]</sup> a TB test in accordance with the guidelines for Testing for TB Infection,<sup>[21]</sup> and additional evaluation for TB disease as needed (e.g. chest x-ray for HCP with a positive TB test)<sup>[18]</sup> The CDC recommends either a blood test, also known as an interferon-gamma release assay (IGRA), or a skin test, also known as a Mantoux tuberculin skin test (TST).<sup>[21]</sup> A TB blood test for baseline testing does not require two-step testing. If the skin test method is used to test HCP upon hire, then two-step testing should be used. A one-step test is not recommended<sup>[18]</sup>

The CDC has outlined further specifics on recommended testing for several scenarios.<sup>[22]</sup> In summary:

1. Previous documented positive skin test (TST) then a further TST is not recommended
2. Previous documented negative TST within 12 months before employment OR at least two documented negative TSTs ever then a single TST is recommended
3. All other scenarios, with the exception of programs using blood tests, the recommended testing is a two-step TST

According to these recommended testing guidelines any two negative TST results within 12 months of each other constitute a two-step TST.

For annual screening, testing, and education, the only recurring requirement for all HCP is to receive TB education annually.<sup>[18]</sup> While the CDC offers education materials, there is not a well defined requirement as to what constitutes a satisfactory annual education. Annual TB testing is no longer recommended unless there is a known exposure or ongoing transmission at a healthcare facility. Should an HCP be considered at increased occupational risk for TB annual screening may be considered. For HCP with a documented history of a positive TB test result do not need to be re-tested but should instead complete a TB symptom evaluation. It is assumed that any HCP who has undergone a chest x-ray test has had a previous positive test result. When considering mental health you may see your doctor to be evaluated at your discretion. It is recommended to see someone at least once a year in order to make sure that there has not been any sudden changes.<sup>[23]</sup>

## **Psychosocial hazards**

[edit]

Occupational stress and occupational burnout are highly prevalent among health professionals.<sup>[24]</sup> Some studies suggest that workplace stress is pervasive in the health care industry because of inadequate staffing levels, long work hours, exposure to infectious diseases and hazardous substances leading to illness or death, and in some countries threat of malpractice litigation. Other stressors include the emotional labor of caring for ill people and high patient loads. The consequences of this stress can include substance abuse, suicide, major depressive disorder, and anxiety, all of which occur at higher rates in health professionals than the general working population. Elevated levels of stress are also linked to high rates of burnout, absenteeism and diagnostic errors, and reduced rates of patient satisfaction.<sup>[25]</sup> In Canada, a national report (*Canada's Health Care Providers*) also indicated higher rates of absenteeism due to illness or disability among health care workers compared to the rest of the working population, although those working in health care reported similar levels of good health and fewer reports of being injured at work.

[<sup>26</sup>]

There is some evidence that cognitive-behavioral therapy, relaxation training and therapy (including meditation and massage), and modifying schedules can reduce stress and burnout among multiple sectors of health care providers. Research is ongoing in this area, especially with regards to physicians, whose occupational stress and burnout is less researched compared to other health professions[<sup>27</sup>]

Healthcare workers are at higher risk of on-the-job injury due to violence. Drunk, confused, and hostile patients and visitors are a continual threat to providers attempting to treat patients. Frequently, assault and violence in a healthcare setting goes unreported and is wrongly assumed to be part of the job[<sup>28</sup>] Violent incidents typically occur during one-on-one care; being alone with patients increases healthcare workers' risk of assault.[<sup>29</sup>] In the United States, healthcare workers experience 2/3 of nonfatal workplace violence incidents.[<sup>28</sup>] Psychiatric units represent the highest proportion of violent incidents, at 40%; they are followed by geriatric units (20%) and the emergency department (10%). Workplace violence can also cause psychological trauma.[<sup>29</sup>]

Health care professionals are also likely to experience sleep deprivation due to their jobs. Many health care professionals are on a shift work schedule, and therefore experience misalignment of their work schedule and their circadian rhythm. In 2007, 32% of healthcare workers were found to get fewer than 6 hours of sleep a night. Sleep deprivation also predisposes healthcare professionals to make mistakes that may potentially endanger a patient.[<sup>30</sup>]

## **COVID pandemic**

[edit]

Especially in times like the present (2020), the hazards of health professional stem into the mental health. Research from the last few months highlights that COVID-19

has contributed greatly to the degradation of mental health in healthcare providers. This includes, but is not limited to, anxiety, depression/burnout, and insomnia.<sup>[citation needed]</sup>

A study done by Di Mattei et al. (2020) revealed that 12.63% of COVID nurses and 16.28% of other COVID healthcare workers reported extremely severe anxiety symptoms at the peak of the pandemic.<sup>[31]</sup> In addition, another study was conducted on 1,448 full time employees in Japan. The participants were surveyed at baseline in March 2020 and then again in May 2020. The result of the study showed that psychological distress and anxiety had increased more among healthcare workers during the COVID-19 outbreak.<sup>[32]</sup>

Similarly, studies have also shown that following the pandemic, at least one in five healthcare professionals report symptoms of anxiety.<sup>[33]</sup> Specifically, the aspect of "anxiety was assessed in 12 studies, with a pooled prevalence of 23.2%" following COVID.<sup>[33]</sup> When considering all 1,448 participants that percentage makes up about 335 people.

## **Abuse by patients**

[edit]

- The patients are selecting victims who are more vulnerable. For example, Cho said that these would be the nurses that are lacking experience or trying to get used to their new roles at work.<sup>[34]</sup>
- Others authors that agree with this are Vento, Cainelli, & Vallone and they said that, the reason patients have caused danger to health care workers is because of insufficient communication between them, long waiting lines, and overcrowding in waiting areas.<sup>[35]</sup> When patients are intrusive and/or violent toward the faculty, this makes the staff question what they should do about taking care of a patient.
- There have been many incidents from patients that have really caused some health care workers to be traumatized and have so much self doubt. Goldblatt and other authors said that there was a lady who was giving birth, her husband

said, "Who is in charge around here"? "Who are these sluts you employ here"<sup>[5]</sup>  
This was very avoidable to have been said to the people who are taking care of your wife and child.

## **Physical and chemical hazards**

[edit]

Slips, trips, and falls are the second-most common cause of worker's compensation claims in the US and cause 21% of work absences due to injury. These injuries most commonly result in strains and sprains; women, those older than 45, and those who have been working less than a year in a healthcare setting are at the highest risk<sup>[36]</sup>

An epidemiological study published in 2018 examined the hearing status of noise-exposed health care and social assistance (HSA) workers sector to estimate and compare the prevalence of hearing loss by subsector within the sector. Most of the HSA subsector prevalence estimates ranged from 14% to 18%, but the Medical and Diagnostic Laboratories subsector had 31% prevalence and the Offices of All Other Miscellaneous Health Practitioners had a 24% prevalence. The Child Day Care Services subsector also had a 52% higher risk than the reference industry<sup>[37]</sup>

Exposure to hazardous drugs, including those for chemotherapy, is another potential occupational risk. These drugs can cause cancer and other health conditions<sup>[38]</sup>

## **Gender factors**

[edit]

Female health care workers may face specific types of workplace-related health conditions and stress. According to the World Health Organization, women predominate in the formal health workforce in many countries and are prone to musculoskeletal injury (caused by physically demanding job tasks such as lifting and moving patients) and burnout. Female health workers are exposed to hazardous

drugs and chemicals in the workplace which may cause adverse reproductive outcomes such as spontaneous abortion and congenital malformations. In some contexts, female health workers are also subject to gender-based violence from coworkers and patients.<sup>[39][40]</sup>

## **Workforce shortages**

[edit]

See also: Health workforce, Doctor shortage, and Nursing shortage

Many jurisdictions report shortfalls in the number of trained health human resources to meet population health needs and/or service delivery targets, especially in medically underserved areas. For example, in the United States, the 2010 federal budget invested \$330 million to increase the number of physicians, physician assistants, nurse practitioners, nurses, and dentists practicing in areas of the country experiencing shortages of trained health professionals. The Budget expands loan repayment programs for physicians, nurses, and dentists who agree to practice in medically underserved areas. This funding will enhance the capacity of nursing schools to increase the number of nurses. It will also allow states to increase access to oral health care through dental workforce development grants. The Budget's new resources will sustain the expansion of the health care workforce funded in the Recovery Act.<sup>[41]</sup> There were 15.7 million health care professionals in the US as of 2011.<sup>[36]</sup>

In Canada, the 2011 federal budget announced a Canada Student Loan forgiveness program to encourage and support new family physicians, physician assistants, nurse practitioners and nurses to practice in underserved rural or remote communities of the country, including communities that provide health services to First Nations and Inuit populations.<sup>[42]</sup>

In Uganda, the Ministry of Health reports that as many as 50% of staffing positions for health workers in rural and underserved areas remain vacant. As of early 2011, the Ministry was conducting research and costing analyses to determine the most appropriate attraction and retention packages for medical officers, nursing officers, pharmacists, and laboratory technicians in the country's rural areas<sup>[43]</sup>

At the international level, the World Health Organization estimates a shortage of almost 4.3 million doctors, midwives, nurses, and support workers worldwide to meet target coverage levels of essential primary health care interventions<sup>[44]</sup> The shortage is reported most severe in 57 of the poorest countries, especially in sub-Saharan Africa.

Nurses are the most common type of medical field worker to face shortages around the world. There are numerous reasons that the nursing shortage occurs globally. Some include: inadequate pay, a large percentage of working nurses are over the age of 45 and are nearing retirement age, burnout, and lack of recognition<sup>[45]</sup>

Incentive programs have been put in place to aid in the deficit of pharmacists and pharmacy students. The reason for the shortage of pharmacy students is unknown but one can infer that it is due to the level of difficulty in the program<sup>[46]</sup>

Results of nursing staff shortages can cause unsafe staffing levels that lead to poor patient care. Five or more incidents that occur per day in a hospital setting as a result of nurses who do not receive adequate rest or meal breaks is a common issue.<sup>[47]</sup>

## **Regulation and registration**

[edit]

Main article: Health professional requisites

Practicing without a license that is valid and current is typically illegal. In most jurisdictions, the provision of health care services is regulated by the government.



Individuals found to be providing medical, nursing or other professional services without the appropriate certification or license may face sanctions and criminal charges leading to a prison term. The number of professions subject to regulation, requisites for individuals to receive professional licensure, and nature of sanctions that can be imposed for failure to comply vary across jurisdictions.

In the United States, under Michigan state laws, an individual is guilty of a felony if identified as practicing in the health profession without a valid personal license or registration. Health professionals can also be imprisoned if found guilty of practicing beyond the limits allowed by their licenses and registration. The state laws define the scope of practice for medicine, nursing, and a number of allied health professions<sup>[48]</sup> *[unreliable source?]* In Florida, practicing medicine without the appropriate license is a crime classified as a third degree felony<sup>[49]</sup> which may give imprisonment up to five years. Practicing a health care profession without a license which results in serious bodily injury classifies as a second degree felony<sup>[49]</sup> providing up to 15 years' imprisonment.

In the United Kingdom, healthcare professionals are regulated by the state; the UK Health and Care Professions Council (HCPC) protects the 'title' of each profession it regulates. For example, it is illegal for someone to call himself an Occupational Therapist or Radiographer if they are not on the register held by the HCPC.

## **See also**

[edit]

- List of healthcare occupations
- Community health center
- Chronic care management
- Electronic superbill
- Geriatric care management
- Health human resources
- Uniform Emergency Volunteer Health Practitioners Act

## References

[edit]

1. ^ "HCWs With Long COVID Report Doubt, Disbelief From Colleagues". *Medscape* . 29 November 2021.
2. ^ World Health Organization, 2006. *World Health Report 2006: working together for health*. Geneva: WHO.
3. ^ "Classifying health workers" (PDF). *World Health Organization*. Geneva. 2010. Archived (PDF) from the original on 2015-08-16. Retrieved 2016-02-13.
4. ^ "Difference Between Psychologists and Psychiatrists". *Psychology.about.com*. 2007. Archived from the original on April 3, 2007. Retrieved March 4, 2007.
5. ^ **a b** Goldblatt, Hadass; Freund, Anat; Drach-Zahavy, Anat; Enosh, Guy; Peterfreund, Ilana; Edlis, Neomi (2020-05-01). "Providing Health Care in the Shadow of Violence: Does Emotion Regulation Vary Among Hospital Workers From Different Professions?". *Journal of Interpersonal Violence*. **35** (9–10): 1908–1933. doi:10.1177/0886260517700620. ISSN 0886-2605. PMID 29294693. S2CID 19304885.
6. ^ Johnson, Cheryl L.; DeMass Martin, Suzanne L.; Markle-Elder, Sara (April 2007). "Stopping Verbal Abuse in the Workplace". *American Journal of Nursing*. **107** (4): 32–34. doi:10.1097/01.naj.0000271177.59574.c5. ISSN 0002-936X. PMID 17413727.
7. ^ Gupta N et al. "Human resources for maternal, newborn and child health: from measurement and planning to performance for improved health outcomes. Archived 2015-09-24 at the Wayback Machine *Human Resources for Health*, 2011, 9(16). Retrieved 20 October 2011.
8. ^ "Ob-Gyn Burnout: Why So Many Doctors Are Questioning Their Calling". *healthcareers.com*. Retrieved 2023-05-22.
9. ^ Araujo de Carvalho, Islene; Epping-Jordan, JoAnne; Pot, Anne Margriet; Kelley, Edward; Toro, Nuria; Thiyagarajan, Jotheeswaran A; Beard, John R (2017-11-01). "Organizing integrated health-care services to meet older people's needs". *Bulletin of the World Health Organization*. **95** (11): 756–763. doi:10.2471/BLT.16.187617 (inactive 5 December 2024). ISSN 0042-9686. PMC 5677611. PMID 29147056.cite journal: CS1 maint: DOI inactive as of December 2024 (link)
10. ^ Gupta N et al. "Health-related rehabilitation services: assessing the global supply of and need for human resources." Archived 2012-07-20 at the Wayback Machine *BMC Health Services Research*, 2011, 11:276. Published 17

October 2011. Retrieved 20 October 2011.

11. ^ "National Occupational Research Agenda for Healthcare and Social Assistance | NIOSH | CDC". [www.cdc.gov](http://www.cdc.gov). 2019-02-15. Retrieved 2019-03-14.
12. ^ Bergman, Michael; Zhuang, Ziqing; Shaffer, Ronald E. (25 July 2013). "Advanced Headforms for Evaluating Respirator Fit". National Institute for Occupational Safety and Health. Archived from the original on 16 January 2015. Retrieved 18 January 2015.
13. ^ FitzGerald, Deirdre; Grainger, Rachel J.; Reid, Alex (2014). "Interventions for preventing the spread of infestation in close contacts of people with scabies". *The Cochrane Database of Systematic Reviews*. **2014** (2): CD009943. doi:10.1002/14651858.CD009943.pub2. ISSN 1469-493X. PMC 10819104. PMID 24566946.
14. ^ Cunningham, Thomas; Burnett, Garrett (17 May 2013). "Does your workplace culture help protect you from hepatitis?". National Institute for Occupational Safety and Health. Archived from the original on 18 January 2015. Retrieved 18 January 2015.
15. ^ **a b** Reddy, Viraj K; Lavoie, Marie-Claude; Verbeek, Jos H; Pahwa, Manisha (14 November 2017). "Devices for preventing percutaneous exposure injuries caused by needles in healthcare personnel". *Cochrane Database of Systematic Reviews*. **2017** (11): CD009740. doi:10.1002/14651858.CD009740.pub3. PMC 6491125. PMID 29190036.
16. ^ **a b** Verbeek, Jos H.; Rajamaki, Blair; Ijaz, Sharea; Sauni, Riitta; Toomey, Elaine; Blackwood, Bronagh; Tikka, Christina; Ruotsalainen, Jani H.; Kilinc Balci, F. Selcen (May 15, 2020). "Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff". *The Cochrane Database of Systematic Reviews*. **2020** (5): CD011621. doi:10.1002/14651858.CD011621.pub5. hdl:1983/b7069408-3bf6-457a-9c6f-ecc38c00ee48. ISSN 1469-493X. PMC 8785899. PMID 32412096. S2CID 218649177.
17. ^ Sosa, Lynn E. (April 2, 2019). "Tuberculosis Screening, Testing, and Treatment of U.S. Health Care Personnel: Recommendations from the National Tuberculosis Controllers Association and CDC, 2019". *MMWR. Morbidity and Mortality Weekly Report*. **68** (19): 439–443. doi:10.15585/mmwr.mm6819a3. PMC 6522077. PMID 31099768.
18. ^ **a b c d** "Testing Health Care Workers | Testing & Diagnosis | TB | CDC". [www.cdc.gov](http://www.cdc.gov). March 8, 2021.
19. ^ "Health Care Personnel (HCP) Baseline Individual TB Risk Assessment" (PDF). [cdc.gov](http://cdc.gov). Retrieved 18 September 2022.

20. ^ "Signs & Symptoms | Basic TB Facts | TB | CDC". [www.cdc.gov](http://www.cdc.gov). February 4, 2021.
21. ^ **a b** "Testing for TB Infection | Testing & Diagnosis | TB | CDC". [www.cdc.gov](http://www.cdc.gov). March 8, 2021.
22. ^ "Guidelines for Preventing the Transmission of Mycobacterium tuberculosis in Health-Care Settings, 2005". [www.cdc.gov](http://www.cdc.gov).
23. ^ Spoorthy, Mamidipalli Sai; Pratapa, Sree Karthik; Mahant, Supriya (June 2020). "Mental health problems faced by healthcare workers due to the COVID-19 pandemic—A review". *Asian Journal of Psychiatry*. **51**: 102119. doi:10.1016/j.ajp.2020.102119. PMC 7175897. PMID 32339895.
24. ^ Ruotsalainen, Jani H.; Verbeek, Jos H.; Mariné, Albert; Serra, Consol (2015-04-07). "Preventing occupational stress in healthcare workers". *The Cochrane Database of Systematic Reviews*. **2015** (4): CD002892. doi:10.1002/14651858.CD002892.pub5. ISSN 1469-493X. PMC 6718215. PMID 25847433.
25. ^ "Exposure to Stress: Occupational Hazards in Hospitals". NIOSH Publication No. 2008–136 (July 2008). 2 December 2008. doi:10.26616/NIOSH PUB2008136. Archived from the original on 12 December 2008.
26. ^ *Canada's Health Care Providers, 2007 (Report)*. Ottawa: Canadian Institute for Health Information. 2007. Archived from the original on 2011-09-27.
27. ^ Ruotsalainen, JH; Verbeek, JH; Mariné, A; Serra, C (7 April 2015). "Preventing occupational stress in healthcare workers". *The Cochrane Database of Systematic Reviews*. **2015** (4): CD002892. doi:10.1002/14651858.CD002892.pub5. PMC 6718215. PMID 25847433.
28. ^ **a b** Hartley, Dan; Ridenour, Marilyn (12 August 2013). "Free On-line Violence Prevention Training for Nurses". National Institute for Occupational Safety and Health. Archived from the original on 16 January 2015. Retrieved 15 January 2015.
29. ^ **a b** Hartley, Dan; Ridenour, Marilyn (September 13, 2011). "Workplace Violence in the Healthcare Setting". NIOSH: Workplace Safety and Health. Medscape and NIOSH. Archived from the original on February 8, 2014.
30. ^ Caruso, Claire C. (August 2, 2012). "Running on Empty: Fatigue and Healthcare Professionals". NIOSH: Workplace Safety and Health. Medscape and NIOSH. Archived from the original on May 11, 2013.
31. ^ Di Mattei, Valentina; Perego, Gaia; Milano, Francesca; Mazzetti, Martina; Taranto, Paola; Di Pierro, Rossella; De Panfilis, Chiara; Madeddu, Fabio; Preti, Emanuele (2021-05-15). "The "Healthcare Workers' Wellbeing (Benessere Operatori)" Project: A Picture of the Mental Health Conditions of Italian Healthcare Workers during the First Wave of the COVID-19 Pandemic". *International Journal of Environmental Research and Public Health*. **18** (10): 5267. doi:

- 10.3390/ijerph18105267. ISSN 1660-4601. PMC 8156728. PMID 34063421.
32. ^ Sasaki, Natsu; Kuroda, Reiko; Tsuno, Kanami; Kawakami, Norito (2020-11-01). "The deterioration of mental health among healthcare workers during the COVID-19 outbreak: A population-based cohort study of workers in Japan". *Scandinavian Journal of Work, Environment & Health*. **46** (6): 639–644. doi:10.5271/sjweh.3922. ISSN 0355-3140. PMC 7737801. PMID 32905601.
  33. ^ **a b** Pappa, Sofia; Ntella, Vasiliki; Giannakas, Timoleon; Giannakoulis, Vassilis G.; Papoutsis, Eleni; Katsaounou, Paraskevi (August 2020). "Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis". *Brain, Behavior, and Immunity*. **88**: 901–907. doi:10.1016/j.bbi.2020.05.026. PMC 7206431. PMID 32437915.
  34. ^ Cho, Hyeonmi; Pavek, Katie; Steege, Linsey (2020-07-22). "Workplace verbal abuse, nurse-reported quality of care and patient safety outcomes among early-career hospital nurses". *Journal of Nursing Management*. **28** (6): 1250–1258. doi:10.1111/jonm.13071. ISSN 0966-0429. PMID 32564407. S2CID 219972442.
  35. ^ Vento, Sandro; Cainelli, Francesca; Vallone, Alfredo (2020-09-18). "Violence Against Healthcare Workers: A Worldwide Phenomenon With Serious Consequences". *Frontiers in Public Health*. **8**: 570459. doi:10.3389/fpubh.2020.570459. ISSN 2296-2565. PMC 7531183. PMID 33072706.
  36. ^ **a b** Collins, James W.; Bell, Jennifer L. (June 11, 2012). "Slipping, Tripping, and Falling at Work". *NIOSH: Workplace Safety and Health*. Medscape and NIOSH. Archived from the original on December 3, 2012.
  37. ^ Masterson, Elizabeth A.; Themann, Christa L.; Calvert, Geoffrey M. (2018-04-15). "Prevalence of Hearing Loss Among Noise-Exposed Workers Within the Health Care and Social Assistance Sector, 2003 to 2012". *Journal of Occupational and Environmental Medicine*. **60** (4): 350–356. doi:10.1097/JOM.0000000000001214. ISSN 1076-2752. PMID 29111986. S2CID 4637417.
  38. ^ Connor, Thomas H. (March 7, 2011). "Hazardous Drugs in Healthcare". *NIOSH: Workplace Safety and Health*. Medscape and NIOSH. Archived from the original on March 7, 2012.
  39. ^ World Health Organization. *Women and health: today's evidence, tomorrow's agenda*. Archived 2012-12-25 at the Wayback Machine Geneva, 2009. Retrieved on March 9, 2011.
  40. ^ Swanson, Naomi; Tisdale-Pardi, Julie; MacDonald, Leslie; Tiesman, Hope M. (13 May 2013). "Women's Health at Work". *National Institute for Occupational Safety and Health*. Archived from the original on 18 January 2015. Retrieved 21 January 2015.

41. ^ "Archived copy" (PDF). *Office of Management and Budget*. Retrieved 2009-03-06 – via *National Archives*.
42. ^ Government of Canada. 2011. *Canada's Economic Action Plan: Forgiving Loans for New Doctors and Nurses in Under-Served Rural and Remote Areas* Ottawa, 22 March 2011. Retrieved 23 March 2011.
43. ^ Rockers P et al. *Determining Priority Retention Packages to Attract and Retain Health Workers in Rural and Remote Areas in Uganda* Archived 2011-05-23 at the Wayback Machine CapacityPlus Project. February 2011.
44. ^ "The World Health Report 2006 - Working together for health". Geneva: WHO: World Health Organization. 2006. Archived from the original on 2011-02-28.
45. ^ Mefoh, Philip Chukwuemeka; Ude, Eze Nsi; Chukwuorji, JohBosco Chika (2019-01-02). "Age and burnout syndrome in nursing professionals: moderating role of emotion-focused coping". *Psychology, Health & Medicine*. **24** (1): 101–107. doi:10.1080/13548506.2018.1502457. ISSN 1354-8506. PMID 30095287. S2CID 51954488.
46. ^ Traynor, Kate (2003-09-15). "Staffing shortages plague nation's pharmacy schools". *American Journal of Health-System Pharmacy*. **60** (18): 1822–1824. doi:10.1093/ajhp/60.18.1822. ISSN 1079-2082. PMID 14521029.
47. ^ Leslie, G. D. (October 2008). "Critical Staffing shortage". *Australian Nursing Journal*. **16** (4): 16–17. doi:10.1016/s1036-7314(05)80033-5. ISSN 1036-7314. PMID 14692155.
48. ^ wiki.bmezine.com --> Practicing Medicine. In turn citing Michigan laws
49. ^ **a b** CHAPTER 2004-256 Committee Substitute for Senate Bill No. 1118 Archived 2011-07-23 at the Wayback Machine State of Florida, Department of State.

## External links

[edit]

- World Health Organization: Health workers

- v
- t
- e

Health care

- Economics
- Equipment
- Guidelines
- Industry
- Philosophy
- Policy
- Providers
- Public health
- Ranking
- Reform
- System

- Medicine

- Nursing

- Pharmacy

**Professions**

- Healthcare science

- Dentistry

- Allied health professions

- Health information management

- Assisted living

- Clinic

- Hospital

- Nursing home

**Settings**

- Medical school (Academic health science centre, Teaching hospital)

- Pharmacy school

- Supervised injection site

- Acute
- Chronic
- End-of-life
- Hospice
- Care**
  - Overutilization
  - Palliative
  - Primary
  - Self
  - Total
  - Bedside manner
- Skills / training**
  - Cultural competence
  - Diagnosis
  - Education
  - Universal precautions
  - 3D bioprinting
  - Artificial intelligence
  - Connected health
- Technology**
  - Digital health
  - Electronic health records
  - mHealth
  - Nanomedicine
  - Telemedicine



- Medical image computing and imaging informatics
- Artificial intelligence in healthcare
- Neuroinformatics in healthcare
- Behavior informatics in healthcare
- Computational biology in healthcare
- Translational bioinformatics
- Translational medicine
- health information technology
- Telemedicine
- Public health informatics
- Health information management
- Consumer health informatics
- United States
  - reform debate in the United States
  - History of public health in the United States

**Health informatics**

**By country**

- United Kingdom
- Canada
- Australia
- New Zealand
- (Category Health care by country)


-  **Category** image not found or type unknown

**Authority control databases: National** ○ Czech Republic  
**Edit this at Wikidata**

**About malocclusion**


"Deep bite" and "Buck teeth" redirect here. For the village, see Deep Bight, Newfoundland and Labrador.

## Malocclusion



Malocclusion in 10-year-old girl

**Specialty** Dentistry Image not found or type unknown



[Edit this on Wikidata](#)

Look up ***bucktooth*** in Wiktionary, the free dictionary.

In orthodontics, a **malocclusion** is a misalignment or incorrect relation between the teeth of the upper and lower dental arches when they approach each other as the jaws close. The English-language term dates from 1864;<sup>[1]</sup> Edward Angle (1855–1930), the "father of modern orthodontics",<sup>[2][3]</sup> *[need quotation to verify]* popularised it. The word derives from *mal-* 'incorrect' and *occlusion* 'the manner in which opposing teeth meet'.

The malocclusion classification is based on the relationship of the mesiobuccal cusp of the maxillary first molar and the buccal groove of the mandibular first molar. If this molar relationship exists, then the teeth can align into normal occlusion. According to Angle, malocclusion is any deviation of the occlusion from the ideal.<sup>[4]</sup> However, assessment for malocclusion should also take into account aesthetics and the impact on functionality. If these aspects are acceptable to the patient despite meeting the formal definition of malocclusion, then treatment may not be necessary.

It is estimated that nearly 30% of the population have malocclusions that are categorised as severe and definitely benefit from orthodontic treatment<sup>[5]</sup>

## Causes

[edit]

The aetiology of malocclusion is somewhat contentious, however, simply put it is multifactorial, with influences being both genetic<sup>[6]</sup><sup>[unreliable source?]</sup> and environmental.<sup>[7]</sup> Malocclusion is already present in one of the Skhul and Qafzeh hominin fossils and other prehistoric human skulls.<sup>[8]</sup><sup>[9]</sup> There are three generally accepted causative factors of malocclusion:

- Skeletal factors – the size, shape and relative positions of the upper and lower jaws. Variations can be caused by environmental or behavioral factors such as muscles of mastication, nocturnal mouth breathing, and cleft lip and cleft palate.
- Muscle factors – the form and function of the muscles that surround the teeth. This could be impacted by habits such as finger sucking, nail biting, pacifier and tongue thrusting<sup>[10]</sup>
- Dental factors – size of the teeth in relation to the jaw, early loss of teeth could result in spacing or mesial migration causing crowding, abnormal eruption path or timings, extra teeth (supernumeraries), or too few teeth (hypodontia)

There is not one single cause of malocclusion, and when planning orthodontic treatment it is often helpful to consider the above factors and the impact they have played on malocclusion. These can also be influenced by oral habits and pressure resulting in malocclusion.<sup>[11]</sup><sup>[12]</sup>

## Behavioral and dental factors

[edit]

In the active skeletal growth,<sup>[13]</sup> mouthbreathing, finger sucking, thumb sucking, pacifier sucking, onychophagia (nail biting), dermatophagia, pen biting, pencil biting, abnormal posture, deglutition disorders and other habits greatly influence the development of the face and dental arches.<sup>[14][15][16][17][18]</sup> Pacifier sucking habits are also correlated with otitis media.<sup>[19][20]</sup> Dental caries, periapical inflammation and tooth loss in the deciduous teeth can alter the correct permanent teeth eruptions.

## **Primary vs. secondary dentition**

[edit]

Malocclusion can occur in primary and secondary dentition.

In primary dentition malocclusion is caused by:

- Underdevelopment of the dentoalveolar tissue.
- Over development of bones around the mouth.
- Cleft lip and palate.
- Overcrowding of teeth.
- Abnormal development and growth of teeth.

In secondary dentition malocclusion is caused by:

- Periodontal disease.
- Overeruption of teeth.<sup>[21]</sup>
- Premature and congenital loss of missing teeth.

## **Signs and symptoms**

[edit]

Malocclusion is a common finding,<sup>[22][23]</sup> although it is not usually serious enough to require treatment. Those who have more severe malocclusions, which present as a

part of craniofacial anomalies, may require orthodontic and sometimes surgical treatment (orthognathic surgery) to correct the problem.

The ultimate goal of orthodontic treatment is to achieve a stable, functional and aesthetic alignment of teeth which serves to better the patient's dental and total health.<sup>[24]</sup> The symptoms which arise as a result of malocclusion derive from a deficiency in one or more of these categories.<sup>[25]</sup>

The symptoms are as follows:

- Tooth decay (caries): misaligned teeth will make it more difficult to maintain oral hygiene. Children with poor oral hygiene and diet will be at an increased risk.
- Periodontal disease: irregular teeth would hinder the ability to clean teeth meaning poor plaque control. Additionally, if teeth are crowded, some may be more buccally or lingually placed, there will be reduced bone and periodontal support. Furthermore, in Class III malocclusions, mandibular anterior teeth are pushed labially which contributes to gingival recession and weakens periodontal support.
- Trauma to anterior teeth: Those with an increased overjet are at an increased risk of trauma. A systematic review found that an overjet of greater than 3mm will double the risk of trauma.
- Masticatory function: people with anterior open bites, large increased & reverse overjet and hypodontia will find it more difficult to chew food.
- Speech impairment: a lisp is when the incisors cannot make contact, orthodontics can treat this. However, other forms of misaligned teeth will have little impact on speech and orthodontic treatment has little effect on fixing any problems.
- Tooth impaction: these can cause resorption of adjacent teeth and other pathologies for example a dentigerous cyst formation.
- Psychosocial wellbeing: malocclusions of teeth with poor aesthetics can have a significant effect on self-esteem.

Malocclusions may be coupled with skeletal disharmony of the face, where the relations between the upper and lower jaws are not appropriate. Such skeletal disharmonies often distort sufferer's face shape, severely affect aesthetics of the face, and may be coupled with mastication or speech problems. Most skeletal malocclusions can only be treated by orthognathic surgery.<sup>[citation needed]</sup>

## Classification

[edit]

Depending on the sagittal relations of teeth and jaws, malocclusions can be divided mainly into three types according to Angle's classification system published 1899. However, there are also other conditions, e.g. *crowding of teeth*, not directly fitting into this classification.

Many authors have tried to modify or replace Angle's classification. This has resulted in many subtypes and new systems (see section below: *Review of Angle's system of classes*).

A deep bite (also known as a Type II Malocclusion) is a condition in which the upper teeth overlap the lower teeth, which can result in hard and soft tissue trauma, in addition to an effect on appearance.<sup>[26]</sup> It has been found to occur in 15–20% of the US population.<sup>[27]</sup>

An open bite is a condition characterised by a complete lack of overlap and occlusion between the upper and lower incisors.<sup>[28]</sup> In children, open bite can be caused by prolonged thumb sucking.<sup>[29]</sup> Patients often present with impaired speech and mastication.<sup>[30]</sup>

## Overbites

[edit]

This is a vertical measurement of the degree of overlap between the maxillary incisors and the mandibular incisors. There are three features that are analysed in the classification of an overbite:


- Degree of overlap: edge to edge, reduced, average, increased
- Complete or incomplete: whether there is contact between the lower teeth and the opposing teeth/tissue (hard palate or gingivae) or not.
- Whether contact is traumatic or atraumatic

An average overbite is when the upper anterior teeth cover a third of the lower teeth. Covering less than this is described as 'reduced' and more than this is an 'increased' overbite. No overlap or contact is considered an 'anterior open bite'<sup>[25][31][32]</sup>


### Angle's classification method

[edit]

This section **may be too technical for most readers to understand**. Please **help improve** it to make it understandable to non-experts, without removing the technical details. *(September 2023) (Learn how and when to remove this message)*



Class I with severe crowding and labially erupted canines



Class II molar relationship

Edward Angle, who is considered the father of modern orthodontics, was the first to classify malocclusion. He based his classifications on the relative position of the maxillary first molar.<sup>[33]</sup> According to Angle, the mesiobuccal cusp of the upper first molar should align with the buccal groove of the mandibular first molar. The teeth should all fit on a line of occlusion which, in the upper arch, is a smooth curve through the central fossae of the posterior teeth and cingulum of the canines and incisors, and in the lower arch, is a smooth curve through the buccal cusps of the posterior teeth and incisal edges of the anterior teeth. Any variations from this resulted in malocclusion types. It is also possible to have different classes of malocclusion on left and right sides.

- **Class I** (Neutroclusion): Here the molar relationship of the occlusion is normal but the incorrect line of occlusion or as described for the maxillary first molar, but the other teeth have problems like spacing, crowding, over or under eruption, etc.
- **Class II** (Distocclusion (retrognathism, overjet, overbite)): In this situation, the mesiobuccal cusp of the upper first molar is not aligned with the mesiobuccal groove of the lower first molar. Instead it is anterior to it. Usually the mesiobuccal cusp rests in between the first mandibular molars and second premolars. There are two subtypes:
  - Class II Division 1: The molar relationships are like that of Class II and the anterior teeth are protruded.
  - Class II Division 2: The molar relationships are Class II but the central are retroclined and the lateral teeth are seen overlapping the centrals.



- **Class III:** (Mesioocclusion (prognathism, anterior crossbite, negative overjet, underbite)) In this case the upper molars are placed not in the mesiobuccal groove but posteriorly to it. The mesiobuccal cusp of the maxillary first molar lies posteriorly to the mesiobuccal groove of the mandibular first molar. Usually seen as when the lower front teeth are more prominent than the upper front teeth. In this case the patient very often has a large mandible or a short maxillary bone.

## **Review of Angle's system of classes and alternative systems**

[edit]

A major disadvantage of Angle's system of classifying malocclusions is that it only considers two dimensions along a spatial axis in the sagittal plane in the terminal occlusion, but occlusion problems can be three-dimensional. It does not recognise deviations in other spatial axes, asymmetric deviations, functional faults and other therapy-related features.

Angle's classification system also lacks a theoretical basis; it is purely descriptive. Its much-discussed weaknesses include that it only considers static occlusion, it does not account for the development and causes (aetiology) of occlusion problems, and it disregards the proportions (or relationships in general) of teeth and face<sup>[34]</sup> Thus, many attempts have been made to modify the Angle system or to replace it completely with a more efficient one,<sup>[35]</sup> but Angle's classification continues to be popular mainly because of its simplicity and clarity.<sup>[citation needed]</sup>

Well-known modifications to Angle's classification date back to Martin Dewey (1915) and Benno Lischer (1912, 1933). Alternative systems have been suggested by, among others, Simon (1930, the first three-dimensional classification system), Jacob A. Salzmann (1950, with a classification system based on skeletal structures) and James L. Ackerman and William R. Proffit (1969).<sup>[36]</sup>

## **Incisor classification**

[edit]

Besides the molar relationship, the British Standards Institute Classification also classifies malocclusion into incisor relationship and canine relationship.

- Class I: The lower incisor edges occlude with or lie immediately below the cingulum plateau of the upper central incisors
- Class II: The lower incisor edges lie posterior to the cingulum plateau of the upper incisors
  - Division 1 – the upper central incisors are proclined or of average inclination and there is an increase in overjet
  - Division 2 – The upper central incisors are retroclined. The overjet is usually minimal or may be increased.
- Class III: The lower incisor edges lie anterior to the cingulum plateau of the upper incisors. The overjet is reduced or reversed.

## **Canine relationship by Ricketts**

[edit]

- Class I: Mesial slope of upper canine coincides with distal slope of lower canine
- Class II: Mesial slope of upper canine is ahead of distal slope of lower canine
- Class III: Mesial slope of upper canine is behind to distal slope of lower canine

## **Crowding of teeth**

[edit]

Dental crowding is defined by the amount of space that would be required for the teeth to be in correct alignment. It is obtained in two ways: 1) by measuring the amount of space required and reducing this from calculating the space available via the width of the teeth, or 2) by measuring the degree of overlap of the teeth.

The following criterion is used:[<sup>25</sup>]

- 0–4mm = Mild crowding
- 4–8mm = Moderate crowding
- >8mm = Severe crowding

## Causes

[edit]

Genetic (inheritance) factors, extra teeth, lost teeth, impacted teeth, or abnormally shaped teeth have been cited as causes of crowding. Ill-fitting dental fillings, crowns, appliances, retainers, or braces as well as misalignment of jaw fractures after a severe injury are also known to cause crowding.[<sup>26</sup>] Tumors of the mouth and jaw, thumb sucking, tongue thrusting, pacifier use beyond age three, and prolonged use of a bottle have also been identified.[<sup>26</sup>]

Lack of masticatory stress during development can cause tooth overcrowding[<sup>37</sup>][<sup>38</sup>] Children who chewed a hard resinous gum for two hours a day showed increased facial growth.[<sup>37</sup>] Experiments in animals have shown similar results. In an experiment on two groups of rock hyraxes fed hardened or softened versions of the same foods, the animals fed softer food had significantly narrower and shorter faces and thinner and shorter mandibles than animals fed hard food.[<sup>37</sup>][<sup>39</sup>]<sup>[failed verification]</sup>

A 2016 review found that breastfeeding lowers the incidence of malocclusions developing later on in developing infants.[<sup>40</sup>]

During the transition to agriculture, the shape of the human mandible went through a series of changes. The mandible underwent a complex shape changes not matched by the teeth, leading to incongruity between the dental and mandibular form. These changes in human skulls may have been "driven by the decreasing bite forces required to chew the processed foods eaten once humans switched to growing different types of cereals, milking and herding animals about 10,000 years ago.[<sup>38</sup>][<sup>41</sup>]

## Treatment

[edit]

Orthodontic management of the condition includes dental braces, lingual braces, clear aligners or palatal expanders.<sup>[42]</sup> Other treatments include the removal of one or more teeth and the repair of injured teeth. In some cases, surgery may be necessary.<sup>[43]</sup>

## Treatment

[edit]

Malocclusion is often treated with orthodontics,<sup>[42]</sup> such as tooth extraction, clear aligners, or dental braces,<sup>[44]</sup> followed by growth modification in children or jaw surgery (orthognathic surgery) in adults. Surgical intervention is used only in rare occasions. This may include surgical reshaping to lengthen or shorten the jaw. Wires, plates, or screws may be used to secure the jaw bone, in a manner like the surgical stabilization of jaw fractures. Very few people have "perfect" alignment of their teeth with most problems being minor that do not require treatment.<sup>[37]</sup>

## Crowding

[edit]

Crowding of the teeth is treated with orthodontics, often with tooth extraction, clear aligners, or dental braces, followed by growth modification in children or jaw surgery (orthognathic surgery) in adults. Surgery may be required on rare occasions. This may include surgical reshaping to lengthen or shorten the jaw (orthognathic surgery). Wires, plates, or screws may be used to secure the jaw bone, in a manner similar to the surgical stabilization of jaw fractures. Very few people have "perfect" alignment of their teeth. However, most problems are very minor and do not require treatment.<sup>[39]</sup>

]

## **Class I**

[edit]

While treatment is not crucial in class I malocclusions, in severe cases of crowding can be an indication for intervention. Studies indicate that tooth extraction can have benefits to correcting malocclusion in individuals.<sup>[45][46]</sup> Further research is needed as reoccurring crowding has been examined in other clinical trials.<sup>[45][47]</sup>

## **Class II**

[edit]

A few treatment options for class II malocclusions include:

1. Functional appliance which maintains the mandible in a postured position to influence both the orofacial musculature and dentoalveolar development prior to fixed appliance therapy. This is ideally done through pubertal growth in pre-adolescent children and the fixed appliance during permanent dentition <sup>[48]</sup> Different types of removable appliances include Activator, Bionatar, Medium opening activator, Herbst, Frankel and twin block appliance with the twin block being the most widely used one.<sup>[49]</sup>
2. Growth modification through headgear to redirect maxillary growth
3. Orthodontic camouflage so that jaw discrepancy no longer apparent
4. Orthognathic surgery – sagittal split osteotomy mandibular advancement carried out when growth is complete where skeletal discrepancy is severe in anterior–posterior relationship or in vertical direction. Fixed appliance is required before, during and after surgery.
5. Upper Removable Appliance – limited role in contemporary treatment of increased overjets. Mostly used for very mild Class II, overjet due to incisor proclination, favourable overbite.

## **Class II Division 1**

[edit]

Low- to moderate- quality evidence suggests that providing early orthodontic treatment for children with prominent upper front teeth (class II division 1) is more effective for reducing the incidence of incisal trauma than providing one course of orthodontic treatment in adolescence.<sup>[50]</sup> There do not appear to be any other advantages of providing early treatment when compared to late treatment.<sup>[50]</sup> Low-quality evidence suggests that, compared to no treatment, late treatment in adolescence with functional appliances is effective for reducing the prominence of upper front teeth.<sup>[50]</sup>

## **Class II Division 2**

[edit]

Treatment can be undertaken using orthodontic treatments using dental braces.<sup>[51]</sup> While treatment is carried out, there is no evidence from clinical trials to recommend or discourage any type of orthodontic treatment in children.<sup>[51]</sup> A 2018 Cochrane systematic review anticipated that the evidence base supporting treatment approaches is not likely to improve occlusion due to the low prevalence of the condition and the ethical difficulties in recruiting people to participate in a randomized controlled trials for treating this condition.<sup>[51]</sup>

## **Class III**

[edit]

The British Standard Institute (BSI) classify class III incisor relationship as the lower incisor edge lies anterior to the cingulum plateau of the upper incisors, with reduced

or reversed over jet.[<sup>52</sup>] The skeletal facial deformity is characterized by mandibular prognathism, maxillary retrognathism or a combination of the two. This affects 3–8% of UK population with a higher incidence seen in Asia.[<sup>53</sup>]

One of the main reasons for correcting Class III malocclusion is aesthetics and function. This can have a psychological impact on the person with malocclusion resulting in speech and mastication problems as well. In mild class III cases, the patient is quite accepting of the aesthetics and the situation is monitored to observe the progression of skeletal growth.[<sup>54</sup>]

Maxillary and mandibular skeletal changes during prepubertal, pubertal and post pubertal stages show that class III malocclusion is established before the prepubertal stage.[<sup>55</sup>] One treatment option is the use of growth modification appliances such as the Chin Cap which has greatly improved the skeletal framework in the initial stages. However, majority of cases are shown to relapse into inherited class III malocclusion during the pubertal growth stage and when the appliance is removed after treatment.[<sup>55</sup>]

Another approach is to carry out orthognathic surgery, such as a bilateral sagittal split osteotomy (BSSO) which is indicated by horizontal mandibular excess. This involves surgically cutting through the mandible and moving the fragment forward or backwards for desired function and is supplemented with pre and post surgical orthodontics to ensure correct tooth relationship. Although the most common surgery of the mandible, it comes with several complications including: bleeding from inferior alveolar artery, unfavorable splits, condylar resorption, avascular necrosis and worsening of temporomandibular joint.[<sup>56</sup>]

Orthodontic camouflage can also be used in patients with mild skeletal discrepancies. This is a less invasive approach that uses orthodontic brackets to correct malocclusion and try to hide the skeletal discrepancy. Due to limitations of orthodontics, this option is more viable for patients who are not as concerned about the aesthetics of their facial appearance and are happy to address the malocclusion

only, as well as avoiding the risks which come with orthognathic surgery. Cephalometric data can aid in the differentiation between the cases that benefit from ortho-surgical or orthodontic treatment only (camouflage); for instance, examining a large group of orthognathic patient with Class III malocclusions they had average ANB angle of  $-3.57^\circ$  (95% CI,  $-3.92^\circ$  to  $-3.21^\circ$ ).<sup>[57]</sup>

## Deep bite

[edit]

The most common corrective treatments available are fixed or removal appliances (such as dental braces), which may or may not require surgical intervention. At this time there is no robust evidence that treatment will be successful.<sup>[51]</sup>

## Open bite

[edit]

An open bite malocclusion is when the upper teeth don't overlap the lower teeth. When this malocclusion occurs at the front teeth it is known as anterior open bite. An open bite is difficult to treat due to multifactorial causes, with relapse being a major concern. This is particularly so for an anterior open bite.<sup>[58]</sup> Therefore, it is important to carry out a thorough initial assessment in order to obtain a diagnosis to tailor a suitable treatment plan.<sup>[58]</sup> It is important to take into consideration any habitual risk factors, as this is crucial for a successful outcome without relapse. Treatment approach includes behavior changes, appliances and surgery. Treatment for adults include a combination of extractions, fixed appliances, intermaxillary elastics and orthognathic surgery.<sup>[30]</sup> For children, orthodontics is usually used to compensate for continued growth. With children with mixed dentition, the malocclusion may resolve on its own as the permanent teeth erupt. Furthermore, should the malocclusion be caused by childhood habits such as digit, thumb or pacifier sucking, it may result in resolution as the habit is stopped. Habit deterrent



appliances may be used to help in breaking digit and thumb sucking habits. Other treatment options for patients who are still growing include functional appliances and headgear appliances.

## **Tooth size discrepancy**

[edit]

Identifying the presence of tooth size discrepancies between the maxillary and mandibular arches is an important component of correct orthodontic diagnosis and treatment planning.

To establish appropriate alignment and occlusion, the size of upper and lower front teeth, or upper and lower teeth in general, needs to be proportional. Inter-arch tooth size discrepancy (ITSD) is defined as a disproportion in the mesio-distal dimensions of teeth of opposing dental arches. The prevalence is clinically significant among orthodontic patients and has been reported to range from 17% to 30%<sup>[59]</sup>


Identifying inter-arch tooth size discrepancy (ITSD) before treatment begins allows the practitioner to develop the treatment plan in a way that will take ITSD into account. ITSD corrective treatment may entail demanding reduction (interproximal wear), increase (crowns and resins), or elimination (extractions) of dental mass prior to treatment finalization.<sup>[60]</sup>

Several methods have been used to determine ITSD. Of these methods the one most commonly used is the Bolton analysis. Bolton developed a method to calculate the ratio between the mesiodistal width of maxillary and mandibular teeth and stated that a correct and harmonious occlusion is possible only with adequate proportionality of tooth sizes.<sup>[60]</sup> Bolton's formula concludes that if in the anterior portion the ratio is less than 77.2% the lower teeth are too narrow, the upper teeth are too wide or there is a combination of both. If the ratio is higher than 77.2% either the lower teeth are too wide, the upper teeth are too narrow or there is a combination of both.<sup>[59]</sup>

## Other conditions

[edit]

Further information: Open bite malocclusion



Open bite treatment after eight months of braces.

Other kinds of malocclusions can be due to or horizontal, vertical, or transverse skeletal discrepancies, including skeletal asymmetries.

Increased vertical growth causes a long facial profile and commonly leads to an open bite malocclusion, while decreased vertical facial growth causes a short facial profile and is commonly associated with a deep bite malocclusion. However, there are many other more common causes for open bites (such as tongue thrusting and thumb sucking) and likewise for deep bites.<sup>[61][62][63]</sup>

The upper or lower jaw can be overgrown (macrognathia) or undergrown (micrognathia).<sup>[62][61][63]</sup> It has been reported that patients with micrognathia are also affected by retrognathia (abnormal posterior positioning of the mandible or maxilla relative to the facial structure).<sup>[62]</sup> These patients are majorly predisposed to a class II malocclusion. Mandibular macrognathia results in prognathism and predisposes patients to a class III malocclusion.<sup>[64]</sup>

Most malocclusion studies to date have focused on Class III malocclusions. Genetic studies for Class II and Class I malocclusion are more rare. An example of hereditary mandibular prognathism can be seen amongst the Hapsburg Royal family where one third of the affected individuals with severe class III malocclusion had one parent with a similar phenotype <sup>[65]</sup>

The frequent presentation of dental malocclusions in patients with craniofacial birth defects also supports a strong genetic aetiology. About 150 genes are associated with craniofacial conditions presenting with malocclusions<sup>[66]</sup> Micrognathia is a commonly recurring craniofacial birth defect appearing among multiple syndromes.

For patients with severe malocclusions, corrective jaw surgery or orthognathic surgery may be carried out as a part of overall treatment, which can be seen in about 5% of the general population.<sup>[62]</sup><sup>[61]</sup><sup>[63]</sup>

## See also

[edit]

- Crossbite
- Elastics
- Facemask (orthodontics)
- Maximum intercuspation
- Mouth breathing
- Occlusion (dentistry)

## References

[edit]

1. <sup>^</sup> "malocclusion". *Oxford English Dictionary (Online ed.)*. Oxford University Press. (Subscription or participating institution membership required.)
2. <sup>^</sup> Bell B (September 1965). "Paul G. Spencer". *American Journal of Orthodontics*. **51** (9): 693–694. doi:10.1016/0002-9416(65)90262-9. PMID 14334001.
3. <sup>^</sup> Gruenbaum T (2010). "Famous Figures in Dentistry". *Mouth – JASDA*. **30** (1): 18.
4. <sup>^</sup> Hurt MA (2012). "Weedon D. Weedon's Skin Pathology. 3rd ed. London: Churchill Livingstone Elsevier, 2010". *Dermatology Practical & Conceptual*. **2** (1): 79–82. doi:10.5826/dpc.0201a15. PMC 3997252.
5. <sup>^</sup> Borzabadi-Farahani, A (2011). "An Overview of Selected Orthodontic Treatment Need Indices". In Naretto, Silvano (ed.). *Principles in Contemporary Orthodontics*. IntechOpen Limited. pp. 215–236. doi:10.5772/19735. ISBN 978-953-307-687-4.
6. <sup>^</sup> "How genetics can affect your teeth". *Orthodontics Australia*. 2018-11-25. Retrieved 2020-11-16.

7. ^ Corruccini RS, Potter RH (August 1980). "Genetic analysis of occlusal variation in twins". *American Journal of Orthodontics*. **78** (2): 140–54. doi:10.1016/0002-9416(80)90056-1. PMID 6931485.
8. ^ Sarig, Rachel; Slon, Viviane; Abbas, Janan; May, Hila; Shpack, Nir; Vardimon, Alexander Dan; HersHKovitz, Israel (2013-11-20). "Malocclusion in Early Anatomically Modern Human: A Reflection on the Etiology of Modern Dental Misalignment". *PLOS ONE*. **8** (11): e80771. Bibcode:2013PLoSO...880771S. doi:10.1371/journal.pone.0080771. ISSN 1932-6203. PMC 3835570. PMID 24278319.
9. ^ Pajević, Tina; Juloski, Jovana; Glišić, Branislav (2019-08-29). "Malocclusion from the prehistoric to the medieval times in Serbian population: Dentoalveolar and skeletal relationship comparisons in samples". *Homo: Internationale Zeitschrift für die vergleichende Forschung am Menschen*. **70** (1): 31–43. doi:10.1127/homo/2019/1009. ISSN 1618-1301. PMID 31475289. S2CID 201203069.
10. ^ Moimaz SA, Garbin AJ, Lima AM, Lolli LF, Saliba O, Garbin CA (August 2014). "Longitudinal study of habits leading to malocclusion development in childhood". *BMC Oral Health*. **14** (1): 96. doi:10.1186/1472-6831-14-96. PMC 4126276. PMID 25091288.
11. ^ Klein ET (1952). "Pressure Habits, Etiological Factors in Malocclusion". *Am. J. Orthod.* **38** (8): 569–587. doi:10.1016/0002-9416(52)90025-0.
12. ^ Graber TM. (1963). "The "Three m's": Muscles, Malformation and Malocclusion". *Am. J. Orthod.* **49** (6): 418–450. doi:10.1016/0002-9416(63)90167-2. hdl:2027.42/32220. S2CID 57626540.
13. ^ Björk A, Helm S (April 1967). "Prediction of the age of maximum puberal growth in body height" (PDF). *The Angle Orthodontist*. **37** (2): 134–43. PMID 4290545.
14. ^ Brucker M (1943). "Studies on the Incidence and Cause of Dental Defects in Children: IV. Malocclusion" (PDF). *J Dent Res*. **22** (4): 315–321. doi:10.1177/00220345430220041201. S2CID 71368994.
15. ^ Calisti LJ, Cohen MM, Fales MH (1960). "Correlation between malocclusion, oral habits, and socio-economic level of preschool children". *Journal of Dental Research*. **39** (3): 450–4. doi:10.1177/00220345600390030501. PMID 13806967. S2CID 39619434.
16. ^ Subtelny JD, Subtelny JD (October 1973). "Oral habits--studies in form, function, and therapy". *The Angle Orthodontist*. **43** (4): 349–83. PMID 4583311.
17. ^ Aznar T, Galán AF, Marín I, Domínguez A (May 2006). "Dental arch diameters and relationships to oral habits". *The Angle Orthodontist*. **76** (3): 441–5. PMID 16637724.
18. ^ Yamaguchi H, Sueishi K (May 2003). "Malocclusion associated with abnormal posture". *The Bulletin of Tokyo Dental College*. **44** (2): 43–54. doi:

10.2209/tdcpublication.44.43. PMID 12956088.

19. ^ Wellington M, Hall CB (February 2002). "Pacifier as a risk factor for acute otitis media". *Pediatrics*. **109** (2): 351–2, author reply 353. doi:10.1542/peds.109.2.351. PMID 11826228.
20. ^ Rovers MM, Numans ME, Langenbach E, Grobbee DE, Verheij TJ, Schilder AG (August 2008). "Is pacifier use a risk factor for acute otitis media? A dynamic cohort study". *Family Practice*. **25** (4): 233–6. doi:10.1093/fampra/cmn030. PMID 18562333.
21. ^ Hamish T (1990). *Occlusion*. Parkins, B. J. (2nd ed.). London: Wright. ISBN 978-0723620754. OCLC 21226656.
22. ^ Thilander B, Pena L, Infante C, Parada SS, de Mayorga C (April 2001). "Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development". *European Journal of Orthodontics*. **23** (2): 153–67. doi:10.1093/ejo/23.2.153. PMID 11398553.
23. ^ Borzabadi-Farahani A, Borzabadi-Farahani A, Eslamipour F (October 2009). "Malocclusion and occlusal traits in an urban Iranian population. An epidemiological study of 11- to 14-year-old children". *European Journal of Orthodontics*. **31** (5): 477–84. doi:10.1093/ejo/cjp031. PMID 19477970.
24. ^ "5 reasons you should see an orthodontist". *Orthodontics Australia*. 2017-09-27. Retrieved 2020-08-18.
25. ^ **a b c** Oliver RG (December 2001). "An Introduction to Orthodontics, 2nd edn". *Journal of Orthodontics*. **28** (4): 320. doi:10.1093/ortho/28.4.320.
26. ^ **a b c** Millett DT, Cunningham SJ, O'Brien KD, Benson PE, de Oliveira CM (February 2018). "Orthodontic treatment for deep bite and retroclined upper front teeth in children". *The Cochrane Database of Systematic Reviews*. **2** (3): CD005972. doi:10.1002/14651858.cd005972.pub4. PMC 6491166. PMID 29390172.
27. ^ Brunelle JA, Bhat M, Lipton JA (February 1996). "Prevalence and distribution of selected occlusal characteristics in the US population, 1988-1991". *Journal of Dental Research*. 75 Spec No (2 Suppl): 706–13. doi:10.1177/002203459607502S10. PMID 8594094. S2CID 30447284.
28. ^ de Castilho LS, Abreu MH, Pires e Souza LG, Romualdo LT, Souza e Silva ME, Resende VL (January 2018). "Factors associated with anterior open bite in children with developmental disabilities". *Special Care in Dentistry*. **38** (1): 46–50. doi:10.1111/scd.12262. PMID 29278267. S2CID 42747680.
29. ^ Feres MF, Abreu LG, Insabralde NM, Almeida MR, Flores-Mir C (June 2016). "Effectiveness of the open bite treatment in growing children and adolescents. A systematic review". *European Journal of Orthodontics*. **38** (3): 237–50.

doi:10.1093/ejo/cjv048. PMC 4914905. PMID 26136439.

30. ^ **a b** Cambiano AO, Janson G, Lorenzoni DC, Garib DG, Dávalos DT (2018). "Nonsurgical treatment and stability of an adult with a severe anterior open-bite malocclusion". *Journal of Orthodontic Science*. **7**: 2. doi:10.4103/jos.JOS\_69\_17. PMC 5952238. PMID 29765914.
31. ^ Houston, W. J. B. (1992-02-01). "Book Reviews". *The European Journal of Orthodontics*. **14** (1): 69. doi:10.1093/ejo/14.1.69.
32. ^ Hamdan AM, Lewis SM, Kelleher KE, Elhady SN, Lindauer SJ (November 2019). "Does overbite reduction affect smile esthetics?". *The Angle Orthodontist*. **89** (6): 847–854. doi:10.2319/030819-177.1. PMC 8109173. PMID 31306077.
33. ^ "Angle's Classification of Malocclusion". Archived from the original on 2008-02-13 . Retrieved 2007-10-31.
34. ^ Sunil Kumar (Ed.): Orthodontics. New Delhi 2008, 624 p., ISBN 978–81–312–1054–3, p. 127
35. ^ Sunil Kumar (Ed.): Orthodontics. New Delhi 2008, p. 123. A list of 18 approaches to modify or replace Angle's system is given here with further references at the end of the book.
36. ^ Gurkeerat Singh: Textbook of Orthodontics, p. 163–170, with further references on p. 174.
37. ^ **a b c d** Lieberman, D (May 2004). "Effects of food processing on masticatory strain and craniofacial growth in a retrognathic face". *Journal of Human Evolution*. **46** (6): 655–77. doi:10.1016/s0047-2484(04)00051-x. PMID 15183669.
38. ^ **a b** Ingervall B, Bitsanis E (February 1987). "A pilot study of the effect of masticatory muscle training on facial growth in long-face children" (PDF). *European Journal of Orthodontics*. **9** (1): 15–23. doi:10.1093/ejo/9.1.15. PMID 3470182.
39. ^ **a b** Rosenberg J (2010-02-22). "Malocclusion of teeth". *Medline Plus*. Retrieved 2012-02-06.
40. ^ Victora CG, Bahl R, Barros AJ, França GV, Horton S, Krasevec J, Murch S, Sankar MJ, Walker N, Rollins NC (January 2016). "Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect". *Lancet*. **387** (10017): 475–90. doi:10.1016/s0140-6736(15)01024-7. PMID 26869575.
41. ^ Quaglio CL, de Freitas KM, de Freitas MR, Janson G, Henriques JF (June 2011). "Stability and relapse of maxillary anterior crowding treatment in class I and class II Division 1 malocclusions". *American Journal of Orthodontics and Dentofacial Orthopedics*. **139** (6): 768–74. doi:10.1016/j.ajodo.2009.10.044. PMID 21640883.

42. ^ **a b** "Dental Crowding: Causes and Treatment Options". *Orthodontics Australia*. 2020-06-29. Retrieved 2020-11-19.
43. ^ "Malocclusion of teeth: MedlinePlus Medical Encyclopedia". *medlineplus.gov*. Retrieved 2021-04-07.
44. ^ "Can Buck Teeth Be Fixed? Causes & Treatment Options". *Orthodontics Australia*. 2021-07-01. Retrieved 2021-10-11.
45. ^ **a b** Alam, MK (October 2018). "Treatment of Angle Class I malocclusion with severe crowding by extraction of four premolars: a case report". *Bangladesh Journal of Medical Science*. **17** (4): 683–687. doi:10.3329/bjms.v17i4.38339.
46. ^ Persson M, Persson EC, Skagius S (August 1989). "Long-term spontaneous changes following removal of all first premolars in Class I cases with crowding". *European Journal of Orthodontics*. **11** (3): 271–82. doi:10.1093/oxfordjournals.ejo.a035995. PMID 2792216.
47. ^ von Cramon-Taubadel N (December 2011). "Global human mandibular variation reflects differences in agricultural and hunter-gatherer subsistence strategies". *Proceedings of the National Academy of Sciences of the United States of America* **108** (49): 19546–51. Bibcode:2011PNAS..10819546V. doi:10.1073/pnas.1113050108. PMC 3241821. PMID 22106280.
48. ^ Nayak KU, Goyal V, Malviya N (October 2011). "Two-phase treatment of class II malocclusion in young growing patient". *Contemporary Clinical Dentistry*. **2** (4): 376–80. doi:10.4103/0976-237X.91808. PMC 3276872. PMID 22346172.
49. ^ "Treatment of class ii malocclusions". 2013-11-14.
50. ^ **a b c** Pinhasi R, Eshed V, von Cramon-Taubadel N (2015-02-04). "Incongruity between affinity patterns based on mandibular and lower dental dimensions following the transition to agriculture in the Near East, Anatolia and Europe". *PLOS ONE*. **10** (2): e0117301. Bibcode:2015PLoSO..1017301P. doi:10.1371/journal.pone.0117301. PMC 4317182. PMID 25651540.
51. ^ **a b c d** Batista KB, Thiruvengkatachari B, Harrison JE, O'Brien KD (March 2018). "Orthodontic treatment for prominent upper front teeth (Class II malocclusion) in children and adolescents". *The Cochrane Database of Systematic Reviews*. **2018** (3): CD003452. doi:10.1002/14651858.cd003452.pub4. PMC 6494411. PMID 29534303.
52. ^ CLASSIFICATION OF SKELETAL AND DENTAL MALOCCLUSION: REVISITED; Mageet, Adil Osman (2016). "Classification of Skeletal and Dental Malocclusion: Revisited". *Stomatology Edu Journal*. **3** (2): 205–211. doi:10.25241/2016.3(2).11.
53. ^ *Esthetics and biomechanics in orthodontics*. Nanda, Ravindra,, Preceded by (work): Nanda, Ravindra. (Second ed.). St. Louis, Missouri. 2014-04-10. ISBN978-0-323-22659-2. OCLC 880707123.cite book: CS1 maint: location missing

publisher (link) CS1 maint: others (link)

54. ^ Eslami S, Faber J, Fateh A, Sheikholaemmeh F, Grassia V, Jamilian A (August 2018). "Treatment decision in adult patients with class III malocclusion: surgery versus orthodontics". *Progress in Orthodontics*. **19** (1): 28. doi:10.1186/s40510-018-0218-0. PMC 6070451. PMID 30069814.
55. ^ **a b** Uner O, Yüksel S, Uçüncü N (April 1995). "Long-term evaluation after chin cap treatment". *European Journal of Orthodontics*. **17** (2): 135–41. doi:10.1093/ejo/17.2.135. PMID 7781722.
56. ^ Ravi MS, Shetty NK, Prasad RB (January 2012). "Orthodontics-surgical combination therapy for Class III skeletal malocclusion". *Contemporary Clinical Dentistry*. **3** (1): 78–82. doi:10.4103/0976-237X.94552. PMC 3341765. PMID 22557903.
57. ^ Borzabadi Farahani A, Olkun HK, Eslamian L, Eslamipour F (2024). "A retrospective investigation of orthognathic patients and functional needs". *Australasian Orthodontic Journal*. **40**: 111–120. doi:10.2478/aoj-2024-0013.
58. ^ **a b** Atsawasuwana P, Hohlt W, Evans CA (April 2015). "Nonsurgical approach to Class I open-bite malocclusion with extrusion mechanics: a 3-year retention case report". *American Journal of Orthodontics and Dentofacial Orthopedics*. **147** (4): 499–508. doi:10.1016/j.ajodo.2014.04.024. PMID 25836010.
59. ^ **a b** Grauer D, Heymann GC, Swift EJ (June 2012). "Clinical management of tooth size discrepancies". *Journal of Esthetic and Restorative Dentistry*. **24** (3): 155–9. doi:10.1111/j.1708-8240.2012.00520.x. PMID 22691075. S2CID 11482185.
60. ^ **a b** Cançado RH, Gonçalves Júnior W, Valarelli FP, Freitas KM, Crêspo JA (2015). "Association between Bolton discrepancy and Angle malocclusions". *Brazilian Oral Research*. **29**: 1–6. doi:10.1590/1807-3107BOR-2015.vol29.0116. PMID 26486769.
61. ^ **a b c** Harrington C, Gallagher JR, Borzabadi-Farahani A (July 2015). "A retrospective analysis of dentofacial deformities and orthognathic surgeries using the index of orthognathic functional treatment need (IOFTN)". *International Journal of Pediatric Otorhinolaryngology*. **79** (7): 1063–6. doi:10.1016/j.ijporl.2015.04.027. PMID 25957779.
62. ^ **a b c d** Posnick JC (September 2013). "Definition and Prevalence of Dentofacial Deformities". *Orthognathic Surgery: Principles and Practice*. Amsterdam: Elsevier. pp. 61–68. doi:10.1016/B978-1-4557-2698-1.00003-4. ISBN 978-145572698-1.
63. ^ **a b c** Borzabadi-Farahani A, Eslamipour F, Shahmoradi M (June 2016). "Functional needs of subjects with dentofacial deformities: A study using the index of orthognathic functional treatment need (IOFTN)". *Journal of Plastic, Reconstructive & Aesthetic Surgery*. **69** (6): 796–801. doi:10.1016/j.bjps.2016.03.008. PMID 27068664.



64. <sup>^</sup> Purkait, S (2011). *Essentials of Oral Pathology 4th Edition*.
65. <sup>^</sup> Joshi N, Hamdan AM, Fakhouri WD (December 2014). "Skeletal malocclusion: a developmental disorder with a life-long morbidity". *Journal of Clinical Medicine Research*. **6** (6): 399–408. doi:10.14740/jocmr1905w. PMC 4169080. PMID 25247012.
66. <sup>^</sup> Moreno Uribe LM, Miller SF (April 2015). "Genetics of the dentofacial variation in human malocclusion". *Orthodontics & Craniofacial Research*. **18** Suppl 1 (S1): 91–9. doi:10.1111/ocr.12083. PMC 4418210. PMID 25865537.


## Further reading

[edit]


- Peter S. Ungar, "The Trouble with Teeth: Our teeth are crowded, crooked and riddled with cavities. It hasn't always been this way", *Scientific American*, vol. 322, no. 4 (April 2020), pp. 44–49. "Our teeth [...] evolved over hundreds of millions of years to be incredibly strong and to align precisely for efficient chewing. [...] Our dental disorders largely stem from a shift in the oral environment caused by the introduction of softer, more sugary foods than the ones our ancestors typically ate."

## External links

[edit]



- **ICD-10:** K07.3, K07.4, K07.5, D K07.6
- **ICD-9-CM:** 524.4
- **MeSH:** D008310



Wikimedia Commons has media related to ***Malocclusion***.

- v
- t

- e

## Orthodontics

- Bolton analysis
- Cephalometric analysis
- Cephalometry
- Dentition analysis
- Failure of eruption of teeth
- Diagnosis**
  - Little's Irregularity Index
  - Malocclusion
  - Scissor bite
  - Standard anatomical position
  - Tooth ankylosis
  - Tongue thrust
  - Overbite
  - Overjet
  - Open bite
  - Crossbite
  - Dental crowding
  - Dental spacing
- Conditions**
  - Bimaxillary Protrusion
  - Prognathism
  - Retrognathism
  - Maxillary hypoplasia
  - Condylar hyperplasia
  - Overeruption
  - Mouth breathing
  - Temporomandibular dysfunction

- ACCO appliance
- Archwire
- Activator appliance
- Braces
- Damon system
- Elastics
- Frankel appliance
- Invisalign
- Lingual arch
- Lip bumper
- Herbst Appliance
- List of orthodontic functional appliances
- List of palatal expanders
- Lingual braces
- Headgear
- Orthodontic technology
- Orthodontic spacer
- Palatal lift prosthesis
- Palatal expander
- Quad helix
- Retainer
- SureSmile
- Self-ligating braces
- Splint activator
- Twin Block Appliance

## **Appliances**

**Procedures**

- Anchorage (orthodontics)
- Cantilever mechanics
- Fiberotomy
- Interproximal reduction
- Intrusion (orthodontics)
- Molar distalization
- SARPE
- Serial extraction
- Beta-titanium
- Nickel titanium
- Stainless steel

**Materials**

- TiMolium
- Elgiloy
- Ceramic
- Composite
- Dental elastics

- Edward Angle
- Spencer Atkinson
- Clifford Ballard
- Raymond Begg
- Hans Peter Bimler
- Samir Bishara
- Arne Björk
- Charles B. Bolton
- Holly Broadbent Sr.
- Allan G. Brodie
- Charles J. Burstone
- Peter Buschang
- Calvin Case
- Harold Chapman (Orthodontist)
- David Di Biase
- Jean Delaire
- Terry Dischinger
- William B. Downs
- John Nutting Farrar
- Rolf Frankel
- Sheldon Friel
- Thomas M. Graber
- Charles A. Hawley
- Reed Holdaway
- John Hooper (Orthodontist)
- Joseph Jarabak
- Harold Kesling
- Albert Ketcham
- Juri Kurol
- Craven Kurz
- Benno Lischer
- James A. McNamara
- Birte Melsen
- Robert Moyers
- Hayes Nance

**Notable  
contributors**

- American Association of Orthodontists
- American Board of Orthodontics
- British Orthodontic Society
- Organizations**
  - Canadian Association of Orthodontists
  - Indian Orthodontic Society
  - Italian Academy of Orthodontic Technology
  - Society for Orthodontic Dental Technology (Germany)
  - American Journal of Orthodontics and Dentofacial Orthopedics
- Journals**
  - The Angle Orthodontist
  - Journal of Orthodontics
- Institution**
  - Angle School of Orthodontia

- v
- t
- e

Dental disease involving the jaw

- General**
  - Jaw abnormality
  - malocclusion
  - Orthodontics
  - Gnathitis
- Size**
  - Micrognathism
  - Maxillary hypoplasia
  - Cherubism
- Maxilla and Mandible**
  - Congenital epulis
  - Torus mandibularis
  - Torus palatinus

## Other

- Jaw and base of cranium
  - Prognathism
  - Retrognathism
- Dental arch
  - Crossbite
  - Overbite
- Temporomandibular joint disorder

Check our other pages :

- [Clear Aligners for Subtle Tooth Movement](#)
- [Collaborative Approaches to Pediatric Growth Management](#)
- [Shared Planning for Improved Patient Experiences](#)

IQDENT – Ortodontska Klinika

Phone : +385953817015

City : Zagreb

State : Hrvatska

Zip : 10000

Address : IQDENT – Ortodontska Klinika

Company Website : <https://iqdent.hr/>

USEFUL LINKS

[Orthodontic treatment can help improve your child's smile](#)

[Orthodontic treatment for children](#)

[Sitemap](#)

[Privacy Policy](#)

[About Us](#)

Follow us