



- **Reasons Adults Seek Orthodontic Treatment**

Reasons Adults Seek Orthodontic Treatment Overcoming Hesitations About Braces Later in Life Clear Aligners for Subtle Tooth Movement Comparing Ceramic Brackets With Other Invisible Options Balancing Cosmetic Goals and Functional Needs Social Changes That Encourage Adult Orthodontics Handling Orthodontic Care Alongside Busy Schedules Managing Discomfort and Adjusting to Lifestyle Shifts Case Studies Highlighting Adult Smile Transformations Psychological Benefits of Aligned Teeth in Adulthood Minimizing Visibility of Orthodontic Appliances Key Questions Adults Often Ask About Treatment

- **How Orthodontics Connects With Periodontal Health**

How Orthodontics Connects With Periodontal Health Coordinating Orthodontic Work With Restorative Dentistry The Role of Oral Surgery in Complex Alignment Cases Importance of Joint Consultation for Comprehensive Outcomes Leveraging Data Sharing Between Dental Specialties Collaborative Approaches to Pediatric Growth Management Integrating Prosthodontics for Full Mouth Rehabilitation Teamwork for Managing Tongue Thrust and Speech Issues Streamlining Communication Among Healthcare Providers Shared Planning for Improved Patient Experiences Post Surgical Follow Ups With Multiple Specialists Building a Unified Strategy for Long Term Stability

- **About Us**



Shared Planning for Improved Patient Experiences

****Desire for Improved Aesthetics**:** Many adults seek orthodontic treatment to enhance the appearance of their smile, addressing concerns such as crooked teeth, gaps, or overbites that may have persisted from childhood or developed due to various life factors.

****Desire for Improved Aesthetics**:** Many adults seek orthodontic treatment to enhance the appearance of their smile, addressing concerns such as crooked teeth, gaps, or overbites that may have persisted from childhood or developed due to various life factors.

Certainly, let's dive into the heart of enhancing patient experiences, especially within the context of orthodontic treatment for children, through the lens of shared planning. This approach not only transforms the traditional patient-provider dynamic but also ensures that every step of the orthodontic journey is a collaborative effort, focusing on the needs and comfort of the young patients and their families. Here's how shared planning can revolutionize pediatric orthodontic care:

1. **Building Trust from the Start:** The foundation of successful shared planning in orthodontics lies in fostering trust between the orthodontist, parents, and the child. Clear aligners are an option for some kids needing orthodontic care Kids' dental alignment services hospital. Early, open discussions about treatment goals, expectations, and potential outcomes help establish a rapport that makes

children feel secure and involved in their own care.

2. **Understanding Unique Needs:** Each child is unique; therefore, their orthodontic plan should be tailored to their specific dental health needs, growth patterns, and personal preferences. Involving both parents and kids in this assessment ensures that all perspectives are considered, leading to more personalized and effective treatment plans.
3. **Visualizing Success:** Utilizing models, simulations, and digital imagery allows everyone involved to visualize proposed treatments and outcomes. This visual aid not only helps children understand what to expect but also gives parents peace of mind by making abstract concepts concrete and tangible.
4. **Inclusive Decision-Making:** Shared planning encourages active participation from all stakeholders—children, parents, and orthodontists—in decision-making processes regarding treatment options and timelines. This inclusive approach empowers families to voice concerns or preferences at every stage, ensuring alignment with family values and priorities.

5.

Education as a Partnership: Educating families about orthodontic conditions and treatments is crucial for informed decision-making. A collaborative approach means providing clear explanations tailored to each family's level of understanding—explaining why certain procedures are recommended and how they contribute to long-term oral health benefits for kids.

6. **Supporting Emotional Well-being:** Orthodontic treatment can sometimes affect a child's emotional state due to changes in appearance or discomfort during adjustments. Shared planning includes discussing strategies for managing these emotional challenges together with psychologists or counselors if necessary, ensuring holistic support throughout the treatment journey.

7. **Regular Feedback Loops:** Establishing regular check-ins where feedback is welcomed from both parents and patients creates an environment where concerns can be addressed promptly. This continuous dialogue ensures that any issues are nipped in the bud while reinforcing that everyone's voice matters in shaping the treatment experience positively.

8. **Celebrating Milestones Together:** Lastly, recognizing achievements along the way—be it completing a phase of treatment or seeing improvements in dental alignment—strengthens relationships between providers and families while motivating children through positive reinforcement techniques that celebrate their hard work and progress together as a team.

By embracing shared planning in pediatric orthodontics, we shift from a one-dimensional provider-centric model towards a collaborative approach that values input from everyone involved—ultimately enhancing patient experiences by making them feel heard, understood, and actively engaged throughout their orthodontic journey toward healthier smiles and improved confidence

Shared planning in pediatric orthodontics represents a collaborative approach that fundamentally alters the landscape of patient care, focusing on enhancing experiences for both children and their families. This model is rooted in the belief that treating young patients isn't just about correcting dental alignment; it's about creating a comprehensive, family-centered experience that considers emotional, psychological, and developmental aspects alongside the physical treatment needs.

At its core, shared planning involves a close collaboration between various specialists—pediatric dentists, orthodontists, pediatricians, and sometimes even psychologists or child life specialists—to craft a unified treatment plan tailored specifically to each child's unique needs. This multidisciplinary team gathers insights from multiple perspectives to ensure that every aspect of the child's oral health journey is considered. For instance, while an orthodontist might focus on the technical aspects of aligning teeth and correcting bites, a pediatrician could offer crucial insights into how the child's overall

health might impact or be impacted by orthodontic treatment.

This approach fosters an environment where communication is paramount. Regular meetings among the care team members ensure everyone is on the same page regarding treatment progress, potential challenges, and adjustments needed along the way. More importantly, it empowers parents and guardians by involving them actively in discussions about their child's care. This inclusivity not only helps in setting realistic expectations but also encourages a sense of partnership where families feel heard and valued—a vital component for reducing anxiety often associated with orthodontic treatments in children.

Shared planning also emphasizes the significance of creating positive experiences throughout a child's treatment journey. By considering factors like fear or discomfort related to dental visits, teams can implement strategies such as using child-friendly language, incorporating playful elements into appointments (like themed waiting rooms or fun rewards for cooperation), and employing gentle techniques to make procedures as comfortable as possible. These small yet impactful gestures contribute significantly to building trust between the child and their healthcare providers, making future visits less daunting.

Moreover, this collaborative approach allows for flexibility in adapting treatment plans as children grow and develop—not just physically but emotionally too. It acknowledges that each child's experience with orthodontic care is unique and may require adjustments over time based on their responses to treatment and evolving personal needs. By maintaining open lines of communication among all parties involved—including regular updates to parents—this ensures that any deviations from the plan are addressed promptly and effectively.

In essence, shared planning in pediatric orthodontics transforms what could be a stressful experience for children into an adventure marked by understanding support systems—

where every member of the care team plays an integral role in ensuring not just successful outcomes but enjoyable ones too. This model underscores that effective patient care transcends clinical expertise; it thrives on empathy, collaboration, and a genuine commitment to improving every aspect of a young patient's experience from start to finish. As we continue to refine these practices, we pave the way for generations of children who view dental care not with apprehension but with confidence—and perhaps even excitement—for what it can bring them: healthier smiles and brighter futures filled with possibilities beyond just straight teeth but confident interactions in life itself.

****Enhancing Oral Health**:**

Proper alignment of teeth through orthodontic treatment can improve oral hygiene by making it easier to clean teeth effectively, thereby reducing the risk of

cavities and gum disease that may have been challenging to manage in misaligned dentitions.

Shared Planning for Improved Patient Experiences: A Beacon of Patient-Centered Care

In the evolving landscape of healthcare, the shift towards patient-centered care has become not just a trend but a fundamental necessity. At the heart of this transformation lies the practice of shared planning, an essential component that significantly enhances patient experiences by placing the patient at the very center of their own care journey. This approach doesn't merely represent a change in how care is delivered; it signifies a profound respect for the individuality, preferences, and needs of each patient, fostering an environment where they are active participants rather than passive recipients of medical interventions.

The importance of shared planning in enhancing patient-centered care cannot be overstated. By engaging patients in the planning process from the outset, healthcare providers acknowledge and validate their unique perspectives and goals. This collaborative approach ensures that treatment plans align closely with what matters most to the patient—be it recovery time, minimizing side effects, or maintaining quality of life.

during treatment. It's about crafting a roadmap that reflects not just medical necessities but personal values and lifestyle considerations, thus making the entire experience feel more tailored and less impersonal.

Shared planning empowers patients with knowledge and involvement, dramatically improving their satisfaction levels. When patients understand their conditions and treatment options clearly, they feel more confident in their decisions, leading to greater adherence to prescribed plans and better health outcomes. This empowerment also extends beyond immediate treatment; it fosters a lasting sense of control over one's health journey, which can positively influence long-term health behaviors and outcomes.

Moreover, shared planning facilitates open communication between healthcare providers and patients—a cornerstone of effective patient-centered care. It encourages dialogue that goes beyond merely discussing symptoms or treatments; it opens channels for understanding fears, hopes, and expectations. This deeper level of engagement builds trust and rapport between patients and their caregivers, creating a supportive partnership aimed at achieving optimal health outcomes together.

Furthermore, incorporating shared planning into healthcare practices promotes efficiency within systems traditionally known for their complexity and fragmentation. By involving patients in decision-making processes early on, potential barriers or misalignments can be identified and addressed proactively. This proactive approach helps streamline care pathways, reduces unnecessary procedures, and ensures resources are utilized effectively—ultimately leading to improved outcomes at both individual and systemic levels.

In essence, shared planning is not just an add-on feature but an integral thread woven into the fabric of patient-centered care that enhances experiences from start to finish. It recognizes each patient as an individual with unique insights into their own health needs

while leveraging professional expertise to guide them through complex medical landscapes. As we continue to advance our healthcare systems toward more human-centric models, embracing shared planning will undoubtedly play a pivotal role in realizing healthier individuals within thriving communities—a testament to how thoughtful collaboration can transform lives for the better.

****Addressing Chronic Dental Issues**:** Adults may seek orthodontics to resolve long-standing dental problems such as bite issues (overbites, underbites, crossbites) that can lead to jaw pain, headaches, and digestive difficulties if left untreated.

Collaborative approaches in the realm of healthcare, particularly when it comes to shared planning for improved patient experiences, represent a transformative shift from traditional, siloed practices. This method emphasizes teamwork across various healthcare disciplines and with patients themselves, fostering an environment where diverse perspectives converge to enhance care delivery. The benefits of such collaborative efforts are manifold, touching on quality of care, efficiency, patient satisfaction, and even financial outcomes.

Firstly, collaboration significantly enhances the quality of care that patients receive. By bringing together doctors, nurses, therapists, social workers, and other essential healthcare professionals—alongside patients and their families—the planning process becomes comprehensive. Each participant contributes unique insights into the patient's needs and circumstances. This holistic view ensures that treatment plans are not only medically sound but also consider the broader context of a patient's life, including psychological and social factors. Such comprehensive care planning leads to more effective outcomes and a greater likelihood of adherence to treatment regimens.

Moreover, collaborative approaches streamline processes within healthcare settings, leading to improved efficiency. When teams communicate openly and regularly share information about patient progress and needs, duplication of efforts is minimized while critical steps are ensured. This coordination can lead to quicker responses to changes in a patient's condition and smoother transitions between different stages of care or facilities—critical aspects that can significantly affect patient outcomes and satisfaction.

Patient satisfaction stands as another crucial benefit of shared planning initiatives. When patients feel involved in their care plan, they often experience higher levels of satisfaction. Being heard and having their preferences considered cultivates a sense of partnership with their healthcare providers rather than mere passivity as recipients of care. This

engagement empowers patients, making them active participants in decisions affecting their health journey—a dynamic that not only boosts morale but also encourages better engagement with treatment plans.

Additionally, collaborative approaches have shown promise in optimizing resource use within healthcare systems, contributing positively to financial sustainability without compromising quality. By ensuring that all angles are covered through collective expertise and by reducing inefficiencies through streamlined communication and processes, these methods help manage costs effectively. This is crucial given the ongoing challenges faced by healthcare systems globally regarding resource allocation and expenditure control.

In essence, embracing collaborative approaches in shared planning for improved patient experiences unlocks a synergy among stakeholders that enriches every aspect of healthcare delivery—from the quality and effectiveness of care to the efficiency of operations and ultimately to enhancing patient satisfaction and well-being. As we navigate the complexities of modern healthcare challenges, this human-centric model proves invaluable in crafting solutions that truly serve those at the heart of our endeavors: the patients themselves.

****Correcting Speech**

Impediments:** Misaligned teeth or jaw structures can

**contribute to speech
difficulties; orthodontic
treatment can correct these
issues, improving articulation
and overall communication
skills.**

Shared planning, particularly in the context of orthodontic treatment for children, stands as a pivotal approach to enhancing communication among all stakeholders—parents, children, and orthodontists. This collaborative process isn't merely about scheduling appointments or discussing treatment options; it's a holistic strategy aimed at improving patient experiences by fostering an environment of mutual understanding, respect, and involvement. Let's delve into how shared planning can weave a more cohesive tapestry for everyone involved in the journey toward a healthier smile.

Firstly, shared planning opens up a dialogue that is essential for setting realistic expectations. When parents and orthodontists collaborate to outline treatment goals and timelines, they provide children with a clearer picture of what to expect. This transparency

reduces anxiety often associated with dental procedures, especially for children who might fear the unknown. By being included in discussions about potential outcomes and the steps required to achieve them, children feel more engaged and less intimidated by the process. This engagement can lead to better cooperation during treatments, making visits to the orthodontist less stressful for everyone involved.

Moreover, involving parents in the planning stages ensures that their insights are considered crucial components of the treatment plan. Parents often have a deep understanding of their child's personality, habits, and responses to stress or discomfort. Sharing this knowledge with orthodontists allows for personalized care that caters not just to dental needs but also to emotional and psychological well-being. For instance, if a child tends to be particularly anxious about appointments, strategies can be developed collectively—like incorporating distraction techniques or scheduling shorter sessions—to create a more comfortable experience tailored specifically to that child's needs.

Furthermore, shared planning promotes continuous communication channels among all parties. Regular updates on progress can help keep parents informed while allowing them to voice concerns or observations promptly. This ongoing dialogue ensures that any adjustments needed in treatment plans can be made swiftly and effectively without waiting until problems escalate. Such proactive communication helps maintain alignment between home care instructions given by orthodontists and actual practices at home by both parents and children—a critical factor in achieving successful outcomes.

Additionally, when children see their parents actively participating in their care plan discussions, it instills a sense of importance regarding oral health from an early age. Children learn valuable lessons about responsibility and teamwork through witnessing collaborative efforts towards achieving health goals. This learning experience extends beyond orthodontic treatment; it lays down foundational principles of health management that they will carry into adulthood.

In conclusion, shared planning serves as a cornerstone for improved patient experiences within orthodontic care by fostering enhanced communication among parents, children, and orthodontists alike. It transforms what could be perceived as mere medical appointments into cooperative endeavors grounded in mutual respect and understanding. By embracing this approach we not only refine the mechanics of dental care but significantly elevate the overall experience creating happier patients, more satisfied parents, and ultimately, healthier smiles that last a lifetime.

****Preventive Measures**

Against Tooth Wear:**

Properly aligned teeth are less prone to excessive wear and tear; adults may pursue orthodontic treatment to prevent premature tooth

degradation and associated costs of restorative dentistry.

Collaboration, especially in the context of healthcare, particularly when focusing on shared planning for improved patient experiences, plays a pivotal role in fostering a supportive environment that significantly reduces anxiety among young patients. This approach isn't merely about involving various stakeholders—it's about weaving a tapestry of support that makes the often intimidating world of medical care feel more approachable and less daunting. Let's unpack how this collaborative spirit transforms the experience for young patients.

First off, collaboration brings together diverse professionals—doctors, nurses, child life specialists, psychologists, and even family members—each contributing their unique perspective and expertise. This multidisciplinary approach ensures that the care plan addresses not just the medical needs but also the emotional and psychological well-being of the patient. For young patients, who might struggle to articulate their fears or anxieties about medical procedures or hospital stays, this comprehensive support system acts as a safety net. Knowing that every aspect of their care is thoughtfully coordinated can instill a sense of security and understanding that they are not alone in this journey.

Moreover, shared planning sessions often include discussions about the patient's preferences, fears, and questions. By actively involving young patients in conversations about their treatment plans, healthcare providers empower them to have a say in their own care. This empowerment is crucial; it transforms the passive role of receiving

treatment into an active participation in decision-making processes. When children feel heard and involved, their anxiety diminishes because they gain a sense of control over their situation—a powerful antidote to fear and uncertainty.

Additionally, collaboration encourages an environment where open communication thrives. When healthcare teams regularly meet to discuss patient progress and concerns, it fosters transparency and trust between providers and patients (and their families). Young patients observe this open dialogue and feel more comfortable expressing their own worries or asking questions without fear of judgment. This culture of openness validates their feelings and experiences, making them feel understood and valued as individuals rather than just cases to be managed.

Social support also plays a vital role within this collaborative framework. Involving family members in planning sessions acknowledges the critical role they play in a child's life—and consequently, in their healing process. Families can provide comfort through familiarity; knowing loved ones are actively engaged can significantly ease anxiety for young patients who might otherwise feel isolated or overwhelmed by unfamiliar surroundings and routines.

Lastly, let's consider how collaboration extends beyond immediate medical needs to encompass educational elements tailored for young patients. Interactive workshops or preparatory sessions led by child life specialists can demystify hospital procedures using age-appropriate language and activities. Such initiatives not only educate but also entertain, turning potentially anxiety-provoking scenarios into manageable experiences through playful engagement—a brilliant strategy to normalize what might otherwise seem alienating or scary.

In essence, collaboration within shared planning for improved patient experiences serves as a powerful tool for creating an environment where young patients feel supported rather

than overwhelmed by anxiety. It harnesses the collective expertise of various professionals while prioritizing open communication and active participation from both patients and their families. By transforming healthcare from an impersonal service into a collaborative endeavor centered around understanding individual needs and fostering emotional well-being, we can significantly mitigate anxiety—allowing youthful spirits to navigate medical challenges with greater confidence

****Facilitating Better Chewing Efficiency**
Orthodontic treatment can improve bite function and alignment, allowing for more efficient chewing which is crucial for digestion and overall**

nutritional health in adulthood.

Involving children in treatment decisions is a pivotal yet often overlooked aspect of shared planning aimed at improving patient experiences, particularly within pediatric healthcare settings. This approach not only respects the evolving autonomy of children but also enriches the therapeutic process by integrating their perspectives and preferences into care plans. The rationale behind this inclusive strategy is multifaceted, touching on psychological, developmental, and ethical dimensions that underscore its importance in modern healthcare practices.

Firstly, considering the psychological impact, children who are engaged in discussions about their treatment feel a sense of empowerment and control over their health journey. This engagement can significantly mitigate anxiety and fear associated with medical procedures and hospital environments, which are often intimidating for young patients. By explaining treatments in age-appropriate language and involving them in decision-making processes, healthcare providers can alleviate some of the psychological burdens that accompany illness and treatment, fostering a more positive mindset crucial for recovery.

Moreover, from a developmental standpoint, involving children in these discussions aligns with their cognitive growth stages. As children mature, their ability to understand complex information and participate in meaningful dialogue improves. By respecting these developmental milestones and adapting communication accordingly, healthcare

professionals not only facilitate better comprehension of health issues but also encourage active participation in their own care. This practice nurtures self-advocacy skills early on—a trait invaluable as they transition into adulthood managing their health independently.

Ethically speaking, the principle of respect for persons underpins the justification for involving children in treatment decisions. This principle emphasizes dignity and autonomy, suggesting that individuals capable of making informed choices should have a say in matters affecting their well-being. While children may not always possess the full capacity to make definitive medical decisions independently, incorporating their views demonstrates respect for their emerging autonomy and personal agency—a cornerstone of ethical patient care.

Incorporating children into treatment planning also enhances communication between healthcare providers, families, and patients themselves. It encourages collaborative discussions where everyone's insights contribute to a comprehensive care plan tailored to the child's unique needs and desires. Such an environment fosters trust among all parties involved—children feel heard; parents gain reassurance through shared decision-making; and healthcare providers craft more personalized treatment strategies that consider emotional as well as physical health aspects.

To effectively implement this approach, it's essential for healthcare systems to develop guidelines supporting age-appropriate communication techniques and tools for assessing children's understanding and preferences accurately. Training programs for medical staff can focus on enhancing interpersonal skills necessary for engaging with young patients sensitively yet effectively. Additionally, creating resources—like visual aids or storytelling methods—that cater to different age groups can bridge communication gaps between complex medical information and child-friendly explanations.

In conclusion, involving children in treatment decisions is not merely an act of courtesy but a fundamental shift towards patient-centered care that acknowledges the individuality of each young patient. By prioritizing inclusivity from an early stage of healthcare encounters, we foster environments where growing individuals feel valued participants rather than passive recipients of care. This approach promises richer therapeutic experiences characterized by enhanced cooperation between families, caregivers, and young patients themselves—a cornerstone for improved patient experiences across pediatrics comprehensively addressing emotional well-being alongside physical health outcomes.

Shared planning for improved patient experiences is a concept that hinges on the fundamental belief that when patients are actively involved in their healthcare journey, the outcomes are not just medically better but profoundly more satisfying. This approach emphasizes the importance of crafting explanations that resonate with a patient's age and understanding, and equally crucial, involving them in key decision-making processes. The significance of these elements cannot be overstated, as they form the bedrock upon which trust, empowerment, and ultimately, enhanced patient experiences are built.

Firstly, let's unpack the power of age-appropriate explanations. Imagine a young child facing a medical procedure; their understanding and emotional readiness differ vastly from an adult or an elderly person. Tailoring explanations to match a patient's developmental stage ensures they grasp what's happening in a way that's comprehensible and less intimidating. For children, using simple language, visuals, and relatable analogies can demystify complex medical jargon. For adults, while simplicity is key, providing detailed information about procedures, potential outcomes, and risks empowers them to make informed decisions. Elderly patients might benefit from explanations that consider possible cognitive changes or hearing impairments, ensuring clarity without overwhelming them with details. By doing so, healthcare providers bridge the communication gap, instilling confidence and reducing anxiety—a cornerstone of positive patient experiences.

Moreover, involving patients in decision-making isn't merely about ticking boxes for compliance; it's about recognizing each individual as an active participant in their healthcare journey. When patients are included in planning their treatment—be it choosing between different therapeutic options or discussing lifestyle changes—they feel valued and respected as collaborators rather than passive recipients of care. This sense of agency significantly impacts their engagement levels; studies show that involved patients adhere better to treatment plans and report higher satisfaction with their care experiences. It fosters a partnership where patients feel empowered to voice concerns, ask questions, and share preferences—ultimately leading to personalized care plans that align closely with their values and lifestyles.

The synergy between age-appropriate communication and shared decision-making creates an environment ripe for improved patient experiences. It encourages transparency from healthcare providers who strive to meet patients where they are—cognitively and emotionally—and invites them into a dialogue about their health journey. Such an approach not only enhances understanding but also fortifies the trust between patient and provider—a critical component of effective healthcare delivery. When individuals feel understood and included in decisions affecting their well-being, they're more likely to navigate their health challenges with resilience and positivity.

In conclusion, shared planning for improved patient experiences thrives on the twin pillars of age-appropriate explanations and meaningful involvement in decision-making processes. These strategies transform healthcare from a one-dimensional service into a collaborative journey where every patient feels seen, heard, and valued at every step of the way. As we continue to evolve our approaches to care delivery, let us remember that at its heart lies the human element—our capacity to connect, communicate, and empower those we serve.

When we embark on the journey of discussing "Shared Planning for Improved Patient Experiences," especially with children, it's crucial to adopt an approach that's not only

educational but also engaging and relatable. The goal here is to ensure that young minds not only understand the importance of shared planning in healthcare settings but also feel a personal stake in their own care experiences. To achieve this, employing a variety of tools and methods can make the concept vivid and memorable. Let's explore some effective strategies tailored to captivate children while imparting valuable insights into shared planning.

Storytelling with Visual Aids

One powerful method is through storytelling, complemented by visual aids. Picture books or comic strips can be incredibly effective. Imagine a story where a character, perhaps a brave little superhero, visits the hospital. Each page could illustrate different steps of shared planning—from discussing concerns with doctors and nurses to choosing between fun activities during recovery. Visual aids like these not only capture attention but also help children visualize themselves in similar situations, making the concept more relatable and less intimidating.

Interactive Models and Puppets

Interactive models and puppets bring an element of play into learning. For instance, using dolls or stuffed animals as 'patients' allows children to role-play scenarios involving shared planning with healthcare providers represented by adults dressed as doctors or nurses. This hands-on approach lets kids experiment with communication, expressing their fears or wishes in a safe environment. The tangible nature of models helps solidify abstract concepts into something concrete they can manipulate and understand better.

Digital Tools and Games

In our increasingly digital world, leveraging technology can be highly engaging for children. Interactive games or apps designed around themes of health and patient experiences offer immersive learning opportunities. These platforms could simulate patient visits where children must decide which questions to ask healthcare providers based on pre-set criteria related to shared planning principles. Such gamified learning not only educates but also makes the process enjoyable, encouraging repeated engagement to master the concepts further.

Collaborative Art Projects

Artistic expression is another powerful tool for engagement and understanding among children. Organizing collaborative art projects where kids create murals or posters about what they believe makes a good patient experience can be both fun and enlightening. This activity encourages discussion about elements like clear communication, empathy from healthcare providers, and involving family members in decision-making processes—core components of shared planning. Displaying these creations in waiting areas or classrooms reinforces the message while giving children a sense of pride in contributing to important conversations about healthcare experiences.

Conclusion

Engaging children in discussions about "Shared Planning for Improved Patient Experiences" requires creativity and adaptability in our teaching methods. By integrating storytelling with visual aids, interactive models and puppetry, digital tools through games, and

collaborative art projects,

Setting Realistic Expectations Together: A Pillar for Shared Planning in Enhancing Patient Experiences

In the intricate dance of healthcare, where patients, families, and medical professionals intertwine their efforts to navigate the often complex landscape of health and healing, the significance of setting realistic expectations cannot be overstated. This principle forms a cornerstone in the broader strategy of shared planning aimed at improving patient experiences. At its heart, this approach is about fostering an environment where all parties involved align their hopes and understandings to create a journey that is not only medically sound but also emotionally and psychologically supportive.

The essence of shared planning revolves around collaboration—bringing together diverse perspectives to chart a course that accommodates the multifaceted needs of patients. When it comes to setting realistic expectations, it's crucial to acknowledge that each patient's experience is unique, influenced by their personal health history, emotional state, and sometimes even their cultural background. By engaging patients and their families in discussions early on, healthcare teams can gain invaluable insights into what matters most to them. This might include recovery timelines, pain management preferences, or even simple aspects like maintaining a sense of normalcy during hospital stays.

Realistic expectations are built on honest communication. It involves clearly explaining treatment options, potential outcomes, including both successes and possible challenges. By doing so, healthcare providers empower patients with knowledge that reduces anxiety stemming from uncertainty. When patients understand what to expect—be it the rigors of rehabilitation after surgery or the nuances of managing a chronic condition—they are better equipped to prepare mentally and emotionally. This preparedness significantly enhances their experience within the healthcare system.

Moreover, setting realistic expectations opens avenues for proactive problem-solving. When everyone involved has a clear understanding of what can be realistically achieved within given timeframes or circumstances, identifying potential hurdles becomes easier. This allows for timely interventions and adjustments in care plans before issues escalate into larger concerns affecting patient satisfaction and outcomes negatively.

Additionally, fostering an environment where expectations are set collaboratively nurtures trust between patients and healthcare providers—a vital ingredient for successful healing journeys. Trust encourages open dialogue about concerns or dissatisfactions as they arise rather than letting them fester silently. This continuous feedback loop not only improves individual patient experiences but also contributes valuable insights for systemic improvements across healthcare services.

In essence, "Setting Realistic Expectations Together" is more than just a strategy; it's a commitment to partnership in healthcare—a recognition that improved patient experiences emerge from shared understanding and collective effort. As we continue refining our approaches to healthcare delivery, embracing this principle will remain pivotal in crafting experiences that honor the dignity and needs of every individual under our care. Through collaboration rooted in honesty and mutual respect, we pave the way for healing environments where hope thrives alongside realism—a balance essential for true progress in patient-centered care.

Setting achievable treatment goals in a healthcare context, particularly when involving both parents and children, is a delicate yet crucial process that embodies the essence of shared planning for improved patient experiences. This collaborative approach not only ensures that the treatment plan is tailored to the unique needs of the child but also fosters a supportive environment where every family member feels heard and involved. Here's how this inclusive process can unfold, step by step, to create a harmonious and effective pathway toward health and wellness.

Step 1: Establish Open Communication

The journey begins with creating an atmosphere of open and honest communication. It's essential to hold a meeting where all parties—parents, children, and healthcare providers—feel welcome to express their thoughts, concerns, and expectations without judgment. This initial dialogue sets the tone for collaboration and mutual respect, making everyone feel valued in the decision-making process.

Step 2: Understand the Child's Perspective

Children often have unique insights into what they need or desire regarding their treatment. Engaging them directly—depending on their age and understanding—through conversations or play therapies can reveal their feelings about their condition and what they hope to achieve through treatment. This step is pivotal; it ensures that the child's voice is central to the planning process.

Step 3: Gather Input from Parents

Parents bring invaluable perspectives based on their observations of their child's daily needs and challenges. Their input should be sought regarding practical considerations at home, emotional support required, and any previous experiences with treatments that were successful or less so. Understanding these viewpoints helps in aligning professional recommendations with real-life scenarios at home.

Step 4: Collaboratively Identify Goals

With insights from both parents and children, the next step is to collaboratively identify achievable treatment goals. These should be Specific, Measurable, Achievable, Relevant, and Time-bound (SMART). For instance, instead of a vague goal like "feeling better," a SMART goal might be "reduce fever episodes to once a month within three months." This clarity ensures everyone understands what success looks like.

Step 5: Customize Treatment Plans

Once goals are set, tailor the treatment plan to accommodate these objectives while considering lifestyle factors discussed earlier. This might involve adjusting medication schedules to fit family routines or incorporating specific activities that aid recovery while being enjoyable for the child. Personalization here is key; one-size-fits-all approaches often falter in fostering engagement and compliance.

Step 6: Establish Support Systems

Recognizing that achieving these goals won't solely rely on medical interventions but also on emotional and practical support systems is crucial. Encourage family involvement in supporting each other during treatments—whether it's attending appointments together or helping with home care routines—and explore community resources like support groups or educational workshops that can provide additional assistance and encouragement.

Step 7: Monitor Progress Together

Setting up regular check-ins where progress towards goals can be reviewed collectively ensures transparency and adjustability in the plan as needed. These sessions should

remain positive spaces focusing on achievements no matter how small while reassessing strategies if certain milestones aren't met as expected—a flexible approach keeps hopes alive without fostering frustration or disappointment.

Step 8: Celebrate Successes Big and Small

Lastly, acknowledging successes along the way is vital for maintaining morale and motivation across all family members involved in this journey. Cele

The journey of enhancing patient experiences through shared planning is much like embarking on a grand expedition—a voyage where every participant, from healthcare providers to patients and their families, plays a pivotal role in navigating the complex terrain of medical care. At the heart of this expedition lies a crucial compass: realistic timelines and milestones. These elements are not just administrative tools; they are the very heartbeat that sustains motivation and fosters collaboration throughout the shared planning process.

To appreciate the significance of realistic timelines and milestones, one must first understand the intricate dynamics at play in healthcare environments. Patient experiences are multifaceted, influenced by everything from communication quality to the efficiency of service delivery. Shared planning emerges as a beacon in this complexity, bringing together diverse perspectives to craft care paths tailored to individual needs. However, without a clear roadmap—one marked by achievable milestones and grounded in realistic timeframes—these collaborative efforts risk floundering amidst uncertainty and disheartenment.

Realistic timelines act as guiding stars, illuminating what can be reasonably expected within certain periods. They provide clarity amidst the often murky waters of healthcare

operations, where delays and unforeseen challenges are commonplace. When teams—comprising doctors, nurses, administrative staff, and patients—are presented with achievable deadlines for various stages of plan implementation, it instills a sense of direction and purpose. This clarity prevents burnout by ensuring that efforts are aligned with feasible outcomes rather than chasing unattainable perfection within unrealistic time constraints.

Milestones serve as checkpoints along this journey—moments to pause, reflect, and celebrate progress made. Each milestone reached fuels motivation; it's tangible proof that collective effort translates into tangible improvements in patient experiences. These moments of recognition encourage team members to persist through challenges knowing their hard work is making a difference in someone's life—a powerful motivator indeed. Moreover, these achievements foster an environment ripe for learning and adaptation; teams can analyze what worked well and what didn't, refining their approaches for future endeavors with renewed vigor and insight.

Furthermore, shared planning bolstered by realistic timelines and milestones enhances accountability among all stakeholders involved. When everyone understands what is expected at each phase—and when those phases are scheduled realistically—it cultivates a culture where each member feels responsible for contributing to success. This shared sense of responsibility nurtures deeper engagement from all parties involved—patients feel heard and included in decisions about their care; providers find satisfaction in delivering on promises made during these collaborative sessions; administrators observe smoother operations that align with strategic goals set forth by the institution.

In essence, realistic timelines and well-defined milestones breathe life into shared planning initiatives aimed at improving patient experiences. They act as stabilizers against frustration and demotivation by providing structure amidst complexity while simultaneously serving as motivational beacons that guide teams toward common

objectives. As we navigate the evolving landscape of healthcare delivery focused increasingly on patient-centricity, let us remember: it's not merely about reaching destinations but also about savoring each step taken together towards better health outcomes—and it's through these structured yet flexible pathways that we truly embark on journeys marked by meaningful progress and sustained motivation for all involved.

Customizing treatment plans to cater to individual needs is a cornerstone of modern healthcare, deeply embedded in the philosophy of shared planning for improved patient experiences. This approach recognizes that every patient is unique, carrying with them a tapestry of personal health histories, preferences, and lifestyle factors that significantly influence their journey toward wellness. By embracing this individuality, healthcare providers can craft treatment plans that are not just effective but also resonate on a personal level with the patients they serve.

The essence of customization lies in understanding the whole person—not just their illness or condition. It involves gathering comprehensive information about a patient's medical history, genetic predispositions, lifestyle choices, psychological state, and social environment. This holistic view allows healthcare professionals to tailor interventions that align closely with an individual's specific needs and circumstances. For instance, a dietary plan for managing diabetes might consider not only the nutritional aspects but also cultural food preferences and the patient's ability to access certain foods, ensuring adherence and effectiveness without undue hardship.

Moreover, customized treatment plans foster a collaborative relationship between patients and healthcare providers. Engaging patients in the planning process empowers them by making them active participants in their care rather than passive receivers of treatments. This shared decision-making enhances patient satisfaction and compliance since individuals are more likely to follow through with plans they have helped create. It also opens channels for ongoing dialogue where patients can voice concerns or adjustments needed as their condition evolves or as they experience different outcomes.

from their treatment.

The benefits extend beyond improved compliance; personalized care can lead to better health outcomes overall. When treatments are aligned with an individual's unique profile, there's often an increase in efficacy and a reduction in adverse effects or unnecessary procedures. This precision medicine approach not only focuses on curing diseases but also on enhancing quality of life throughout the treatment journey.

However, achieving this level of customization requires resources—time for thorough assessment, expertise in various disciplines to address multifaceted health needs, and sometimes advanced technology for personalized diagnostics and monitoring. Healthcare systems must invest in training staff to be more attuned to individual patient nuances and integrating diverse data sources seamlessly into care planning processes.

In conclusion, customizing treatment plans for individual needs is not merely an idealistic goal but a practical approach that significantly enhances patient experiences in shared care planning initiatives. It reflects respect for each person's unique story while aiming for optimal health outcomes through collaboration and personalization. As we continue advancing in medical science and technology, prioritizing this individualized approach will remain crucial in delivering compassionate, effective healthcare tailored precisely to those we strive to help—our patients themselves.

Shared planning in the realm of pediatric dentistry, particularly when focusing on improving patient experiences, represents a paradigm shift from a one-size-fits-all approach to a more nuanced, personalized strategy that caters to each child's unique dental and emotional needs. This method recognizes the fundamental truth that every child is different; they come with their own set of challenges, fears, dental health histories, and developmental stages that influence how they experience dental care. By embracing shared planning—a collaborative process involving dentists, parents, and sometimes even

the children themselves—we can craft tailored approaches that not only address immediate dental concerns but also foster a positive relationship with dental care that lasts a lifetime.

The essence of shared planning lies in communication and cooperation among all stakeholders involved in a child's dental health. Dentists, armed with their expertise, initiate discussions with parents to understand the broader context of the child's health and emotional well-being. This conversation might reveal specific anxieties about dental visits stemming from past experiences or general fears common among children. Simultaneously, dentists share their observations and recommendations based on the child's current dental condition, growth patterns, and any special needs that might affect treatment approaches.

Incorporating the child's perspective is pivotal. Depending on their age and comfort level, involving children in the planning process can significantly alleviate anxiety by giving them a sense of control over what happens to them. Simple acts like discussing what they can expect during their visit or allowing them to choose certain aspects—like picking a favorite character sticker to place on their chart—can transform an intimidating experience into something more comfortable and familiar.

Tailored approaches resulting from shared planning may manifest in various ways: adjusting appointment times to avoid peak anxiety periods for some children; using storytelling or gentle explanations suited to a child's understanding level; employing specialized techniques or tools designed for sensitive mouths or those with special needs; or creating distraction methods like using technology or engaging games during procedures. These personalized strategies acknowledge each child's unique threshold for stress and adapt accordingly, ensuring that every visit is as comfortable as possible while achieving necessary dental care objectives.

Moreover, shared planning encourages continuous feedback loops where parents can voice their observations about their child's responses at home after visits, enabling adjustments to be made promptly for subsequent appointments. Dentists can utilize this information to fine-tune their approach further, creating an evolving plan that respects both the changing needs of the child's oral health and their emotional journey through these experiences.

In conclusion, shared planning for improved patient experiences in pediatric dentistry exemplifies compassionate care by weaving together professional insight with personal considerations specific to each young patient. It emphasizes collaboration as key to unlocking tailored approaches that not only address immediate dental needs but also nurture positive associations with oral healthcare from an early age. Through thoughtful engagement with families and careful attention to individual differences in dental health and emotional resilience, we pave the way for healthier smiles grounded in trust and understanding—a foundation crucial for lifelong oral wellness.

Personalized treatment, especially when woven into the fabric of shared planning, has become a beacon of hope in transforming patient experiences across various healthcare settings. This approach underscores the significance of tailoring medical care to the unique needs, preferences, and life circumstances of each individual. To truly appreciate its impact, let's delve into a couple of compelling case studies that illuminate how shared planning can lead to remarkable success in personalized treatment.

Case Study 1: Chronic Pain Management

Consider Sarah, a 45-year-old woman battling chronic back pain for over a decade. Traditional treatment methods had offered little relief, leading to frustration and a sense of helplessness. Enter shared planning, where her healthcare team—comprising a primary care physician, physical therapist, psychologist, and an occupational therapist—came

together to craft a comprehensive plan tailored specifically to Sarah's lifestyle and pain triggers.

During their initial meeting, the team didn't just focus on medication or surgery; they explored Sarah's daily routines, work environment, and emotional well-being. They discovered that long hours at her desk job exacerbated her pain and that stress significantly heightened her discomfort levels. Armed with this insight, they devised a multifaceted approach: ergonomic adjustments at work, targeted physical therapy sessions focusing on strengthening core muscles, mindfulness practices for stress management, and periodic medication reviews to ensure efficacy without over-reliance on painkillers.

The shared planning process empowered Sarah to take an active role in her treatment journey. Regular follow-ups allowed for adjustments based on her progress and feedback. Over time, not only did Sarah experience substantial reductions in pain but also regained control over her life. The collaborative effort created a supportive network that addressed not just the physical aspect but also the emotional burdens associated with chronic illness—a testament to how personalized treatment through shared planning can profoundly enhance patient experiences.

Case Study 2: Diabetes Management Among Adolescents

Shifting focus to another scenario involving young patients—let's look at Jake, a 16-year-old diagnosed with Type 1 diabetes. Managing diabetes during adolescence comes with unique challenges; it's not merely about regulating blood sugar but also navigating social dynamics and emerging independence. Here again, shared planning played a pivotal role in personalizing Jake's treatment plan around his lifestyle and developmental needs.

Jake's care team included an endocrinologist, dietitian, school nurse educator, and importantly—his parents—and even involved Jake directly in discussions from an early stage to foster ownership over his health management. Recognizing that peer interactions heavily influence dietary choices and activity levels for adolescents, the team integrated strategies like using technology (a user-friendly app for tracking glucose levels) into Jake's routine without making it feel overly restrictive or adult-centric. They discussed realistic goals aligned with his school schedule and social activities while ensuring his family was equipped to support him without hovering excessively—a delicate balance essential during teenage years.

The personalized approach paid dividends; Jake felt understood rather than just treated as a case file. His blood sugar management improved significantly as he learned to make informed choices that fit seamlessly into his life rather than being seen as burdensome obligations. More importantly, he developed confidence in managing his condition independently—a crucial skill for long-term health success beyond adolescence.

These case studies exemplify how personalized treatment through shared planning not only addresses medical needs but transforms patient experiences by fostering engagement empowerment and holistic care that respects each

Incorporating Feedback Loops in Shared Planning for Improved Patient Experiences: A Human Approach

In the ever-evolving landscape of healthcare, the notion of shared planning has emerged as a pivotal strategy aimed at enhancing patient experiences. At its core, shared planning involves collaborative efforts among healthcare providers, patients, and sometimes their families or caregivers, to craft personalized care plans that resonate with the unique needs and preferences of each individual. Central to this approach is the integration of feedback loops—a dynamic mechanism designed to ensure continuous improvement and

responsiveness to patient experiences. This essay explores how incorporating feedback loops into shared planning not only enriches the quality of care but also fosters a deeper sense of partnership between healthcare providers and patients.

Feedback loops are essentially mechanisms that facilitate the collection, analysis, and application of patient feedback within the planning process. These loops can manifest in various forms—ranging from direct conversations post-care sessions to structured surveys and digital platforms that allow real-time input. The essence lies not merely in gathering data but in creating an environment where every piece of feedback is valued and acted upon, reinforcing a culture of responsiveness and respect for patient perspectives.

The first significant benefit of integrating feedback loops into shared planning is the enhancement of personalized care. By actively listening to patient feedback, healthcare teams gain invaluable insights into what works well and what areas need refinement from the patient's viewpoint. This understanding allows for tailored adjustments to care plans that better align with individual preferences and life circumstances, contributing significantly to improved satisfaction levels. Patients who feel their voices are heard are more likely to engage positively with their treatment plans, adhering more closely to prescribed regimens and actively participating in their health journey.

Moreover, feedback loops foster a climate of transparency and trust between patients and providers. When patients see that their opinions lead to tangible changes in their care experience, it strengthens their confidence in the healthcare system. This trust is foundational for effective patient-provider relationships; it encourages open communication about concerns or needs that might otherwise go unaddressed. In turn, this transparency helps identify systemic issues within healthcare delivery that might negatively impact patient experiences across broader populations—not just individuals—and prompts necessary reforms.

Additionally, these loops serve as educational tools for both patients and providers alike. Patients become more informed about their health conditions and treatment options as they provide insights based on lived experiences. Conversely, healthcare professionals gain deeper understanding through direct engagement with patients' perspectives, enabling them to refine practices based on real-world implications rather than theoretical assumptions alone. This mutual learning process enriches shared planning by ensuring it remains grounded in practical realities rather than abstract objectives.

However, implementing effective feedback loops requires careful consideration and strategic design. It necessitates creating accessible channels for feedback collection that accommodate diverse patient needs—considering factors like language barriers or technological proficiency—and ensuring confidentiality and respect in all interactions. Moreover, organizations must establish clear pathways for translating feedback into actionable improvements while maintaining accountability for these changes within the system. Without such thoughtful execution, there's a risk that feedback becomes mere lip service rather than a catalyst for meaningful change.

In conclusion, incorporating feedback loops into shared planning represents a powerful stride toward improving patient experiences by promoting personalization, trust, transparency, and ultimately, positive Outcomes.

Shared planning in the context of healthcare, particularly when focusing on ensuring patient satisfaction and enhancing experiences throughout treatment, hinges critically on the establishment of robust mechanisms for ongoing feedback. This approach is not merely about collecting patient opinions; it's about crafting a dynamic dialogue that allows for continuous improvement and adaptation of treatment plans to better suit individual needs. Here's how such a system can be effectively implemented, ensuring that treatment journeys are as personalized and satisfying as possible.

Regular Check-ins: The Foundation of Feedback

The cornerstone of ongoing feedback mechanisms is regular, scheduled check-ins between patients and their healthcare providers. These shouldn't be mere formalities but opportunities for open conversation. During these sessions, patients should feel encouraged to voice any discomfort, concerns, or positive experiences they've encountered since the last visit. Healthcare providers, in turn, should actively listen, demonstrating empathy and understanding while noting down key points for consideration in adjusting treatment plans. These check-ins can occur at various touchpoints—after significant milestones in treatment, during routine appointments, or even via scheduled phone calls or video consultations for those with mobility challenges or busy schedules.

Utilizing Technology for Seamless Communication

In our increasingly digital world, leveraging technology can significantly enhance feedback mechanisms. Patient portals equipped with secure messaging systems allow individuals to communicate with their care teams outside traditional office hours. This immediacy ensures that concerns are addressed promptly, fostering a sense of being heard and valued. Additionally, incorporating digital surveys tailored to specific stages of treatment can provide structured yet flexible feedback opportunities without overwhelming patients with lengthy questionnaires at every visit. These tools not only streamline communication but also ensure data is collected systematically for analysis and action.

Empowering Patients Through Education

For feedback to be meaningful and actionable, patients must feel informed and empowered regarding their health journey. Educational sessions—whether one-on-one with healthcare providers or through group workshops—can demystify treatment processes and outcomes, encouraging patients to engage more deeply in shared planning discussions. Understanding the rationale behind certain treatments or changes in plan helps patients articulate their experiences more accurately and contribute constructively to the decision-making process regarding their care.

Adaptive Treatment Planning: A Response to Feedback

The true essence of shared planning lies in its adaptability based on the feedback received. Treatment plans should not be static documents but living guides that evolve with the patient's progress and changing needs. Healthcare teams must cultivate a mindset of flexibility—ready to pivot therapies, medication regimens, or even support services based on ongoing input from patients. This might involve revisiting goals regularly to ensure they remain aligned with what matters most to the patient at each stage of their journey. It's crucial that these adaptations are communicated clearly back to the patient, reinforcing their active role in shaping their care experience.

Creating a Culture of Openness and Responsiveness

Lastly, fostering an organizational culture that values transparency and responsiveness is pivotal for successful shared planning initiatives aimed at improving patient experiences through ongoing feedback mechanisms. Everyone involved—from front-line staff to senior management—must recognize the importance of patient input as vital data driving improvements across all facets of care delivery. Encouraging a culture where feedback is seen as an asset rather than merely data points empowers both staff and patients alike to work collaboratively toward optimal outcomes grounded in personal satisfaction and well-

being.

In conclusion, by implementing regular check-ins paired with technological aids for seamless communication

Shared planning for improved patient experiences is a multifaceted approach that hinges on the foundational principle of communication—particularly, the regular check-ins between healthcare providers, patients, and their families. In an era where patient-centered care is not just a goal but a necessity, these check-ins serve as vital touchpoints that ensure everyone involved in a patient's care journey is aligned, informed, and engaged. The importance of such practices cannot be overstated; they are the linchpin in fostering an environment where patients feel heard, understood, and cared for, thereby significantly enhancing their overall experience within the healthcare system.

Regular check-ins create an open channel for dialogue. They allow patients and their families to voice concerns, ask questions, and express feelings about their treatment plans or hospital stay without feeling rushed or dismissed. This open line of communication is crucial because it empowers patients to take an active role in their health management. When individuals feel they have a say in their care plan, their adherence to treatments often improves, and so does their satisfaction with the service received.

Moreover, these interactions provide healthcare professionals with invaluable insights into the patient's perspective. Understanding a patient's concerns—be it related to pain management, emotional well-being, or logistical issues like appointment scheduling—enables staff to tailor services more effectively. It transforms generic care protocols into personalized experiences that resonate with each individual's unique needs and circumstances. This personalized approach not only addresses immediate concerns but also builds trust and rapport between patients and caregivers—a cornerstone of healing

and recovery.

In addition to addressing immediate issues, regular check-ins foster a sense of community and support around the patient. Involving parents or guardians in these discussions ensures that all voices at the table are considered when making decisions about care plans. This inclusivity can alleviate anxiety for families navigating complex health situations and ensures that every aspect of the patient's well-being—physical, emotional, and social—is taken into account during shared planning sessions.

The benefits extend beyond individual encounters; they contribute significantly to institutional learning and improvement. Feedback gathered through these interactions can highlight systemic areas needing enhancement or innovation within healthcare services. By listening closely to patient experiences and concerns during regular check-ins, healthcare organizations can identify trends that point towards broader improvements in policies, procedures, and even staff training programs aimed at enhancing empathy and effective communication skills among personnel.

In essence, prioritizing regular check-ins as part of shared planning initiatives represents a profound commitment to improving patient experiences from multiple angles—it nurtures trust between providers and patients, enhances personalization of care delivery models, fosters community within clinical settings, and drives continuous improvement across healthcare systems. As we move forward in our quest for excellence in health care provisioning, let us remember: it's not just about treating illnesses; it's about nurturing relationships built on respectful dialogue and shared understanding—a foundation upon which improved health outcomes are sure to follow.

In the rapidly evolving landscape of healthcare, the fusion of technology and patient engagement stands as a beacon of transformative potential, especially when it comes to shared planning for improved patient experiences. This synergy isn't merely about

adopting the latest gadgets or software; it's about crafting a more personalized, efficient, and empathetic healthcare journey for every individual. Let's delve into how leveraging technology can revolutionize the way we plan and deliver care, making it more aligned with patients' needs and expectations.

At the heart of this transformation lies the ability to communicate more effectively. Traditional methods often leave patients feeling like passive recipients of care rather than active participants in their health journey. With technology, we can bridge this gap through user-friendly platforms that facilitate seamless communication between patients and healthcare providers. Imagine an app where patients can easily schedule appointments, request prescription refills, or even send quick updates about their condition to their healthcare team. Such tools not only enhance convenience but also foster a sense of partnership, where both parties are on the same page regarding treatment plans and goals.

Moreover, data analytics and artificial intelligence offer profound insights that can tailor experiences to individual needs. By analyzing vast amounts of patient data—preferences, past interactions, health outcomes—healthcare providers can anticipate needs and customize care plans proactively. For instance, an AI-driven system might recognize patterns indicating a patient is at risk of non-adherence to medication schedules due to lifestyle factors and suggest interventions before issues arise. This predictive approach not only improves outcomes but also empowers patients by showing they are central to the decision-making process.

Telehealth services have emerged as a pivotal tool in enhancing engagement through technology. The pandemic has accelerated their adoption, demonstrating how virtual consultations can make healthcare more accessible, particularly for those with mobility challenges or living in remote areas. Beyond convenience, telehealth allows for continuous care monitoring and adjustments without geographical constraints, ensuring that every

patient's voice is heard and considered in their care plan—regardless of location.

Furthermore, incorporating feedback mechanisms directly within these technological platforms is crucial for fostering an environment of shared planning. Patients should feel empowered to rate their experiences, provide suggestions for improvements, or express concerns instantly. This two-way feedback loop not only enhances service quality but also builds trust—a cornerstone of effective patient-provider relationships. When patients see their input leading to tangible changes in their care experience, it reinforces the notion that they are valued partners rather than mere recipients of services.

However, while embracing technology promises immense benefits, it's essential to navigate potential pitfalls carefully—ensuring data privacy and security remain paramount while designing these systems. Patients must trust that their sensitive information is safeguarded as diligently as their health is cared for. Moreover, we must remember that technology should complement rather than replace human interaction; empathetic human touch remains irreplaceable in healing and support processes.

In conclusion, leveraging technology for enhanced engagement in shared planning represents a significant stride towards improving patient experiences in healthcare settings. By harnessing communication tools, predictive analytics, telehealth capabilities—and most importantly—by involving patients actively in their own care journeys—we pave the way for a future where healthcare is not just about treating illnesses but about nurturing well-being through collaboration and understanding. As we embrace these advancements thoughtfully

Shared planning, particularly in the healthcare sector, represents a transformative approach to enhancing patient experiences by fostering collaboration among patients, caregivers, and healthcare professionals. In this evolving landscape, technology plays a pivotal role, offering innovative tools that streamline communication and coordination,

ultimately leading to improved outcomes and heightened satisfaction for all parties involved. Let's explore how digital simulations, mobile applications, and virtual consultations are reshaping the realm of shared planning.

Digital simulations emerge as a powerful tool in preparing both patients and healthcare providers for various medical procedures or treatment plans. These simulations can provide immersive experiences that mirror real-life scenarios, allowing patients to visualize their treatment journey and understand potential outcomes. For instance, virtual reality (VR) can simulate postoperative recovery processes or the intricacies of managing chronic diseases. By engaging with these realistic simulations, patients become active participants in their care planning, which can significantly reduce anxiety and enhance their preparedness. Moreover, healthcare providers benefit from these simulations by gaining insights into patient perspectives, enabling more empathetic and tailored planning sessions.

Apps designed for health management have revolutionized how patients engage with their health plans outside of clinical settings. These applications often feature personalized dashboards where patients can track medications, appointments, vital signs, and even communicate directly with their healthcare team. Such apps empower individuals to take ownership of their health journeys by providing accessible information at their fingertips. Shared planning is further facilitated as these platforms allow for seamless integration of data from various sources—be it wearable health monitors or electronic health records—ensuring everyone involved has up-to-date insights into the patient's condition and progress. This accessibility fosters proactive discussions during consultations and encourages ongoing collaboration between visits.

Virtual consultations have emerged as a cornerstone in facilitating shared planning across geographical barriers. The COVID-19 pandemic accelerated the adoption of telehealth services, proving not just a necessity but a valuable addition to traditional care models.

Through video conferencing platforms tailored for healthcare interactions, patients can participate in consultations from the comfort of their homes. This accessibility is particularly beneficial for those with mobility issues or living in remote areas where accessing specialized care might otherwise be challenging. Virtual consultations make it easier to involve family members or caregivers in discussions without logistical hurdles, ensuring that shared decisions are made collaboratively and inclusively—a critical aspect of enhancing patient experiences.

The convergence of these technological tools—digital simulations, apps for health management, and virtual consultations—paves the way for an integrated approach to shared planning in healthcare. By leveraging these innovations effectively, healthcare systems can create environments where patient voices are central to decision-making processes. This shift not only improves individual patient experiences but also contributes to overall systemic efficiency by reducing unnecessary hospital visits and enhancing continuity of care across different settings and practitioners. As we continue to embrace these technologies with thoughtfulness and intention, we move closer to realizing a future where shared planning isn't just an ideal but a standard practice that genuinely enhances the quality of care delivered to every individual seeking health support.

Shared planning in the context of healthcare represents a transformative approach to patient care, fundamentally shifting the dynamics from a one-sided medical directive to a collaborative partnership between healthcare providers and patients. This shift is significantly amplified by the advent and integration of advanced technologies, which hold the promise of not only improving understanding but also enhancing involvement in the treatment process. The synergy between shared planning and technology can redefine patient experiences, making them more informed, engaged, and ultimately, more satisfied with their healthcare journey.

The cornerstone of this enhancement lies in **digital health tools**, which include electronic health records (EHRs), patient portals, mobile health apps, and telehealth services. These

technologies provide platforms where patients can access comprehensive information about their health conditions, treatment options, and potential outcomes in an easily digestible format. Imagine a scenario where a patient diagnosed with diabetes can log into a secure patient portal to review their latest blood glucose readings, educational materials on diet management, and interact directly with their healthcare team to adjust their treatment plan—all from the comfort of home. Such accessibility demystifies complex medical jargon and empowers patients to take an active role in managing their health.

Moreover, **telehealth** has revolutionized how patients engage with their healthcare providers. Virtual consultations break down geographical barriers and allow for more frequent communication without the logistical challenges of travel. Patients can discuss symptoms, ask questions about medications or therapies, and receive immediate feedback—fostering a sense of being heard and understood. This continuous dialogue ensures that treatment plans are not only tailored to individual needs but also adjusted swiftly in response to changing health statuses or concerns.

Another pivotal aspect is **patient education technology**. Interactive tutorials, virtual reality simulations for surgical procedures, or gamified approaches to medication adherence educate patients in ways traditional methods cannot match. By engaging multiple senses and offering interactive learning experiences, these technologies enhance comprehension and retention of critical health information. When patients understand why certain treatments are necessary or how lifestyle changes can impact their condition positively, they're more likely to adhere to prescribed regimens actively participate in shared decision-making processes.

Furthermore, **wearable technology** contributes significantly by providing real-time monitoring data that both patients and providers can access. Devices tracking heart rates, physical activity levels, or sleep patterns offer tangible insights into how daily activities affect one's health status. This data-driven approach not only personalizes treatment

plans but also encourages patients to see the direct impact of their actions on their well-being—a powerful motivator for sustained engagement in treatment processes.

In essence, technology's role in facilitating shared planning isn't merely about adding digital tools; it's about fostering an environment where patients feel valued partners rather than passive recipients of care. Through improved understanding bolstered by accessible information and increased involvement facilitated by interactive platforms and devices, both parties contribute equally toward crafting effective treatment strategies tailored uniquely to individual needs. This collaborative spirit not only enhances patient experiences but also leads to better outcomes through increased adherence to treatments, more proactive management of conditions, and ultimately—a deeper satisfaction with the care received.

As we continue advancing into an era where technology permeates every aspect of life—its integration into healthcare through shared planning presents an exciting frontier—a frontier poised to redefine what it means for patients to be truly involved in their own healing journeys, making each step forward not just medically sound but deeply personal.

Evaluating Success: Measuring Patient Experience in Shared Planning for Improved Patient Experiences

In the ever-evolving landscape of healthcare, the concept of shared planning has emerged as a beacon of hope for enhancing patient experiences. This collaborative approach, where patients, families, and healthcare providers work together to design care plans, is not merely a trend but a fundamental shift towards person-centered care. To truly gauge the success of such initiatives, it's vital to delve into the nuances of measuring patient experience effectively. This essay explores how shared planning can serve as both a tool for improvement and a metric for success in evaluating patient experiences.

At its core, shared planning is about listening—truly hearing what patients have to say about their health journeys and involving them in decisions that directly affect their well-being. When patients feel heard and valued, it naturally leads to higher satisfaction levels. However, satisfaction alone doesn't paint a complete picture; it's essential to look deeper into various facets that contribute to an overall positive experience. Herein lies the challenge—and opportunity—of measurement.

One effective approach is through comprehensive feedback mechanisms that capture both quantitative and qualitative data. Surveys are a common method, offering structured insights into patient perceptions regarding communication, empathy, and the clarity of information provided by healthcare teams. Yet, these should be complemented with open-ended questions that allow patients to share their stories—the unexpected moments of kindness or frustration that significantly impact their experiences but might not surface in numerical ratings.

Moreover, incorporating direct observations from healthcare providers can offer another layer of understanding. Observing interactions during shared planning sessions can reveal much about the dynamics at play—how inclusively patients are engaged, how their insights are integrated into care plans, and the overall atmosphere of respect and collaboration fostered during these meetings. These observations can highlight areas where processes might need refinement for even better outcomes.

Another crucial aspect is tracking long-term outcomes tied to patient experience metrics derived from shared planning initiatives. For instance, monitoring readmission rates post-discharge can provide insight into how well care plans align with patient needs outside the clinical setting—a tangible measure of effectiveness beyond immediate hospital stays. Similarly, looking at adherence to treatment plans initiated through shared decision-making illuminates whether patients feel empowered enough to manage their health proactively post-discharge.

It's also important to recognize that measuring success isn't just about identifying areas needing improvement—it's equally about celebrating achievements and sharing learnings across different healthcare settings. By establishing benchmarks and best practices derived from successful shared planning cases, healthcare institutions can inspire one another toward continuous enhancement of patient experiences everywhere.

In conclusion, evaluating success in measuring patient experience within the framework of shared planning requires an holistic approach—one that values both qualitative narratives and quantitative indicators while remaining vigilant on long-term outcomes tied to those initial interactions between patients and their care teams. As we strive toward more person-centered care models, embracing this multifaceted evaluation strategy ensures we're not just meeting standards but exceeding expectations in delivering truly meaningful patient experiences that resonate long after clinical visits end.

Shared planning, a collaborative approach involving patients, healthcare providers, and sometimes families or care teams in the decision-making process regarding treatment and care plans, has emerged as a pivotal strategy to enhance patient experiences. The essence of this approach lies in recognizing patients as active participants rather than passive recipients of care. To effectively assess the impact of shared planning on improving patient experiences, it's vital to adopt a multifaceted approach that encompasses both qualitative and quantitative metrics, ensuring a comprehensive understanding of its benefits and areas for improvement.

Quantitative Metrics:

1. **Patient Satisfaction Scores:** Implementing standardized surveys post-treatment or during follow-ups can provide concrete data on how shared planning influences

patient satisfaction. Questions should specifically target aspects like involvement in decision-making, understanding of the treatment plan, and overall experience with the care process.

2. **Readmission Rates:** A decrease in readmission rates can indicate that patients are better informed and more engaged in their care processes due to shared planning. This metric reflects not just improved patient experiences but also enhanced health outcomes.

3. **Adherence to Treatment Plans:** Tracking how well patients adhere to prescribed treatments can serve as an indirect measure of their engagement level during the planning phase. Greater adherence often correlates with patients feeling more invested in their care plans due to having a say in their development.

Qualitative Metrics:

1. **Patient Narratives:** Collecting stories or testimonials from patients about their experiences with shared planning allows for a deeper insight into how this approach affects their emotional and psychological well-being alongside physical health outcomes. These narratives can highlight aspects like empowerment, trust in

caregivers, and perceived control over health decisions that quantitative data might overlook.

2. **Focus Groups and Interviews:** Conducting discussions with patients who have experienced shared planning can yield rich insights into what works well and what could be improved within the framework. These sessions can explore nuanced feelings about collaboration, communication quality, and overall satisfaction with the care received.

3. **Healthcare Provider Feedback:** Gathering perspectives from healthcare professionals involved in the shared planning process provides an external viewpoint on its effectiveness. Providers can share observations on changes in patient engagement levels, communication dynamics, and any challenges encountered during implementation, offering a balanced view of shared planning's impact on patient experiences from both sides of the consultation room.

Methods for Assessment:

To effectively utilize these metrics and methods for assessing the effectiveness of shared planning on patient experiences:

- **Longitudinal Studies:** Implement ongoing assessments over extended periods to capture long-term impacts on patient satisfaction and health outcomes.
- **Control Groups:** Where feasible, compare outcomes between groups experiencing traditional care versus those involved in shared planning to isolate variables affecting patient experience improvements.
- **Continuous Feedback Loops:** Establish systems for regular feedback collection post-implementation phases to allow for iterative improvements based on real-time insights from both patients and providers.
- **Multidisciplinary Analysis Teams:** Form teams composed of healthcare providers, patient advocates, and researchers to analyze data comprehensively from various angles—medical efficacy, emotional impact, logistical feasibility—ensuring all facets are considered when evaluating success or areas needing attention.

In conclusion, assessing the effectiveness of shared planning for improved patient experiences requires a blend of measurable outcomes—like satisfaction scores and

readmission rates—and qualitative insights drawn from personal narratives and provider observations. By employing these diverse metrics through structured

Shared planning for improved patient experiences is a cornerstone of modern healthcare, reflecting a profound shift towards patient-centered care. This approach isn't merely about treating illnesses; it's about crafting a journey where patients feel valued, heard, and actively involved in their care processes. At the heart of this transformation lies the principle of continuous improvement, fueled by learning from patient feedback and outcomes. This iterative process not only enhances the quality of care but also fosters a culture of respect and collaboration between healthcare providers and patients.

The essence of shared planning revolves around building partnerships with patients from the outset. It involves engaging them in discussions about their health goals, preferences, and concerns, ensuring that every plan is tailored to their unique needs and circumstances. This collaborative approach begins with open communication; healthcare teams must create an environment where patients feel comfortable sharing their thoughts and feelings without fear of judgment. Feedback mechanisms—whether through direct conversations, surveys, or digital platforms—become vital tools in this process, allowing patients to articulate their experiences and suggestions for improvement continually.

Learning from patient feedback is not just about acknowledging what's working well but critically examining areas that fall short. Outcomes, both positive and negative, are invaluable data points that guide refinement efforts. When patients share their experiences—be it the warmth of a nurse's reassurance or the frustration of lengthy wait times—these insights illuminate paths for enhancement. Healthcare providers must cultivate a mindset geared towards learning; every piece of feedback is an opportunity to understand better how services can be fine-tuned to meet patient expectations more effectively.

This continuous improvement loop is dynamic and multifaceted. It requires not only collecting feedback but also analyzing it thoughtfully to identify patterns and underlying issues. Teams must then brainstorm innovative solutions collaboratively with patients, ensuring that changes are meaningful and impactful. For instance, if multiple patients mention difficulty navigating a clinic's layout, perhaps a redesign project could involve current patients in the planning stages to ensure the new layout truly meets their needs.

Furthermore, sharing these improvements back with patients demonstrates commitment to their voices shaping their care experience—a powerful affirmation of their importance in the healthcare ecosystem. Transparency in implementing changes fosters trust and encourages ongoing engagement from patients who see tangible results stemming from their input.

In conclusion, shared planning for improved patient experiences hinges on embracing continuous improvement as a core value—one deeply rooted in listening to patient feedback and measuring outcomes against those very insights. By creating a culture where every voice matters and every experience counts, healthcare providers can refine future approaches continuously. This not only elevates the standard of care but also builds lasting relationships based on mutual respect and collaboration—a true testament to putting the 'patient' at the heart of healthcare delivery.

Shared planning in pediatric orthodontic care represents a transformative approach that significantly enhances patient experiences by fostering a collaborative environment among all stakeholders involved in a child's dental journey. This methodology isn't merely about coordinating appointments and treatments; it's about creating a comprehensive, patient-centered framework where every voice—from the child to the parents, and from the orthodontist to the dental hygienist—contributes to crafting an individualized plan that prioritizes not just clinical outcomes, but also emotional and psychological wellbeing.

The essence of shared planning lies in its ability to bridge gaps between different perspectives, merging professional expertise with personal insights. When orthodontists engage directly with parents and children, they gain invaluable insights into the family's expectations, concerns, and daily routines. This understanding allows for treatments that are not only clinically sound but also seamlessly integrated into the child's life, reducing anxiety and increasing compliance. For instance, discussing treatment timelines during a family meeting can help align expectations and prepare everyone for what lies ahead, transforming potential sources of stress into manageable milestones.

Moreover, shared planning encourages an interdisciplinary approach, involving various specialists such as pediatric dentists, oral surgeons, and even psychologists when necessary. This teamwork ensures that every aspect of a child's oral health is considered holistically. It opens doors for innovative solutions tailored specifically to each child's unique needs—be it adapting treatment plans for children with special healthcare needs or incorporating cutting-edge technology to make procedures less intimidating.

Furthermore, this collaborative framework empowers patients by making them active participants in their care journey rather than passive recipients of treatment. When children understand their condition and the reasons behind certain procedures, they're more likely to feel confident and cooperative during treatments. Engaging them in discussions about their preferences for appliances or colors on braces can turn an otherwise daunting experience into something they look forward to—a rite of passage marked by small victories and personalized touches that affirm their individuality.


Parents play a pivotal role in this process too. Their involvement ensures continuity of care at home and reinforces the importance of oral hygiene practices post-treatment. Shared planning sessions provide them with clear guidance on maintaining oral health between visits and recognizing signs that might necessitate immediate attention—a partnership that extends beyond the clinic walls into everyday life.

In essence, shared planning redefines pediatric orthodontic care by placing empathy and collaboration at its core. It acknowledges that every smile is unique—not just in terms of dental structure but also in terms of emotional significance. By weaving together the threads of professional insight with personal narratives, we create a tapestry of care that respects individual differences while aiming for universally positive outcomes: healthy smiles backed by confidence and joyful experiences throughout the orthodontic journey.

About dentistry

- Sub-Millimeter Surgical Dexterity
- Knowledge of human health, disease, pathology, and anatomy
- Communication/Interpersonal Skills
- Analytical Skills
- Critical Thinking
- Empathy/Professionalism
- Private practices
- Primary care clinics
- Hospitals
- Physician
- dental assistant
- dental technician
- dental hygienist
- various dental specialists

Dentistry



A dentist treats a patient with the help of a dental assistant.

Occupation

- Dentist
- Dental Surgeon
- Doctor

Names

[¹][^{nb 1}]

Occupation type

Profession

Activity sectors

Health care, Anatomy, Physiology, Pathology, Medicine, Pharmacology, Surgery

Description

Competencies

Education required

Dental Degree


Fields of employment

Related jobs

ICD-9-CM 23-24

MeSH D003813

[[edit on Wikidata](#)]



An oral surgeon and dental assistant removing a wisdom tooth

Dentistry, also known as **dental medicine** and **oral medicine**, is the branch of medicine focused on the teeth, gums, and mouth. It consists of the study, diagnosis, prevention, management, and treatment of diseases, disorders, and conditions of the mouth, most commonly focused on dentition (the development and arrangement of teeth) as well as the oral mucosa.^[2] Dentistry may also encompass other aspects of the craniofacial complex including the temporomandibular joint. The practitioner is called a dentist.

The history of dentistry is almost as ancient as the history of humanity and civilization, with the earliest evidence dating from 7000 BC to 5500 BC.^[3] Dentistry is thought to have been the first specialization in medicine which has gone on to develop its own accredited degree with its own specializations.^[4] Dentistry is often also understood to subsume the now largely defunct medical specialty of stomatology (the study of the mouth and its disorders and diseases) for which reason the two terms are used interchangeably in certain regions.

However, some specialties such as oral and maxillofacial surgery (facial reconstruction) may require both medical and dental degrees to accomplish. In European history, dentistry is considered to have stemmed from the trade of barber surgeons.^[5]

Dental treatments are carried out by a dental team, which often consists of a dentist and dental auxiliaries (such as dental assistants, dental hygienists, dental technicians, and dental therapists). Most dentists either work in private practices (primary care), dental hospitals, or (secondary care) institutions (prisons, armed forces bases, etc.).

The modern movement of evidence-based dentistry calls for the use of high-quality scientific research and evidence to guide decision-making such as in manual tooth conservation, use of fluoride water treatment and fluoride toothpaste, dealing with oral diseases such as tooth decay and periodontitis, as well as systematic diseases such as osteoporosis, diabetes, celiac disease, cancer, and HIV/AIDS which could also affect the oral cavity. Other practices relevant to evidence-based dentistry include radiology of the mouth to inspect teeth deformity or oral malaises, haematology (study of blood) to avoid bleeding complications during dental surgery, cardiology (due to various severe complications arising from dental surgery with patients with heart disease), etc.

Terminology

[edit]

The term dentistry comes from *dentist*, which comes from French *dentiste*, which comes from the French and Latin words for tooth.^[6] The term for the associated scientific study of teeth is **odontology** (from Ancient Greek:

ὀδοντολογία, *odon-* (from *odoús*, lit. 'tooth') – the study of the structure, development, and abnormalities of the teeth.


Dental treatment

[edit]

Dentistry usually encompasses practices related to the oral cavity.^[7] According to the World Health Organization, oral diseases are major public health problems due to their high incidence and prevalence across the globe, with the disadvantaged affected more than other socio-economic groups.^[8]

The majority of dental treatments are carried out to prevent or treat the two most common oral diseases which are dental caries (tooth decay) and periodontal disease (gum disease or pyorrhea). Common treatments involve the restoration of teeth, extraction or surgical removal of teeth, scaling and root planing, endodontic root canal treatment, and cosmetic dentistry^[9]

By nature of their general training, dentists, without specialization can carry out the majority of dental treatments such as restorative (fillings, crowns, bridges), prosthetic (dentures), endodontic (root canal) therapy, periodontal (gum) therapy, and extraction of teeth, as well as performing examinations, radiographs (x-rays), and diagnosis. Dentists can also prescribe medications used in the field such as antibiotics, sedatives, and any other drugs used in patient management. Depending on their licensing boards, general dentists may be required to complete additional training to perform sedation, dental implants, etc.



Irreversible enamel defects caused by an untreated celiac disease. They may be the only clue to its diagnosis, even in absence of gastrointestinal symptoms, but are often confused with fluorosis, tetracycline


discoloration, acid reflux or other causes.^{[10][11][12]} The National Institutes of Health include a dental exam in the diagnostic protocol of celiac disease.^[10]

Dentists also encourage the prevention of oral diseases through proper hygiene and regular, twice or more yearly, checkups for professional cleaning and evaluation. Oral infections and inflammations may affect overall health and conditions in the oral cavity may be indicative of systemic diseases, such as osteoporosis, diabetes, celiac disease or cancer.^{[7][10][13][14]} Many studies have also shown that gum disease is associated with an increased risk of diabetes, heart disease, and preterm birth. The concept that oral health can affect systemic health and disease is referred to as "oral-systemic health".


Education and licensing

[edit]

Main article: Dentistry throughout the world



A sagittal cross-section of a molar tooth; 1: crown, 2: root, 3: enamel, 4: dentin and dentin tubules, 5: pulp chamber, 6: blood vessels and nerve, 7: periodontal ligament, 8: apex and periapical region, 9: alveolar bone



Early dental chair in Pioneer West Museum in Shamrock, Texas

John M. Harris started the world's first dental school in Bainbridge, Ohio, and helped to establish dentistry as a health profession. It opened on 21 February 1828, and today is a dental museum.^[15] The first dental college, Baltimore College of Dental Surgery, opened in Baltimore, Maryland, US in 1840. The second in the United States was the Ohio College of Dental Surgery, established in Cincinnati, Ohio, in 1845.^[16] The Philadelphia College of Dental Surgery followed in 1852.^[17] In 1907, Temple University accepted a bid to incorporate the school.

Studies show that dentists that graduated from different countries,^[18] or even from different dental schools in one country,^[19] may make different clinical decisions for the same clinical condition. For example, dentists that graduated from Israeli dental schools may recommend the removal of asymptomatic impacted third molar (wisdom teeth) more often than dentists that graduated from Latin American or Eastern European dental schools.^[20]

In the United Kingdom, the first dental schools, the London School of Dental Surgery and the Metropolitan School of Dental Science, both in London, opened in 1859.^[21] The British Dentists Act of 1878 and the 1879 Dentists Register limited the title of "dentist" and "dental surgeon" to qualified and registered practitioners.^{[22][23]} However, others could legally describe themselves as "dental experts" or "dental consultants".^[24] The practice of dentistry in the United Kingdom became fully regulated with the 1921 Dentists Act, which required the registration of anyone practising dentistry.^[25] The British Dental Association, formed in 1880 with Sir

John Tomes as president, played a major role in prosecuting dentists practising illegally.^[22] Dentists in the United Kingdom are now regulated by the General Dental Council.


In many countries, dentists usually complete between five and eight years of post-secondary education before practising. Though not mandatory, many dentists choose to complete an internship or residency focusing on specific aspects of dental care after they have received their dental degree. In a few countries, to become a qualified dentist one must usually complete at least four years of postgraduate study;^[26] Dental degrees awarded around the world include the Doctor of Dental Surgery (DDS) and Doctor of Dental Medicine (DMD) in North America (US and Canada), and the Bachelor of Dental Surgery/Baccalaureus Dentalis Chirurgiae (BDS, BDent, BChD, BDS_{Sc}) in the UK and current and former British Commonwealth countries.

All dentists in the United States undergo at least three years of undergraduate studies, but nearly all complete a bachelor's degree. This schooling is followed by four years of dental school to qualify as a "Doctor of Dental Surgery" (DDS) or "Doctor of Dental Medicine" (DMD). Specialization in dentistry is available in the fields of Anesthesiology, Dental Public Health, Endodontics, Oral Radiology, Oral and Maxillofacial Surgery, Oral Medicine, Orofacial Pain, Pathology, Orthodontics, Pediatric Dentistry (Pedodontics), Periodontics, and Prosthodontics.^[27]

Specialties

[edit]

Main article: Specialty (dentistry)



A modern dental clinic in Lappeenranta, Finland

Some dentists undertake further training after their initial degree in order to specialize. Exactly which subjects are recognized by dental registration bodies varies according to location. Examples include:

- Anesthesiology^[28] – The specialty of dentistry that deals with the advanced use of general anesthesia, sedation and pain management to facilitate dental procedures.
- Cosmetic dentistry – Focuses on improving the appearance of the mouth, teeth and smile.
- Dental public health – The study of epidemiology and social health policies relevant to oral health.
- Endodontics (also called *endodontology*) – Root canal therapy and study of diseases of the dental pulp and periapical tissues.
- Forensic odontology – The gathering and use of dental evidence in law. This may be performed by any dentist with experience or training in this field. The function of the forensic dentist is primarily documentation and verification of identity.
- Geriatric dentistry or *geriodontics* – The delivery of dental care to older adults involving the diagnosis, prevention, and treatment of problems associated with normal aging and age-related diseases as part of an interdisciplinary team with other health care professionals.
- Oral and maxillofacial pathology – The study, diagnosis, and sometimes the treatment of oral and maxillofacial related diseases.
- Oral and maxillofacial radiology – The study and radiologic interpretation of oral and maxillofacial diseases.


- Oral and maxillofacial surgery (also called *oral surgery*) – Extractions, implants, and surgery of the jaws, mouth and face.^[nb 2]
- Oral biology – Research in dental and craniofacial biology
- Oral Implantology – The art and science of replacing extracted teeth with dental implants.
- Oral medicine – The clinical evaluation and diagnosis of oral mucosal diseases
- Orthodontics and dentofacial orthopedics – The straightening of teeth and modification of midface and mandibular growth.
- Pediatric dentistry (also called *pedodontics*) – Dentistry for children
- Periodontology (also called *periodontics*) – The study and treatment of diseases of the periodontium (non-surgical and surgical) as well as placement and maintenance of dental implants
- Prosthodontics (also called *prosthetic dentistry*) – Dentures, bridges and the restoration of implants.
 - Some prosthodontists super-specialize in maxillofacial prosthetics, which is the discipline originally concerned with the rehabilitation of patients with congenital facial and oral defects such as cleft lip and palate or patients born with an underdeveloped ear (microtia). Today, most maxillofacial prosthodontists return function and esthetics to patients with acquired defects secondary to surgical removal of head and neck tumors, or secondary to trauma from war or motor vehicle accidents.
- Special needs dentistry (also called *special care dentistry*) – Dentistry for those with developmental and acquired disabilities.
- Sports dentistry – the branch of sports medicine dealing with prevention and treatment of dental injuries and oral diseases associated with sports and exercise.^[29] The sports dentist works as an individual consultant or as a member of the Sports Medicine Team.

- Veterinary dentistry – The field of dentistry applied to the care of animals. It is a specialty of veterinary medicine.^[30]^[31]


History

[edit]

See also: History of dental treatments



A wealthy patient falling over because of having a tooth extracted with such vigour by a fashionable dentist, c. 1790. History of Dentistry.



Farmer at the dentist, Johann Liss, c. 1616–17

Tooth decay was low in pre-agricultural societies, but the advent of farming society about 10,000 years ago correlated with an increase in tooth decay (cavities).^[32] An infected tooth from Italy partially cleaned with flint tools, between 13,820 and 14,160 years old, represents the oldest known dentistry,^[33] although a 2017 study suggests that 130,000 years ago the Neanderthals already used rudimentary dentistry tools.^[34] In Italy evidence dated to the Paleolithic, around 13,000 years ago, points to bitumen used to fill a tooth^[35] and in Neolithic Slovenia, 6500 years ago, beeswax was used to close a fracture in a tooth.^[36] The Indus valley has yielded evidence of dentistry being practised as far back as 7000 BC, during the Stone Age.^[37] The Neolithic site of Mehrgarh (now in Pakistan's south western province of Balochistan) indicates that this form of dentistry involved curing tooth related disorders with bow drills operated, perhaps, by skilled bead-crafters.^[3] The reconstruction of this ancient form of dentistry showed that the methods used were reliable and effective.^[38] The earliest dental filling, made of beeswax, was discovered in Slovenia and dates from 6500 years ago.^[39] Dentistry was practised in prehistoric Malta, as evidenced by a skull which had a dental abscess lanced from the root of a tooth dating back to around 2500 BC.^[40]

An ancient Sumerian text describes a "tooth worm" as the cause of dental caries.^[41] Evidence of this belief has also been found in ancient India, Egypt, Japan, and China. The legend of the worm is also found in the *Homeric Hymns*,^[42] and as late as the 14th century AD the surgeon Guy de Chauliac still promoted the belief that worms cause tooth decay.^[43]

Recipes for the treatment of toothache, infections and loose teeth are spread throughout the Ebers Papyrus, Kahun Papyri, Brugsch Papyrus, and Hearst papyrus of Ancient Egypt.^[44] The Edwin Smith Papyrus, written in the 17th century BC but which may reflect previous manuscripts from as early as 3000 BC, discusses the treatment of dislocated or fractured jaws.^{[44][45]} In the 18th century BC, the Code of Hammurabi referenced dental extraction twice as it related to punishment.^[46]


Examination of the remains of some ancient Egyptians and Greco-Romans reveals early attempts at dental prosthetics.^[47] However, it is possible the prosthetics were prepared after death for aesthetic reasons.^[44]

Ancient Greek scholars Hippocrates and Aristotle wrote about dentistry, including the eruption pattern of teeth, treating decayed teeth and gum disease, extracting teeth with forceps, and using wires to stabilize loose teeth and fractured jaws.^[48] Use of dental appliances, bridges and dentures was applied by the Etruscans in northern Italy, from as early as 700 BC, of human or other animal teeth fastened together with gold bands.^{[49][50][51]} The Romans had likely borrowed this technique by the 5th century BC.^{[50][52]} The Phoenicians crafted dentures during the 6th–4th century BC, fashioning them from gold wire and incorporating two ivory teeth.^[53] In ancient Egypt, Hesy-Ra is the first named "dentist" (greatest of the teeth). The Egyptians bound replacement teeth together with gold wire. Roman medical writer Cornelius Celsus wrote extensively of oral diseases as well as dental treatments such as narcotic-containing emollients and astringents.^[54] The earliest dental amalgams were first documented in a Tang dynasty medical text written by the Chinese physician Su Kung in 659, and appeared in Germany in 1528.^{[55][56]}

During the Islamic Golden Age Dentistry was discussed in several famous books of medicine such as The Canon in medicine written by Avicenna and Al-Tasreef by Al-Zahrawi who is considered the greatest surgeon of the Middle Ages,^[57] Avicenna said that jaw fracture should be reduced according to the occlusal guidance of the teeth; this principle is still valid in modern times. Al-Zahrawi invented over 200 surgical tools that resemble the modern kind.^[58]

Historically, dental extractions have been used to treat a variety of illnesses. During the Middle Ages and throughout the 19th century, dentistry was not a profession in itself, and often dental procedures were performed by barbers or general physicians. Barbers usually limited their practice to extracting teeth which

alleviated pain and associated chronic tooth infection. Instruments used for dental extractions date back several centuries. In the 14th century, Guy de Chauliac most probably invented the dental pelican^[59] (resembling a pelican's beak) which was used to perform dental extractions up until the late 18th century. The pelican was replaced by the dental key^[60] which, in turn, was replaced by modern forceps in the 19th century.^[61]




Dental needle-nose pliers designed by Fauchard in the late 17th century to use in prosthodontics

The first book focused solely on dentistry was the "Artzney Buchlein" in 1530,^[48] and the first dental textbook written in English was called "Operator for the Teeth" by Charles Allen in 1685.^[23]

In the United Kingdom, there was no formal qualification for the providers of dental treatment until 1859 and it was only in 1921 that the practice of dentistry was limited to those who were professionally qualified. The Royal Commission on the National Health Service in 1979 reported that there were then more than twice as many registered dentists per 10,000 population in the UK than there were in 1921.^[62]

Modern dentistry

[edit]




A microscopic device used in dental analysis, c. 1907

It was between 1650 and 1800 that the science of modern dentistry developed. The English physician Thomas Browne in his *A Letter to a Friend* (c. 1656 pub. 1690) made an early dental observation with characteristic humour:

The Egyptian Mummies that I have seen, have had their Mouths open, and somewhat gaping, which affordeth a good opportunity to view and observe their Teeth, wherein 'tis not easie to find any wanting or decayed: and therefore in Egypt, where one Man practised but one Operation, or the Diseases but of single Parts, it must needs be a barren Profession to confine unto that of drawing of Teeth, and little better than to have been Tooth-drawer unto King Pyrrhus, who had but two in his Head.


The French surgeon Pierre Fauchard became known as the "father of modern dentistry". Despite the limitations of the primitive surgical instruments during the late 17th and early 18th century, Fauchard was a highly skilled surgeon who made remarkable improvisations of dental instruments, often adapting tools from

watchmakers, jewelers and even barbers, that he thought could be used in dentistry. He introduced dental fillings as treatment for dental cavities. He asserted that sugar-derived acids like tartaric acid were responsible for dental decay, and also suggested that tumors surrounding the teeth and in the gums could appear in the later stages of tooth decay.^{[63][64]}



Panoramic radiograph of historic dental implants, made 1978

Fauchard was the pioneer of dental prosthesis, and he invented many methods to replace lost teeth. He suggested that substitutes could be made from carved blocks of ivory or bone. He also introduced dental braces, although they were initially made of gold, he discovered that the teeth position could be corrected as the teeth would follow the pattern of the wires. Waxed linen or silk threads were usually employed to fasten the braces. His contributions to the world of dental science consist primarily of his 1728 publication *Le chirurgien dentiste* or *The Surgeon Dentist*. The French text included "basic oral anatomy and function, dental construction, and various operative and restorative techniques, and effectively separated dentistry from the wider category of surgery".^{[63][64]}



A modern dentist's chair

After Fauchard, the study of dentistry rapidly expanded. Two important books, *Natural History of Human Teeth* (1771) and *Practical Treatise on the Diseases of the Teeth* (1778), were published by British surgeon John Hunter. In 1763, he entered into a period of collaboration with the London-based dentist James Spence. He began to theorise about the possibility of tooth transplants from one person to another. He realised that the chances of a successful tooth transplant (initially, at least) would be improved if the donor tooth was as fresh as possible and was matched for size with the recipient. These principles are still used in the transplantation of internal organs. Hunter conducted a series of pioneering operations, in which he attempted a tooth transplant. Although the donated teeth never properly bonded with the recipients' gums, one of Hunter's patients stated that he had three which lasted for six years, a remarkable achievement for the period.^[65]

Major advances in science were made in the 19th century, and dentistry evolved from a trade to a profession. The profession came under government regulation by the end of the 19th century. In the UK, the Dentist Act was passed in 1878 and the British Dental Association formed in 1879. In the same year, Francis Brodie Imlach was the first ever dentist to be elected President of the Royal College of Surgeons (Edinburgh), raising dentistry onto a par with clinical surgery for the first time.^[66]

Hazards in modern dentistry

[edit]

Main article: Occupational hazards in dentistry

Long term occupational noise exposure can contribute to permanent hearing loss, which is referred to as noise-induced hearing loss (NIHL) and tinnitus. Noise exposure can cause excessive stimulation of the hearing mechanism, which damages the delicate structures of the inner ear.^[67] NIHL can occur when an

individual is exposed to sound levels above 90 dBA according to the Occupational Safety and Health Administration (OSHA). Regulations state that the permissible noise exposure levels for individuals is 90 dBA.^[68] For the National Institute for Occupational Safety and Health (NIOSH), exposure limits are set to 85 dBA. Exposures below 85 dBA are not considered to be hazardous. Time limits are placed on how long an individual can stay in an environment above 85 dBA before it causes hearing loss. OSHA places that limitation at 8 hours for 85 dBA. The exposure time becomes shorter as the dBA level increases.

Within the field of dentistry, a variety of cleaning tools are used including piezoelectric and sonic scalers, and ultrasonic scalers and cleaners.^[69] While a majority of the tools do not exceed 75 dBA,^[70] prolonged exposure over many years can lead to hearing loss or complaints of tinnitus.^[71] Few dentists have reported using personal hearing protective devices,^[72]^[73] which could offset any potential hearing loss or tinnitus.


Evidence-based dentistry

[edit]

Main article: Evidence-based dentistry

There is a movement in modern dentistry to place a greater emphasis on high-quality scientific evidence in decision-making. Evidence-based dentistry (EBD) uses current scientific evidence to guide decisions. It is an approach to oral health that requires the application and examination of relevant scientific data related to the patient's oral and medical health. Along with the dentist's professional skill and expertise, EBD allows dentists to stay up to date on the latest procedures and patients to receive improved treatment. A new paradigm for medical education designed to incorporate current research into education and practice was developed to help practitioners provide the best care for their patients.^[74] It was first introduced by Gordon Guyatt and the Evidence-Based Medicine Working

Group at McMaster University in Ontario, Canada in the 1990s. It is part of the larger movement toward evidence-based medicine and other evidence-based practices, especially since a major part of dentistry involves dealing with oral and systemic diseases. Other issues relevant to the dental field in terms of evidence-based research and evidence-based practice include population oral health, dental clinical practice, tooth morphology etc.



A dental chair at the University of Michigan School of Dentistry


Ethical and medicolegal issues

[edit]

Dentistry is unique in that it requires dental students to have competence-based clinical skills that can only be acquired through supervised specialized laboratory training and direct patient care.^[75] This necessitates the need for a scientific and professional basis of care with a foundation of extensive research-based education.^[76] According to some experts, the accreditation of dental schools can enhance the quality and professionalism of dental education.^{[77][78]}

See also

[edit]

-  [Medicine portal](#)
- Dental aerosol
- Dental instrument
- Dental public health

- Domestic healthcare:
 - Dentistry in ancient Rome
 - Dentistry in Canada
 - Dentistry in the Philippines
 - Dentistry in Israel
 - Dentistry in the United Kingdom
 - Dentistry in the United States
- Eco-friendly dentistry
- Geriatric dentistry
- List of dental organizations
- Pediatric dentistry
- Sustainable dentistry
- Veterinary dentistry

Notes

[edit]

1. ^ Whether Dentists are referred to as "Doctor" is subject to geographic variation. For example, they are called "Doctor" in the US. In the UK, dentists have traditionally been referred to as "Mister" as they identified themselves with barber surgeons more than physicians (as do surgeons in the UK, see Surgeon#Titles). However more UK dentists now refer to themselves as "Doctor", although this was considered to be potentially misleading by the British public in a single report (see Costley and Fawcett 2010).
2. ^ The scope of oral and maxillofacial surgery is variable. In some countries, both a medical and dental degree is required for training, and the scope includes head and neck oncology and craniofacial deformity.

References

[edit]

1. ^ Neil Costley; Jo Fawcett (November 2010). *General Dental Council Patient and Public Attitudes to Standards for Dental Professionals, Ethical Guidance and Use of the Term Doctor (PDF) (Report)*. General Dental Council/George Street Research. Archived from the original (PDF) on 4 March 2016. Retrieved 11 January 2017.
2. ^ "Glossary of Dental Clinical and Administrative Terms". American Dental Association. Archived from the original on 6 March 2016. Retrieved 1 February 2014.
3. ^ **a b** "Stone age man used dentist drill". BBC News. 6 April 2006. Retrieved 24 May 2010.
4. ^ Suddick, RP; Harris, NO (1990). "Historical perspectives of oral biology: a series". *Critical Reviews in Oral Biology and Medicine*. **1** (2): 135–51. doi:10.1177/10454411900010020301. PMID 2129621.
5. ^ "When barbers were surgeons and surgeons were barbers". Radio National. 15 April 2015. Retrieved 10 September 2021.
6. ^ "dentistry". Etymonline.com. Retrieved 17 May 2018.
7. ^ **a b** Gambhir RS (2015). "Primary care in dentistry – an untapped potential". *Journal of Family Medicine and Primary Care (Review)*. **4** (1): 13–18. doi:10.4103/2249-4863.152239. PMC 4366984. PMID 25810982.
8. ^ "What is the burden of oral disease?". WHO. Archived from the original on 30 June 2004. Retrieved 6 June 2017.
9. ^ "American Academy of Cosmetic Dentistry | Dental CE Courses". aacd.com. Retrieved 21 October 2019.
10. ^ **a b c** "Diagnosis of Celiac Disease". National Institute of Health (NIH). Archived from the original on 15 May 2017. Retrieved 6 June 2017.cite web: CS1 maint: bot: original URL status unknown (link)
11. ^ *Dental Enamel Defects and Celiac Disease (PDF) (Report)*. National Institute of Health (NIH). Archived from the original (PDF) on 5 March 2016.
12. ^ Pastore L, Carroccio A, Compilato D, Panzarella V, Serpico R, Lo Muzio L (2008). "Oral manifestations of celiac disease". *J Clin Gastroenterol (Review)*. **42** (3): 224–32. doi:10.1097/MCG.0b013e318074dd98. hdl:10447/1671. PMID 18223505. S2CID 205776755.
13. ^ Estrella MR, Boynton JR (2010). "General dentistry's role in the care for children with special needs: a review". *Gen Dent (Review)*. **58** (3): 222–29. PMID 20478802.

14. ^ da Fonseca MA (2010). "Dental and oral care for chronically ill children and adolescents". *Gen Dent (Review)*. **58** (3): 204–09, quiz 210–11. PMID 20478800.
15. ^ Owen, Lorrie K., ed. (1999). *Dictionary of Ohio Historic Places*. Vol. 2. St. Clair Shores: Somerset. pp. 1217–1218.
16. ^ Mary, Otto (2017). *Teeth: the story of beauty, inequality, and the struggle for oral health in America*. New York: The New Press. p. 70. ISBN 978-1-62097-144-4. OCLC 958458166.
17. ^ "History". Pennsylvania School of Dental Medicine. Retrieved 13 January 2016.
18. ^ Zadik Yehuda; Levin Liran (January 2008). "Clinical decision making in restorative dentistry, endodontics, and antibiotic prescription". *J Dent Educ*. **72** (1): 81–86. doi:10.1002/j.0022-0337.2008.72.1.tb04456.x. PMID 18172239.
19. ^ Zadik Yehuda; Levin Liran (April 2006). "Decision making of Hebrew University and Tel Aviv University Dental Schools graduates in every day dentistry—is there a difference?". *J Isr Dent Assoc*. **23** (2): 19–23. PMID 16886872.
20. ^ Zadik Yehuda; Levin Liran (April 2007). "Decision making of Israeli, East European, and South American dental school graduates in third molar surgery: is there a difference?". *J Oral Maxillofac Surg*. **65** (4): 658–62. doi:10.1016/j.joms.2006.09.002. PMID 17368360.
21. ^ Gelbier, Stanley (1 October 2005). "Dentistry and the University of London". *Medical History*. **49** (4): 445–462. doi:10.1017/s0025727300009157. PMC 1251639. PMID 16562330.
22. ^ **a b** Gelbier, S. (2005). "125 years of developments in dentistry, 1880–2005 Part 2: Law and the dental profession". *British Dental Journal*. **199** (7): 470–473. doi:10.1038/sj.bdj.4812875. ISSN 1476-5373. PMID 16215593. The 1879 register is referred to as the "Dental Register".
23. ^ **a b** "The story of dentistry: Dental History Timeline". British Dental Association. Archived from the original on 9 March 2012. Retrieved 2 March 2010.
24. ^ J Menzies Campbell (8 February 1955). "Banning Clerks, Colliers and other Charlatans". *The Glasgow Herald*. p. 3. Retrieved 5 April 2017.
25. ^ "History of Dental Surgery in Edinburgh" (PDF). Royal College of Surgeons of Edinburgh. Retrieved 11 December 2007.
26. ^ "Dentistry (D.D.S. or D.M.D.)" (PDF). Purdue.edu. Archived from the original (PDF) on 9 January 2017. Retrieved 17 May 2018.
27. ^ "Canadian Dental Association". cda-adc.ca. Retrieved 21 October 2019.

28. ^ "Anesthesiology recognized as a dental specialty". www.ada.org. Archived from the original on 21 September 2019. Retrieved 12 March 2019.
29. ^ "Sports dentistry". FDI World Dental Federation. Archived from the original on 23 October 2020. Retrieved 13 July 2020.
30. ^ "AVDC Home". Avdc.org. 29 November 2009. Retrieved 18 April 2010.
31. ^ "EVDC web site". Evdc.info. Archived from the original on 5 September 2018. Retrieved 18 April 2010.
32. ^ Barras, Colin (29 February 2016). "How our ancestors drilled rotten teeth". BBC. Archived from the original on 19 May 2017. Retrieved 1 March 2016.
33. ^ "Oldest Dentistry Found in 14,000-Year-Old Tooth". Discovery Channel. 16 July 2015. Archived from the original on 18 July 2015. Retrieved 21 July 2015.
34. ^ "Analysis of Neanderthal teeth marks uncovers evidence of prehistoric dentistry". The University of Kansas. 28 June 2017. Retrieved 1 July 2017.
35. ^ Oxilia, Gregorio; Fiorillo, Flavia; Boschin, Francesco; Boaretto, Elisabetta; Apicella, Salvatore A.; Matteucci, Chiara; Panetta, Daniele; Pistocchi, Rossella; Guerrini, Franca; Margherita, Cristiana; Andretta, Massimo; Sorrentino, Rita; Boschian, Giovanni; Arrighi, Simona; Dori, Irene (2017). "The dawn of dentistry in the late upper Paleolithic: An early case of pathological intervention at Riparo Fredian". *American Journal of Physical Anthropology*. **163** (3): 446–461. doi:10.1002/ajpa.23216. hdl:11585/600517. ISSN 0002-9483. PMID 28345756.
36. ^ Bernardini, Federico; Tuniz, Claudio; Coppa, Alfredo; Mancini, Lucia; Dreossi, Diego; Eichert, Diane; Turco, Gianluca; Biasotto, Matteo; Terrasi, Filippo; Cesare, Nicola De; Hua, Quan; Levchenko, Vladimir (19 September 2012). "Beeswax as Dental Filling on a Neolithic Human Tooth". *PLOS ONE*. **7** (9): e44904. Bibcode:2012PLoSO...744904B. doi:10.1371/journal.pone.0044904. ISSN 1932-6203. PMC 3446997. PMID 23028670.
37. ^ Coppa, A.; et al. (2006). "Early Neolithic tradition of dentistry". *Nature*. **440** (7085). Springer Science and Business Media LLC: 755–756. doi:10.1038/440755a. ISSN 0028-0836. PMID 16598247.
38. ^ "Dig uncovers ancient roots of dentistry". NBC News. 5 April 2006.
39. ^ Bernardini, Federico; et al. (2012). "Beeswax as Dental Filling on a Neolithic Human Tooth". *PLOS ONE*. **7** (9): e44904. Bibcode:2012PLoSO...744904B. doi:10.1371/journal.pone.0044904. PMC 3446997. PMID 23028670.
40. ^ "700 years added to Malta's history". Times of Malta. 16 March 2018. Archived from the original on 16 March 2018.
41. ^ "History of Dentistry: Ancient Origins". American Dental Association. Archived from the original on 5 July 2007. Retrieved 9 January 2007.

42. ^ TOWNEND, B. R. (1944). "The Story of the Tooth-Worm". *Bulletin of the History of Medicine*. **15** (1): 37–58. ISSN 0007-5140. JSTOR 44442797.
43. ^ Suddick Richard P., Harris Norman O. (1990). "Historical Perspectives of Oral Biology: A Series" (PDF). *Critical Reviews in Oral Biology and Medicine*. **1** (2): 135–51. doi:10.1177/10454411900010020301. PMID 2129621. Archived from the original (PDF) on 18 December 2007.
44. ^ a b c Blomstedt, P. (2013). "Dental surgery in ancient Egypt". *Journal of the History of Dentistry*. **61** (3): 129–42. PMID 24665522.
45. ^ "Ancient Egyptian Dentistry". University of Oklahoma. Archived from the original on 26 December 2007. Retrieved 15 December 2007.
46. ^ Wilwerding, Terry. "History of Dentistry 2001" (PDF). Archived from the original (PDF) on 3 November 2014. Retrieved 3 November 2014.
47. ^ "Medicine in Ancient Egypt 3". Arabworldbooks.com. Retrieved 18 April 2010.
48. ^ a b "History Of Dentistry". Complete Dental Guide. Archived from the original on 14 July 2016. Retrieved 29 June 2016.
49. ^ "History of Dentistry Research Page, Newsletter". Rcpsg.ac.uk. Archived from the original on 28 April 2015. Retrieved 9 June 2014.
50. ^ a b Donaldson, J. A. (1980). "The use of gold in dentistry" (PDF). *Gold Bulletin*. **13** (3): 117–124. doi:10.1007/BF03216551. PMID 11614516. S2CID 137571298.
51. ^ Becker, Marshall J. (1999). Ancient "dental implants": a recently proposed example from France evaluated with other spurious examples (PDF). *International Journal of Oral & Maxillofacial Implants* 14.1.
52. ^ Malik, Ursman. "History of Dentures from Beginning to Early 19th Century". Exhibits. Retrieved 3 May 2023.
53. ^ Renfrew, Colin; Bahn, Paul (2012). *Archaeology: Theories, Methods, and Practice* (6th ed.). Thames & Hudson. p. 449. ISBN 978-0-500-28976-1.
54. ^ "Dental Treatment in the Ancient Times". Dentaltreatment.org.uk. Archived from the original on 1 December 2009. Retrieved 18 April 2010.
55. ^ Bjørklund G (1989). "The history of dental amalgam (in Norwegian)". *Tidsskr Nor Laegeforen*. **109** (34–36): 3582–85. PMID 2694433.
56. ^ Czarnetzki, A.; Ehrhardt S. (1990). "Re-dating the Chinese amalgam-filling of teeth in Europe". *International Journal of Anthropology*. **5** (4): 325–32.
57. ^ Meri, Josef (2005). *Medieval Islamic Civilization: An Encyclopedia* (Routledge Encyclopedias of the Middle Ages). Psychology Press. ISBN 978-0-415-96690-0.
58. ^ Friedman, Saul S. (2006). *A history of the Middle East*. Jefferson, N.C.: Mcfarland. p. 152. ISBN 0786451343.








59. ^ Gregory Ribitzky. "Pelican". Archived from the original on 25 January 2020. Retrieved 23 June 2018.
60. ^ Gregory Ribitzky. "Toothkey". Archived from the original on 23 June 2018. Retrieved 23 June 2018.
61. ^ Gregory Ribitzky. "Forceps". Archived from the original on 23 June 2018. Retrieved 23 June 2018.
62. ^ Royal Commission on the NHS Chapter 9. HMSO. July 1979. ISBN 978-0-10-176150-5. Retrieved 19 May 2015.
63. ^ **a b** André Besombes; Phillipe de Gaillande (1993). *Pierre Fauchard (1678–1761): The First Dental Surgeon, His Work, His Actuality*. Pierre Fauchard Academy.
64. ^ **a b** Bernhard Wolf Weinberger (1941). *Pierre Fauchard, Surgeon-dentist: A Brief Account of the Beginning of Modern Dentistry, the First Dental Textbook, and Professional Life Two Hundred Years Ago*. Pierre Fauchard Academy.
65. ^ Moore, Wendy (30 September 2010). *The Knife Man*. Transworld. pp. 223–24. ISBN 978-1-4090-4462-8. Retrieved 8 March 2012.
66. ^ Dingwall, Helen (April 2004). "A pioneering history: dentistry and the Royal College of Surgeons of Edinburgh" (PDF). *History of Dentistry Newsletter*. No. 14. Archived from the original (PDF) on 1 February 2013.
67. ^ "Noise-Induced Hearing Loss". NIDCD. 18 August 2015.
68. ^ "Occupational Safety and Health Standards | Occupational Safety and Health Administration". Osha.gov.
69. ^ Stevens, M (1999). "Is someone listening to the din of occupational noise exposure in dentistry". *RDH* (19): 34–85.
70. ^ Merrel, HB (1992). "Noise pollution and hearing loss in the dental office". *Dental Assisting Journal*. **61** (3): 6–9.
71. ^ Wilson, J.D. (2002). "Effects of occupational ultrasonic noise exposure on hearing of dental hygienists: A pilot study". *Journal of Dental Hygiene*. **76** (4): 262–69. PMID 12592917.
72. ^ Leggat, P.A. (2007). "Occupational Health Problems in Modern Dentistry: A Review" (PDF). *Industrial Health*. **45** (5): 611–21. doi:10.2486/indhealth.45.611. PMID 18057804. Archived (PDF) from the original on 27 April 2019.
73. ^ Leggat, P.A. (2001). "Occupational hygiene practices of dentists in southern Thailand". *International Dental Journal*. **51** (51): 11–6. doi:10.1002/j.1875-595x.2001.tb00811.x. PMID 11326443.

74. ^ *Evidence-Based Medicine Working Group* (1992). "Evidence-based medicine. A new approach to teaching the practice of medicine". *Journal of the American Medical Association*. **268** (17): 2420–2425. doi:10.1001/jama.1992.03490170092032. PMID 1404801.
75. ^ "Union workers build high-tech dental simulation laboratory for SIU dental school". *The Labor Tribune*. 17 March 2014. Retrieved 10 September 2021.
76. ^ Slavkin, Harold C. (January 2012). "Evolution of the scientific basis for dentistry and its impact on dental education: past, present, and future". *Journal of Dental Education*. **76** (1): 28–35. doi:10.1002/j.0022-0337.2012.76.1.tb05231.x. ISSN 1930-7837. PMID 22262547.
77. ^ Formicola, Allan J.; Bailit, Howard L.; Beazoglou, Tryfon J.; Tedesco, Lisa A. (February 2008). "The interrelationship of accreditation and dental education: history and current environment". *Journal of Dental Education*. **72** (2 Suppl): 53–60. doi:10.1002/j.0022-0337.2008.72.2_suppl.tb04480.x. ISSN 0022-0337. PMID 18250379.
78. ^ Carrassi, A. (2019). "The first 25 year [Internet] Ireland: ADEE (Association for Dental Education in Europe)". Association for Dental Education in Europe. Retrieved 21 October 2019.

External links

[edit]

Dentistry at Wikipedia's sister projects

-  Definitions from Wiktionary
-  Media from Commons
-  Quotations from Wikiquote
-  Texts from Wikisource
-  Textbooks from Wikibooks
-  Resources from Wikiversity
-  Data from Wikidata

- v
- t
- e

Dentistry

Specialties

- Endodontics
- Oral and maxillofacial pathology
- Oral and maxillofacial radiology
- Oral and maxillofacial surgery
- Orthodontics and dentofacial orthopedics
- Pediatric dentistry
- Periodontics
- Prosthodontics
- Dental public health
- Cosmetic dentistry
- Dental implantology
- Geriatric dentistry
- Restorative dentistry
- Forensic odontology
- Dental traumatology
- Holistic dentistry
- Dental extraction
- Tooth filling
- Root canal therapy
- Root end surgery
- Scaling and root planing

Dental surgery

- Teeth cleaning
- Dental bonding
- Tooth polishing
- Tooth bleaching
- Socket preservation
- Dental implant

Organisations

- American Association of Orthodontists
- British Dental Association
- British Dental Health Foundation
- British Orthodontic Society
- Canadian Association of Orthodontists
- Dental Technologists Association
- General Dental Council
- Indian Dental Association
- National Health Service
- Canada
- Philippines

By country

- Israel
- United Kingdom
- United States
- Index of oral health and dental articles
- Outline of dentistry and oral health
- Dental fear
- Dental instruments

See also

- Dental material
- History of dental treatments
 - Ancient Rome
- Infant oral mutilation
- Mouth assessment
- Oral hygiene

- v
- t
- e

Cleft lip and cleft palate

**Related
specialities**

- Advance practice nursing
- Audiology
- Dentistry
- Dietetics
- Genetics
- Oral and maxillofacial surgery
- Orthodontics
- Orthodontic technology
- Otolaryngology
- Pediatrics
- Pediatric dentistry
- Physician
- Plastic surgery
- Psychiatry
- Psychology
- Respiratory therapy
- Social work
- Speech and language therapy
- Hearing loss with craniofacial syndromes

**Related
syndromes**

- Pierre Robin syndrome
- Popliteal pterygium syndrome
- Van der Woude syndrome

**National and
international
organisations**

- Cleft Lip and Palate Association
- Craniofacial Society of Great Britain and Ireland
- Interplast
- North Thames Regional Cleft Lip and Palate Service
- Operation Smile
- Overseas Plastic Surgery Appeal
- Shriners Hospitals for Children
- Smile Train
- Transforming Faces Worldwide
- Smile Angel Foundation (China)

- v
- t
- e

Dental schools

**American
dental
schools**

- UAB
- Arizona
- Augusta (DCG)
- Boston U (Goldman)
- California (UCLA, UCSF)
- Case Western Reserve
- Colorado
- Columbia
- Connecticut
- Creighton
- Detroit Mercy
- East Carolina
- Florida
- Harvard
- Howard
- Illinois–Chicago
- Indiana
- Iowa
- Kentucky
- Lake Erie
- Loma Linda
- Louisville
- LSU Health–New Orleans
- Marquette
- Maryland–Baltimore
- Meharry
- Michigan
- Midwestern
- Minnesota
- Mississippi
- Missouri–Kansas City
- Nebraska–Medical Center
- Nevada–Las Vegas
- New England

	<ul style="list-style-type: none"> ○ Emory ○ Fairleigh Dickinson ○ Georgetown
Defunct	<ul style="list-style-type: none"> ○ Harris
American	<ul style="list-style-type: none"> ○ Loyola
dental	<ul style="list-style-type: none"> ○ Northwestern
schools	<ul style="list-style-type: none"> ○ Ohio College ○ Oral Roberts ○ Pennsylvania College ○ Wash U ○ Alberta ○ British Columbia ○ Dalhousie
Canadian	<ul style="list-style-type: none"> ○ Laval
dental	<ul style="list-style-type: none"> ○ Manitoba
schools	<ul style="list-style-type: none"> ○ McGill ○ Montréal ○ Saskatchewan ○ Toronto ○ Western ○ Aberdeen
	<ul style="list-style-type: none"> ○ Barts and The London School of Medicine and Dentistry
British	<ul style="list-style-type: none"> ○ Glasgow
dental	<ul style="list-style-type: none"> ○ Guy's, King's & St Thomas's
schools	<ul style="list-style-type: none"> ○ Liverpool ○ Newcastle ○ Peninsula College of Medicine and Dentistry ○ UCL Eastman Dental Institute

**Australian
and New
Zealand
dental
schools**

- Sydney
- Melbourne
- Adelaide
- Charles Sturt University
- Griffith University
- James Cook
- La Trobe
- Queensland
- Western Australia
- University of Otago

**South
Korean
dental
schools**

- Chonbuk
- Chonnam
- Chosun
- Dankook
- Gangneung-Wonju
- Kyung Hee
- Kyungpook
- Pusan
- Seoul
- Wonkwang
- Yonsei

- v
- t
- e

Medicine

Surgery

- Cardiac surgery
- Cardiothoracic surgery
- Endocrine surgery
- Eye surgery
- General surgery
 - Colorectal surgery
 - Digestive system surgery
- Neurosurgery
- Oral and maxillofacial surgery
- Orthopedic surgery
- Hand surgery
- Otolaryngology
 - ENT
- Pediatric surgery
- Plastic surgery
- Reproductive surgery
- Surgical oncology
- Transplant surgery
- Trauma surgery
- Urology
 - Andrology
- Vascular surgery
- Allergy / Immunology
- Angiology
- Cardiology
- Endocrinology
- Gastroenterology
 - Hepatology


Internal medicine

- Geriatrics
- Hematology
- Hospital medicine
- Infectious diseases
- Nephrology

Medical education

- Medical school
- Bachelor of Medicine, Bachelor of Surgery
- Bachelor of Medical Sciences
- Master of Medicine
- Master of Surgery
- Doctor of Medicine
- Doctor of Osteopathic Medicine
- MD–PhD
 - Medical Scientist Training Program
- Alternative medicine
- Allied health
- Molecular oncology
- Nanomedicine
- Personalized medicine
- Public health
- Rural health
- Therapy
- Traditional medicine
- Veterinary medicine
- Physician
 - Chief physician
- History of medicine

Related topics

-  **Category**
-  **Commons**
-  **WikiProject**
-  **Portal**
-  **Outline**

Authority control databases  **Edit this at Wikidata**

National

- Germany
- United States
- France
- BnF data
- Czech Republic
- Latvia
- Israel

Other

- Historical Dictionary of Switzerland
- NARA

About dental caries

Redirect to:


- Tooth decay
- **From a page move:** This is a redirect from a page that has been moved (renamed). This page was kept as a redirect to avoid breaking links, both internal and external, that may have been made to the old page name.

About dental braces

This article **needs additional citations for verification**. Please help improve this article by adding citations to reliable sources. Unsourced material may be challenged and removed.



Find sources: "Dental braces" – news · newspapers · books · scholar · JSTOR (August 2016) *(Learn how and when to remove this message)*



Dental braces

Dental braces (also known as **orthodontic braces**, or simply **braces**) are devices used in orthodontics that align and straighten teeth and help position them with regard to a person's bite, while also aiming to improve dental health. They are often used to correct underbites, as well as malocclusions, overbites, open bites, gaps, deep bites, cross bites, crooked teeth, and various other flaws of the teeth and jaw. Braces can be either cosmetic or structural. Dental braces are often used in conjunction with other orthodontic appliances to help widen the palate or jaws and to otherwise assist in shaping the teeth and jaws.

Process

[edit]


The application of braces moves the teeth as a result of force and pressure on the teeth. Traditionally, four basic elements are used: brackets, bonding material, arch wire, and ligature elastic (also called an "O-ring"). The teeth move when the arch wire puts pressure on the brackets and teeth. Sometimes springs or rubber bands are used to put more force in a specific direction.^[1]

Braces apply constant pressure which, over time, moves teeth into the desired positions. The process loosens the tooth after which new bone grows to support the tooth in its new position. This is called bone remodelling. Bone remodelling is a biomechanical process responsible for making bones stronger in response to sustained load-bearing activity and weaker in the absence of carrying a load.


Bones are made of cells called osteoclasts and osteoblasts. Two different kinds of bone resorption are possible: direct resorption, which starts from the lining cells of the alveolar bone, and indirect or retrograde resorption, which occurs when the periodontal ligament has been subjected to an excessive amount and duration of compressive stress.^[2] Another important factor associated with tooth movement is bone deposition. Bone deposition occurs in the distracted periodontal ligament. Without bone deposition, the tooth will loosen, and voids will occur distal to the direction of tooth movement.^[3]

Types

[edit]



"Clear" braces



Upper and Lower Jaw Functional Expanders


- **Traditional metal wired braces** (also known as "train track braces") are stainless-steel and are sometimes used in combination with titanium. Traditional metal braces are the most common type of braces.^[4] These braces have a metal bracket with elastic ties (also known as rubber bands)

holding the wire onto the metal brackets. The second-most common type of braces is self-ligating braces, which have a built-in system to secure the archwire to the brackets and do not require elastic ties. Instead, the wire goes through the bracket. Often with this type of braces, treatment time is reduced, there is less pain on the teeth, and fewer adjustments are required than with traditional braces.

- **Gold-plated stainless steel** braces are often employed for patients allergic to nickel (a basic and important component of stainless steel), but may also be chosen for aesthetic reasons.
- **Lingual braces** are a cosmetic alternative in which custom-made braces are bonded to the back of the teeth making them externally invisible.
- **Titanium braces** resemble stainless-steel braces but are lighter and just as strong. People with allergies to nickel in steel often choose titanium braces, but they are more expensive than stainless steel braces.
- **Customized orthodontic treatment systems** combine high technology including 3-D imaging, treatment planning software and a robot to custom bend the wire. Customized systems such as this offer faster treatment times and more efficient results.^[5]
- **Progressive, clear removable aligners** may be used to gradually move teeth into their final positions. Aligners are generally not used for complex orthodontic cases, such as when extractions, jaw surgery, or palate expansion are necessary.^{[medical citation needed][6]}

Fitting procedure

[edit]



A patient's teeth are prepared for the application of braces.

Orthodontic services may be provided by any licensed dentist trained in orthodontics. In North America, most orthodontic treatment is done by orthodontists, who are dentists in the diagnosis and treatment of *malocclusions*—malalignments of the teeth, jaws, or both. A dentist must complete 2–3 years of additional post-doctoral training to earn a specialty certificate in orthodontics. There are many general practitioners who also provide orthodontic services.


The first step is to determine whether braces are suitable for the patient. The doctor consults with the patient and inspects the teeth visually. If braces are appropriate, a records appointment is set up where X-rays, moulds, and impressions are made. These records are analyzed to determine the problems and the proper course of action. The use of digital models is rapidly increasing in the orthodontic industry. Digital treatment starts with the creation of a three-dimensional digital model of the patient's arches. This model is produced by laser-scanning plaster models created using dental impressions. Computer-automated treatment simulation has the ability to automatically separate the gums and teeth from one another and can handle malocclusions well; this software enables clinicians to ensure, in a virtual setting, that the selected treatment will produce the optimal outcome, with minimal user input.^[*medical citation needed*]

Typical treatment times vary from six months to two and a half years depending on the complexity and types of problems. Orthognathic surgery may be required in extreme cases. About 2 weeks before the braces are applied, orthodontic

spacers may be required to spread apart back teeth in order to create enough space for the bands.

Teeth to be braced will have an adhesive applied to help the cement bond to the surface of the tooth. In most cases, the teeth will be banded and then brackets will be added. A bracket will be applied with dental cement, and then cured with light until hardened. This process usually takes a few seconds per tooth. If required, orthodontic spacers may be inserted between the molars to make room for molar bands to be placed at a later date. Molar bands are required to ensure brackets will stick. Bands are also utilized when dental fillings or other dental works make securing a bracket to a tooth infeasible. Orthodontic tubes (stainless steel tubes that allow wires to pass through them), also known as molar tubes, are directly bonded to molar teeth either by a chemical curing or a light curing adhesive. Usually, molar tubes are directly welded to bands, which is a metal ring that fits onto the molar tooth. Directly bonded molar tubes are associated with a higher failure rate when compared to molar bands cemented with glass ionomer cement. Failure of orthodontic brackets, bonded tubes or bands will increase the overall treatment time for the patient. There is evidence suggesting that there is less enamel decalcification associated with molar bands cemented with glass ionomer cement compared with orthodontic tubes directly cemented to molars using a light cured adhesive. Further evidence is needed to withdraw a more robust conclusion due to limited data.^[7]

An archwire will be threaded between the brackets and affixed with elastic or metal ligatures. Ligatures are available in a wide variety of colours, and the patient can choose which colour they like. Arch wires are bent, shaped, and tightened frequently to achieve the desired results.



Dental braces, with a transparent power chain, removed after completion of treatment.

Modern orthodontics makes frequent use of nickel–titanium archwires and temperature–sensitive materials. When cold, the archwire is limp and flexible, easily threaded between brackets of any configuration. Once heated to body temperature, the arch wire will stiffen and seek to retain its shape, creating constant light force on the teeth.


Brackets with hooks can be placed, or hooks can be created and affixed to the arch wire to affix rubber bands. The placement and configuration of the rubber bands will depend on the course of treatment and the individual patient. Rubber bands are made in different diameters, colours, sizes, and strengths. They are also typically available in two versions: Coloured or clear/opaque.

The fitting process can vary between different types of braces, though there are similarities such as the initial steps of moulding the teeth before application. For example, with clear braces, impressions of a patient's teeth are evaluated to create a series of trays, which fit to the patient's mouth almost like a protective mouthpiece. With some forms of braces, the brackets are placed in a special form that is customized to the patient's mouth, drastically reducing the application time.

In many cases, there is insufficient space in the mouth for all the teeth to fit properly. There are two main procedures to make room in these cases. One is extraction: teeth are removed to create more space. The second is expansion, in which the palate or arch is made larger by using a palatal expander. Expanders can

be used with both children and adults. Since the bones of adults are already fused, expanding the palate is not possible without surgery to separate them. An expander can be used on an adult without surgery but would be used to expand the dental arch, and not the palate.

Sometimes children and teenage patients, and occasionally adults, are required to wear a headgear appliance as part of the primary treatment phase to keep certain teeth from moving (for more detail on headgear and facemask appliances see Orthodontic headgear). When braces put pressure on one's teeth, the periodontal membrane stretches on one side and is compressed on the other. This movement needs to be done slowly or otherwise, the patient risks losing their teeth. This is why braces are worn as long as they are and adjustments are only made every so often.



Young Colombian man during an adjustment visit for his orthodontics

Braces are typically adjusted every three to six weeks. This helps shift the teeth into the correct position. When they get adjusted, the orthodontist removes the coloured or metal ligatures keeping the arch wire in place. The arch wire is then removed and may be replaced or modified. When the archwire has been placed back into the mouth, the patient may choose a colour for the new elastic ligatures,

which are then affixed to the metal brackets. The adjusting process may cause some discomfort to the patient, which is normal.

Post-treatment


[edit]

Patients may need post-orthodontic surgery, such as a fiberotomy or alternatively a gum lift, to prepare their teeth for retainer use and improve the gumline contours after the braces come off. After braces treatment, patients can use a transparent plate to keep the teeth in alignment for a certain period of time. After treatment, patients usually use transparent plates for 6 months. In patients with long and difficult treatment, a fixative wire is attached to the back of the teeth to prevent the teeth from returning to their original state.^[8]

Retainers

[edit]

Main article: Retainer (orthodontic device)



Hawley retainers are the most common type of retainers. This picture shows retainers for the top (right) and bottom (left) of the mouth.

In order to prevent the teeth from moving back to their original position, retainers are worn once the treatment is complete. Retainers help in maintaining and stabilizing the position of teeth long enough to permit the reorganization of the supporting structures after the active phase of orthodontic therapy. If the patient does not wear the retainer appropriately and/or for the right amount of time, the

teeth may move towards their previous position. For regular braces, Hawley retainers are used. They are made of metal hooks that surround the teeth and are enclosed by an acrylic plate shaped to fit the patient's palate. For Clear Removable braces, an Essix retainer is used. This is similar to the original aligner; it is a clear plastic tray that is firmly fitted to the teeth and stays in place without a plate fitted to the palate. There is also a bonded retainer where a wire is permanently bonded to the lingual side of the teeth, usually the lower teeth only.


Headgear

[edit]

Main article: Orthodontic headgear

Headgear needs to be worn between 12 and 22 hours each day to be effective in correcting the overbite, typically for 12 to 18 months depending on the severity of the overbite, how much it is worn and what growth stage the patient is in. Typically the prescribed daily wear time will be between 14 and 16 hours a day and is frequently used as a post-primary treatment phase to maintain the position of the jaw and arch. Headgear can be used during the night while the patient sleeps.^[9]^[better source needed]

Orthodontic headgear usually consists of three major components:



Full orthodontic headgear with head cap, fitting straps, facebow and elastics

1. Facebow: the facebow (or J-Hooks) is fitted with a metal arch onto headgear tubes attached to the rear upper and lower molars. This facebow then extends out of the mouth and around the patient's face. J-Hooks are different in that they hook into the patient's mouth and attach directly to the brace (see photo for an example of J-Hooks).
2. Head cap: the head cap typically consists of one or a number of straps fitting around the patient's head. This is attached with elastic bands or springs to the facebow. Additional straps and attachments are used to ensure comfort and safety (see photo).
3. Attachment: typically consisting of rubber bands, elastics, or springs—joins the facebow or J-Hooks and the head cap together, providing the force to move the upper teeth, jaw backwards.

The headgear application is one of the most useful appliances available to the orthodontist when looking to correct a Class II malocclusion. See more details in the section Orthodontic headgear.

Pre-finisher

[edit]

The pre-finisher is moulded to the patient's teeth by use of extreme pressure on the appliance by the person's jaw. The product is then worn a certain amount of time with the user applying force to the appliance in their mouth for 10 to 15 seconds at a time. The goal of the process is to increase the exercise time in applying the force to the appliance. If a person's teeth are not ready for a proper retainer the orthodontist may prescribe the use of a preformed finishing appliance such as the pre-finisher. This appliance fixes gaps between the teeth, small spaces between the upper and lower jaw, and other minor problems.

Complications and risks

[edit]

A group of dental researchers, Fatma Boke, Cagri Gazioglu, Selvi Akkaya, and Murat Akkaya, conducted a study titled "Relationship between orthodontic treatment and gingival health." The results indicated that some orthodontist treatments result in gingivitis, also known as gum disease. The researchers concluded that functional appliances used to harness natural forces (such as improving the alignment of bites) do not usually have major effects on the gum after treatment.^[10] However, fixed appliances such as braces, which most people get, can result in visible plaque, visible inflammation, and gum recession in a majority of the patients. The formation of plaques around the teeth of patients with braces is almost inevitable regardless of plaque control and can result in mild gingivitis. But if someone with braces does not clean their teeth carefully, plaques will form, leading to more severe gingivitis and gum recession.

Experiencing some pain following fitting and activation of fixed orthodontic braces is very common and several methods have been suggested to tackle this.^{[11][12]}

Pain associated with orthodontic treatment increases in proportion to the amount of force that is applied to the teeth. When a force is applied to a tooth via a brace, there is a reduction in the blood supply to the fibres that attach the tooth to the surrounding bone. This reduction in blood supply results in inflammation and the release of several chemical factors, which stimulate the pain response.

Orthodontic pain can be managed using pharmacological interventions, which involve the use of analgesics applied locally or systemically. These analgesics are divided into four main categories, including opioids, non-steroidal anti-inflammatory drugs (NSAIDs), paracetamol and local anesthesia. The first three of these analgesics are commonly taken systemically to reduce orthodontic pain.^[13]

A Cochrane Review in 2017 evaluated the pharmacological interventions for pain relief during orthodontic treatment. The study concluded that there was moderate-quality evidence that analgesics reduce the pain associated with orthodontic treatment. However, due to a lack of evidence, it was unclear whether systemic NSAIDs were more effective than paracetamol, and whether topical NSAIDs were more effective than local anaesthesia in the reduction of pain associated with orthodontic treatment. More high-quality research is required to investigate these particular comparisons.^[13]


The dental displacement obtained with the orthodontic appliance determines in most cases some degree of root resorption. Only in a few cases is this side effect large enough to be considered real clinical damage to the tooth. In rare cases, the teeth may fall out or have to be extracted due to root resorption.^{[14][15]}

History

[edit]

Ancient

[edit]




Old Braces at a museum in Jbeil, Lebanon

According to scholars and historians, braces date back to ancient times. Around 400–300 BC, Hippocrates and Aristotle contemplated ways to straighten teeth and fix various dental conditions. Archaeologists have discovered numerous mummified ancient individuals with what appear to be metal bands wrapped around their teeth. Catgut, a type of cord made from the natural fibres of an animal's intestines, performed a similar role to today's orthodontic wire in closing gaps in the teeth and mouth.^[16]

The Etruscans buried their dead with dental appliances in place to maintain space and prevent the collapse of the teeth during the afterlife. A Roman tomb was found with a number of teeth bound with gold wire documented as a ligature wire, a small elastic wire that is used to affix the arch wire to the bracket. Even Cleopatra wore a pair. Roman philosopher and physician Aulus Cornelius Celsus first recorded the treatment of teeth by finger pressure. Unfortunately, due to a lack of evidence, poor preservation of bodies, and primitive technology, little research was carried out on dental braces until around the 17th century, although dentistry was making great advancements as a profession by then.^[citation needed]

18th century

[edit]



Portrait of Fauchard from his 1728 edition of *"The Surgical Dentist"*.

Orthodontics truly began developing in the 18th and 19th centuries. In 1669, French dentist Pierre Fauchard, who is often credited with inventing modern orthodontics, published a book entitled *"The Surgeon Dentist"* on methods of straightening teeth. Fauchard, in his practice, used a device called a "Bandeau", a horseshoe-shaped piece of iron that helped expand the palate. In 1754, another French dentist, Louis Bourdet, dentist to the King of France, followed Fauchard's book with *The Dentist's Art*, which also dedicated a chapter to tooth alignment and application. He perfected the "Bandeau" and was the first dentist on record to recommend extraction of the premolar teeth to alleviate crowding and improve jaw growth.

19th century

[edit]

Although teeth and palate straightening and/or pulling were used to improve the alignment of remaining teeth and had been practised since early times,

orthodontics, as a science of its own, did not really exist until the mid-19th century. Several important dentists helped to advance dental braces with specific instruments and tools that allowed braces to be improved.

In 1819, Christophe François Delabarre introduced the wire crib, which marked the birth of contemporary orthodontics, and gum elastics were first employed by Maynard in 1843. Tucker was the first to cut rubber bands from rubber tubing in 1850. Dentist, writer, artist, and sculptor Norman William Kingsley in 1858 wrote the first article on orthodontics and in 1880, his book, *Treatise on Oral Deformities*, was published. A dentist named John Nutting Farrar is credited for writing two volumes entitled, *A Treatise on the Irregularities of the Teeth and Their Corrections* and was the first to suggest the use of mild force at timed intervals to move teeth.

20th century


[edit]

In the early 20th century, Edward Angle devised the first simple classification system for malocclusions, such as Class I, Class II, and so on. His classification system is still used today as a way for dentists to describe how crooked teeth are, what way teeth are pointing, and how teeth fit together. Angle contributed greatly to the design of orthodontic and dental appliances, making many simplifications. He founded the first school and college of orthodontics, organized the American Society of Orthodontia in 1901 which became the American Association of Orthodontists (AAO) in the 1930s, and founded the first orthodontic journal in 1907. Other innovations in orthodontics in the late 19th and early 20th centuries included the first textbook on orthodontics for children, published by J.J. Guilford in 1889, and the use of rubber elastics, pioneered by Calvin S. Case, along with Henry Albert Baker.

Today, space age wires (also known as dental arch wires) are used to tighten braces. In 1959, the Naval Ordnance Laboratory created an alloy of nickel and titanium called Nitinol. NASA further studied the material's physical properties.^[17] In 1979, Dr. George Andreasen developed a new method of fixing braces with the use of the Nitinol wires based on their superelasticity. Andreasen used the wire on some patients and later found out that he could use it for the entire treatment. Andreasen then began using the nitinol wires for all his treatments and as a result, dental doctor visits were reduced, the cost of dental treatment was reduced, and patients reported less discomfort.

See also

[edit]

- o  Medicine portal
- o Mandibular advancement splint
- o Oral and maxillofacial surgery
- o Orthognathic surgery
- o Prosthodontics
- o Trismus
- o Dental implant

References

[edit]

1. ^ "Dental Braces and Retainers". *WebMD*. Retrieved 2020-10-30.
2. ^ Robling, Alexander G.; Castillo, Alesha B.; Turner, Charles H. (2006). "Biomechanical and Molecular Regulation of Bone Remodeling". *Annual Review of Biomedical Engineering*. **8**: 455–498. doi:10.1146/annurev.bioeng.8.061505.095721. PMID 16834564.
3. ^ Toledo SR, Oliveira ID, Okamoto OK, Zago MA, de Seixas Alves MT, Filho RJ, et al. (September 2010). "Bone deposition, bone resorption, and osteosarcoma". *Journal of Orthopaedic Research*. **28** (9): 1142–1148. doi:10.1002/jor.21120. PMID 20225287. S2CID 22660771.


4. ^ "Metal Braces for Teeth: Braces Types, Treatment, Cost in India". Clove Dental . Retrieved 2025-02-06.
5. ^ Saxe, Alana K.; Louie, Lenore J.; Mah, James (2010). "Efficiency and effectiveness of SureSmile". *World Journal of Orthodontics*. **11** (1): 16–22. PMID 20209172.
6. ^ Tamer, Ãfâ€žÃ,Â°pek (December 2019). "Orthodontic Treatment with Clear Aligners and The Scientific Reality Behind Their Marketing: A Literature Review". *Turkish Journal of Orthodontics*. **32** (4): 241–246. doi:10.5152/TurkJOrthod.2019.18083. PMC 7018497. PMID 32110470.
7. ^ Millett DT, Mandall NA, Mattick RC, Hickman J, Glenny AM (February 2017). "Adhesives for bonded molar tubes during fixed brace treatment". *The Cochrane Database of Systematic Reviews*. **2** (3): CD008236. doi:10.1002/14651858.cd008236.pub3. PMC 6464028. PMID 28230910.
8. ^ Rubie J Patrick (2017). "What About Teeth After Braces?" 2017 – "Health Journal Article" Toothcost Archived 2021-10-18 at the Wayback Machine
9. ^ Naten, Joshua. "Braces Headgear (Treatments)". toothcost.com. Archived from the original on 19 October 2021.
10. ^ Boke, Fatma; Gazioglu, Cagri; Akkaya, Sevil; Akkaya, Murat (2014). "Relationship between orthodontic treatment and gingival health: A retrospective study". *European Journal of Dentistry*. **8** (3): 373–380. doi:10.4103/1305-7456.137651. ISSN 1305-7456. PMC 4144137. PMID 25202219.
11. ^ Eslamian L, Borzabadi-Farahani A, Hassanzadeh-Azhiri A, Badiie MR, Fekrazad R (March 2014). "The effect of 810-nm low-level laser therapy on pain caused by orthodontic elastomeric separators". *Lasers in Medical Science*. **29** (2): 559–64. doi:10.1007/s10103-012-1258-1. PMID 23334785. S2CID 25416518.
12. ^ Eslamian L, Borzabadi-Farahani A, Edini HZ, Badiie MR, Lynch E, Mortazavi A (September 2013). "The analgesic effect of benzocaine mucoadhesive patches on orthodontic pain caused by elastomeric separators, a preliminary study". *Acta Odontologica Scandinavica*. **71** (5): 1168–73. doi:10.3109/00016357.2012.757358. PMID 23301559. S2CID 22561192.
13. ^ **a b** Monk AB, Harrison JE, Worthington HV, Teague A (November 2017). "Pharmacological interventions for pain relief during orthodontic treatment". *The Cochrane Database of Systematic Reviews*. **11** (12): CD003976. doi:10.1002/14651858.cd003976.pub2. PMC 6486038. PMID 29182798.
14. ^ Artun J, Smale I, Behbehani F, Doppel D, Van't Hof M, Kuijpers-Jagtman AM (November 2005). "Apical root resorption six and 12 months after initiation of fixed orthodontic appliance therapy". *The Angle Orthodontist*. **75** (6): 919–26.

PMID 16448232.

15. ^ Mavragani M, Vergari A, Selliseth NJ, Bøe OE, Wisth PL (December 2000). "A radiographic comparison of apical root resorption after orthodontic treatment with a standard edgewise and a straight-wire edgewise technique". *European Journal of Orthodontics*. **22** (6): 665–74. doi:10.1093/ejo/22.6.665. PMID 11212602.
16. ^ Wahl N (February 2005). "Orthodontics in 3 millennia. Chapter 1: Antiquity to the mid-19th century". *American Journal of Orthodontics and Dentofacial Orthopedics*. **127** (2): 255–9. doi:10.1016/j.ajodo.2004.11.013. PMID 15750547.
17. ^ "NASA Technical Reports Server (NTRS)". Spinoff 1979. February 1979. Retrieved 2021-03-02.

External links

[edit]

- Useful Resources: FAQ and Downloadable eBooks at Orthodontics Australia
- Orthos Explain: Treatment Options at Orthodontics Australia
-  Media related to Dental braces at Wikimedia Commons
- v
- t
- e

Orthodontics

- Bolton analysis
- Cephalometric analysis
- Cephalometry
- Dentition analysis
- Failure of eruption of teeth

Diagnosis

- Little's Irregularity Index
- Malocclusion
- Scissor bite
- Standard anatomical position
- Tooth ankylosis
- Tongue thrust

Conditions

- Overbite
- Overjet
- Open bite
- Crossbite
- Dental crowding
- Dental spacing
- Bimaxillary Protrusion
- Prognathism
- Retrognathism
- Maxillary hypoplasia
- Condylar hyperplasia
- Overeruption
- Mouth breathing
- Temporomandibular dysfunction

- ACCO appliance
- Archwire
- Activator appliance
- Braces
- Damon system
- Elastics
- Frankel appliance
- Invisalign
- Lingual arch
- Lip bumper
- Herbst Appliance
- List of orthodontic functional appliances

Appliances

- List of palatal expanders
- Lingual braces
- Headgear
- Orthodontic technology
- Orthodontic spacer
- Palatal lift prosthesis
- Palatal expander
- Quad helix
- Retainer
- SureSmile
- Self-ligating braces
- Splint activator
- Twin Block Appliance

Procedures

- Anchorage (orthodontics)
- Cantilever mechanics
- Fiberotomy
- Interproximal reduction
- Intrusion (orthodontics)
- Molar distalization
- SARPE
- Serial extraction

Materials

- Beta-titanium
- Nickel titanium
- Stainless steel
- TiMolium
- Elgiloy
- Ceramic
- Composite
- Dental elastics

- Edward Angle
- Spencer Atkinson
- Clifford Ballard
- Raymond Begg
- Hans Peter Bimler
- Samir Bishara
- Arne Björk
- Charles B. Bolton
- Holly Broadbent Sr.
- Allan G. Brodie
- Charles J. Burstone
- Peter Buschang
- Calvin Case
- Harold Chapman (Orthodontist)
- David Di Biase
- Jean Delaire
- Terry Dischinger
- William B. Downs
- John Nutting Farrar
- Rolf Frankel
- Sheldon Friel
- Thomas M. Graber
- Charles A. Hawley
- Reed Holdaway
- John Hooper (Orthodontist)
- Joseph Jarabak
- Harold Kesling
- Albert Ketcham
- Juri Kurol
- Craven Kurz
- Benno Lischer
- James A. McNamara
- Birte Melsen
- Robert Meyers

**Notable
contributors**

	<ul style="list-style-type: none"> ○ American Association of Orthodontists ○ American Board of Orthodontics ○ British Orthodontic Society
Organizations	<ul style="list-style-type: none"> ○ Canadian Association of Orthodontists ○ Indian Orthodontic Society ○ Italian Academy of Orthodontic Technology ○ Society for Orthodontic Dental Technology (Germany) ○ American Journal of Orthodontics and Dentofacial Orthopedics
Journals	<ul style="list-style-type: none"> ○ The Angle Orthodontist ○ Journal of Orthodontics
Institution	<ul style="list-style-type: none"> ○ Angle School of Orthodontia

	<ul style="list-style-type: none"> ○ Germany ○ United States ○ France
Authority control databases: National	<ul style="list-style-type: none"> ○ BnF data ○ Israel

 [Edit this at Wikidata](#)

Check our other pages :

- [Key Questions Adults Often Ask About Treatment](#)
- [Shared Planning for Improved Patient Experiences](#)
- [Importance of Joint Consultation for Comprehensive Outcomes](#)

IQDENT – Ortodontska Klinika

Phone : +385953817015

City : Zagreb

State : Hrvatska

Zip : 10000

Address : IQDENT – Ortodontska Klinika

Company Website : <https://iqdent.hr/>

USEFUL LINKS

[Orthodontic treatment can help improve your child's smile](#)

[Orthodontic treatment for children](#)

[Sitemap](#)

[Privacy Policy](#)

[About Us](#)

Follow us